



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 14, 2016 – 02:30 PM EDT

PDB ID : 5HN8  
Title : Crystal structure of Plasmodium vivax geranylgeranylpyrophosphate synthase complexed with BPH-1182  
Authors : Liu, Y.-L.; Zhang, Y.; Oldfield, E.  
Deposited on : 2016-01-18  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027939  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027939


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	375	<div> <div>4%</div> <div> <div></div> <div>66%</div> <div>18%</div> <div>•</div> <div>14%</div> </div> </div>
1	B	375	<div> <div>3%</div> <div> <div></div> <div>69%</div> <div>16%</div> <div>•</div> <div>13%</div> </div> </div>
1	C	375	<div> <div>5%</div> <div> <div></div> <div>65%</div> <div>21%</div> <div>•</div> <div>11%</div> </div> </div>
1	D	375	<div> <div>2%</div> <div> <div></div> <div>68%</div> <div>19%</div> <div>•</div> <div>11%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	HXK	A	401	-	-	-	X
2	HXK	B	401	-	-	-	X
2	HXK	C	401	-	-	-	X
2	HXK	D	401	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10771 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Farnesyl pyrophosphate synthase, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	321	Total	C	N	O	S	0	0	0
			2580	1682	405	479	14			
1	B	326	Total	C	N	O	S	0	0	0
			2595	1684	414	483	14			
1	C	334	Total	C	N	O	S	0	0	0
			2662	1730	426	491	15			
1	D	334	Total	C	N	O	S	0	0	0
			2660	1731	424	490	15			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	134	MET	THR	SEE REMARK 999	UNP A5K4U6
A	227	ASP	ASN	SEE REMARK 999	UNP A5K4U6
B	134	MET	THR	SEE REMARK 999	UNP A5K4U6
B	227	ASP	ASN	SEE REMARK 999	UNP A5K4U6
C	134	MET	THR	SEE REMARK 999	UNP A5K4U6
C	227	ASP	ASN	SEE REMARK 999	UNP A5K4U6
D	134	MET	THR	SEE REMARK 999	UNP A5K4U6
D	227	ASP	ASN	SEE REMARK 999	UNP A5K4U6

- Molecule 2 is 2-{[3-hydroxy-5-(octyloxy)benzyl]sulfanyl}benzoic acid (three-letter code: HXK) (formula: C<sub>22</sub>H<sub>28</sub>O<sub>4</sub>S).



- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	60	Total	O	0	0
			60	60		
4	B	19	Total	O	0	0
			19	19		
4	C	38	Total	O	0	0
			38	38		
4	D	44	Total	O	0	0
			44	44		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Farnesyl pyrophosphate synthase, putative




- Molecule 1: Farnesyl pyrophosphate synthase, putative



- Molecule 1: Farnesyl pyrophosphate synthase, putative





- Molecule 1: Farnesyl pyrophosphate synthase, putative



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.07Å 117.22Å 92.79Å 90.00° 116.20° 90.00°	Depositor
Resolution (Å)	31.10 – 2.70 31.10 – 2.70	Depositor EDS
% Data completeness (in resolution range)	92.0 (31.10-2.70) 91.9 (31.10-2.70)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.92 (at 2.72Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.259 , 0.296 0.259 , 0.288	Depositor DCC
$R_{free}$ test set	2075 reflections (5.37%)	DCC
Wilson B-factor (Å <sup>2</sup> )	46.1	Xtriage
Anisotropy	0.144	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 54.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.026 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	10771	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.87% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HXK, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.69	1/2631 (0.0%)	0.69	3/3559 (0.1%)
1	B	0.61	0/2645	0.66	1/3582 (0.0%)
1	C	0.63	1/2712 (0.0%)	0.65	2/3673 (0.1%)
1	D	0.60	0/2712	0.67	1/3675 (0.0%)
All	All	0.63	2/10700 (0.0%)	0.67	7/14489 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	77	CYS	CB-SG	-6.57	1.71	1.82
1	A	114	CYS	CB-SG	-5.39	1.73	1.81

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	376	LEU	CA-CB-CG	5.94	128.97	115.30
1	B	344	VAL	N-CA-C	-5.78	95.38	111.00
1	C	158	LEU	CA-CB-CG	-5.76	102.05	115.30
1	A	158	LEU	CA-CB-CG	-5.74	102.10	115.30
1	A	348	LEU	CA-CB-CG	5.42	127.77	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	343	LYS	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2580	0	2433	60	0
1	B	2595	0	2431	56	0
1	C	2662	0	2528	82	0
1	D	2660	0	2523	82	0
2	A	27	0	0	3	0
2	B	27	0	0	6	0
2	C	27	0	0	2	0
2	D	27	0	0	13	0
3	D	5	0	0	0	0
4	A	60	0	0	11	0
4	B	19	0	0	4	0
4	C	38	0	0	4	0
4	D	44	0	0	5	0
All	All	10771	0	9915	258	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 258 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:401:HXK:CAA	1:C:157:LEU:HD12	1.85	1.06
1:C:283:PHE:CE1	1:C:392:LEU:HG	1.90	1.05
1:D:283:PHE:CD2	2:D:401:HXK:CAH	2.41	1.02
1:A:335:GLY:HA3	4:A:501:HOH:O	1.61	1.01
1:A:291:ASP:OD2	1:A:305:ASP:HB2	1.67	0.95

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/375 (80%)	286 (95%)	15 (5%)	0	100	100
1	B	310/375 (83%)	295 (95%)	13 (4%)	2 (1%)	30	59
1	C	318/375 (85%)	292 (92%)	20 (6%)	6 (2%)	10	25
1	D	320/375 (85%)	300 (94%)	17 (5%)	3 (1%)	21	49
All	All	1249/1500 (83%)	1173 (94%)	65 (5%)	11 (1%)	21	49

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	344	VAL
1	C	291	ASP
1	D	343	LYS
1	D	392	LEU
1	B	291	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/339 (77%)	249 (95%)	12 (5%)	33	64
1	B	258/339 (76%)	244 (95%)	14 (5%)	27	56
1	C	268/339 (79%)	257 (96%)	11 (4%)	37	69
1	D	268/339 (79%)	256 (96%)	12 (4%)	34	65
All	All	1055/1356 (78%)	1006 (95%)	49 (5%)	33	64

5 of 49 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	344	VAL
1	C	172	ARG
1	D	355	ARG
1	B	355	ARG
1	C	185	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	338	ASN
1	C	196	HIS
1	D	337	ASN
1	B	357	HIS
1	B	374	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	HXK	A	401	-	25,28,28	0.94	1 (4%)	31,35,35	1.06	2 (6%)
2	HXK	B	401	-	25,28,28	0.75	0	31,35,35	0.79	0
2	HXK	C	401	-	25,28,28	1.03	1 (4%)	31,35,35	1.36	4 (12%)
2	HXK	D	401	-	25,28,28	0.98	1 (4%)	31,35,35	0.85	0
3	SO4	D	402	-	4,4,4	0.25	0	6,6,6	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HXK	A	401	-	-	0/14/18/18	0/2/2/2
2	HXK	B	401	-	-	0/14/18/18	0/2/2/2
2	HXK	C	401	-	-	0/14/18/18	0/2/2/2
2	HXK	D	401	-	-	0/14/18/18	0/2/2/2
3	SO4	D	402	-	-	0/0/0/0	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	401	HXK	CAZ-SAU	-2.90	1.73	1.77
2	C	401	HXK	CAZ-SAU	-2.21	1.74	1.77
2	A	401	HXK	CAZ-SAU	-2.15	1.74	1.77

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	401	HXK	CAS-SAU-CAZ	-4.09	95.14	102.51
2	A	401	HXK	CAS-SAU-CAZ	-2.62	97.79	102.51
2	C	401	HXK	CAG-CAZ-SAU	-2.09	116.04	121.23
2	A	401	HXK	CAG-CAZ-SAU	-2.08	116.06	121.23
2	C	401	HXK	CAX-CAS-SAU	2.05	118.74	110.81

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	HXK	3	0
2	B	401	HXK	6	0
2	C	401	HXK	2	0
2	D	401	HXK	13	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	321/375 (85%)	0.38	15 (4%) 35 34	21, 49, 70, 80	0
1	B	326/375 (86%)	0.10	11 (3%) 49 49	21, 49, 72, 99	0
1	C	334/375 (89%)	0.32	18 (5%) 29 28	21, 51, 71, 80	0
1	D	334/375 (89%)	0.12	8 (2%) 62 62	21, 50, 71, 80	0
All	All	1315/1500 (87%)	0.23	52 (3%) 42 41	21, 50, 71, 99	0

The worst 5 of 52 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	283	PHE	4.4
1	A	388	LEU	4.1
1	A	345	ILE	4.1
1	C	224	PRO	3.6
1	A	325	PRO	3.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors



of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	HXK	B	401	27/27	0.77	0.29	3.42	31,56,62,63	0
2	HXK	A	401	27/27	0.81	0.28	2.48	34,52,57,59	0
2	HXK	C	401	27/27	0.77	0.31	2.32	47,63,72,72	0
2	HXK	D	401	27/27	0.88	0.24	0.53	38,49,53,54	0
3	SO4	D	402	5/5	0.97	0.09	-3.92	43,43,45,46	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.