



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:33 AM GMT

PDB ID : 2HOR  
Title : Crystal structure of alliinase from garlic- apo form  
Authors : Shimon, L.J.W; Rabinkov, A.; Wilcheck, M.; Mirelman, D.; Frolow, F.  
Deposited on : 2006-07-16  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2475 (1.60-1.60)
Clashscore	102246	2732 (1.60-1.60)
Ramachandran outliers	100387	2654 (1.60-1.60)
Sidechain outliers	100360	2653 (1.60-1.60)
RSRZ outliers	91569	2479 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	427	<div> <div>19%</div> <div>93%</div> <div>6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BMA	A	504	X	-	-	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MAN	A	506	-	-	-	X
3	NAG	A	555	-	-	-	X
4	SO4	A	703	-	-	-	X
7	NO3	A	707	-	X	-	-

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 4122 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alliin lyase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	425	3506	2242	581	657	26	6	15	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	176	ASP	ASN	SEE REMARK 999	UNP Q01594

- Molecule 2 is a polymer of unknown type called SUGAR (6-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	6	71	40	2	29	0	0

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Cl	0	0
			2	2		

- Molecule 7 is NITRATE ION (three-letter code: NO3) (formula: NO<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	N	O	0	0
			4	1	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	496	Total	O	0	0
			496	496		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alliin lyase 1





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.12Å 81.12Å 164.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.60 – 1.60 39.57 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.60-1.60) 99.9 (39.57-1.60)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.47 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.163 , 0.179 0.234 , 0.243	Depositor DCC
$R_{free}$ test set	3663 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	20.4	Xtriage
Anisotropy	0.361	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 55.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 72666 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4122	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, CL, NO3, FUC, ACT, MAN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	2/3640 (0.1%)	0.69	5/4922 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	1	0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	55[A]	GLN	N-CA	-13.55	1.19	1.46
1	A	55[B]	GLN	N-CA	-13.55	1.19	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	55[A]	GLN	CA-C-N	7.45	131.10	116.20
1	A	55[B]	GLN	CA-C-N	7.45	131.10	116.20
1	A	55[A]	GLN	CA-C-O	-7.26	104.85	120.10
1	A	55[B]	GLN	CA-C-O	-7.26	104.85	120.10
1	A	54	ILE	C-N-CA	5.53	135.52	121.70

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	504	BMA	C3

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3506	0	3438	20	0
2	A	71	0	61	0	0
3	A	14	0	13	0	0
4	A	25	0	0	0	0
5	A	4	0	3	0	0
6	A	2	0	0	0	0
7	A	4	0	0	0	0
8	A	496	0	0	5	0
All	All	4122	0	3515	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:354[A]:ARG:NH2	8:A:1172:HOH:O	2.26	0.66
1:A:2[B]:MET:HA	1:A:2[B]:MET:HE3	1.78	0.63
1:A:294:LEU:HG	1:A:298:LYS:HE2	1.82	0.61
1:A:232:HIS:ND1	1:A:318:GLY:HA3	2.18	0.59
1:A:19:ASN:HD22	1:A:21[B]:SER:H	1.51	0.57
1:A:298:LYS:HE3	8:A:899:HOH:O	2.05	0.55
1:A:19:ASN:HD22	1:A:21[A]:SER:H	1.51	0.55
1:A:19:ASN:ND2	1:A:21[B]:SER:H	2.06	0.54
1:A:19:ASN:ND2	1:A:21[A]:SER:H	2.06	0.53
1:A:2[B]:MET:HA	1:A:2[B]:MET:CE	2.39	0.49
1:A:143:LEU:HD22	1:A:221:LYS:HD3	1.97	0.47
1:A:19:ASN:C	1:A:19:ASN:HD22	2.20	0.45
1:A:68:PHE:HB2	1:A:321[A]:LYS:HE2	2.00	0.43
1:A:83:LEU:HD13	1:A:83:LEU:C	2.38	0.43
1:A:74:LYS:NZ	8:A:1138:HOH:O	2.33	0.43
1:A:124:ARG:NH2	1:A:266:LYS:HE2	2.34	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:333:LEU:HD22	1:A:339:PHE:HB2	1.99	0.43
1:A:226[B]:MET:HE1	1:A:240:ALA:HB2	2.02	0.41
1:A:354[B]:ARG:NH1	8:A:946:HOH:O	2.34	0.41
1:A:11[B]:GLU:HG3	8:A:970:HOH:O	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/427 (103%)	426 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	388/375 (104%)	383 (99%)	5 (1%)	76	56

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	LYS
1	A	19	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	25	ARG
1	A	374	LYS
1	A	423	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

6 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	500	1,2	14,14,15	0.87	1 (7%)	15,19,21	0.83	0
2	FUC	A	501	2	10,10,11	0.38	0	14,14,16	1.26	1 (7%)
2	NAG	A	503	2	14,14,15	0.57	0	15,19,21	1.03	1 (6%)
2	BMA	A	504	2	9,10,12	0.68	0	10,13,17	1.03	1 (10%)
2	MAN	A	505	2	11,11,12	0.41	0	14,15,17	0.97	1 (7%)
2	MAN	A	506	2	12,12,12	0.53	0	17,17,17	1.56	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	500	1,2	-	0/6/23/26	0/1/1/1
2	FUC	A	501	2	-	0/0/17/20	0/1/1/1
2	NAG	A	503	2	-	0/6/23/26	0/1/1/1
2	BMA	A	504	2	1/1/3/5	0/2/15/22	0/1/1/1
2	MAN	A	505	2	-	0/2/19/22	0/1/1/1
2	MAN	A	506	2	-	0/2/22/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	NAG	O5-C1	-2.24	1.40	1.43

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	503	NAG	O4-C4-C3	-2.82	103.98	110.34
2	A	504	BMA	C6-C5-C4	-2.04	110.68	113.29
2	A	506	MAN	O5-C5-C6	2.39	112.39	106.36
2	A	506	MAN	C3-C4-C5	2.45	114.47	110.20
2	A	501	FUC	O5-C5-C6	2.56	110.36	106.13
2	A	506	MAN	C4-C3-C2	2.60	115.64	110.79
2	A	505	MAN	C1-O5-C5	2.87	115.89	112.25
2	A	506	MAN	C1-C2-C3	3.76	116.02	110.43

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	504	BMA	C3

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	555	1	14,14,15	0.51	0	15,19,21	1.38	1 (6%)
4	SO4	A	600	-	4,4,4	0.27	0	6,6,6	0.72	0
4	SO4	A	601	-	4,4,4	0.44	0	6,6,6	0.55	0
5	ACT	A	602	-	1,3,3	1.86	0	0,3,3	0.00	-
4	SO4	A	701	-	4,4,4	0.11	0	6,6,6	0.09	0
4	SO4	A	702	-	4,4,4	0.32	0	6,6,6	0.16	0
4	SO4	A	703	-	4,4,4	0.43	0	6,6,6	0.29	0
7	NO3	A	707	-	3,3,3	3.25	3 (100%)	3,3,3	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	555	1	-	0/6/23/26	0/1/1/1
4	SO4	A	600	-	-	0/0/0/0	0/0/0/0
4	SO4	A	601	-	-	0/0/0/0	0/0/0/0
5	ACT	A	602	-	-	0/0/0/0	0/0/0/0
4	SO4	A	701	-	-	0/0/0/0	0/0/0/0
4	SO4	A	702	-	-	0/0/0/0	0/0/0/0
4	SO4	A	703	-	-	0/0/0/0	0/0/0/0
7	NO3	A	707	-	-	0/0/0/0	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	707	NO3	O3-N	2.69	1.39	1.25
7	A	707	NO3	O2-N	2.80	1.39	1.25
7	A	707	NO3	O1-N	4.06	1.41	1.24

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	555	NAG	C1-O5-C5	3.79	117.05	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	425/427 (99%)	0.95	81 (19%) 2 1	23, 31, 37, 48	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	425	LYS	8.4
1	A	1	LYS	6.9
1	A	55[A]	GLN	6.0
1	A	191	ASN	5.7
1	A	424	ALA	5.4
1	A	193	SER	5.2
1	A	423	LYS	5.1
1	A	2[A]	MET	4.2
1	A	422	VAL	4.0
1	A	136	ILE	3.9
1	A	130	VAL	3.9
1	A	152	ASP	3.8
1	A	100	PHE	3.8
1	A	190	VAL	3.7
1	A	336	SER	3.5
1	A	19	ASN	3.4
1	A	135	LEU	3.4
1	A	282	THR	3.4
1	A	285	THR	3.4
1	A	192	VAL	3.3
1	A	332	LEU	3.3
1	A	416	TYR	3.3
1	A	335	GLN	3.1
1	A	331	ALA	3.1
1	A	334	ASP	3.1
1	A	286	PRO	3.1
1	A	93	PHE	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	337	ASP	3.1
1	A	133	THR	3.0
1	A	419	LYS	3.0
1	A	227	VAL	2.9
1	A	284	GLY	2.9
1	A	260	PHE	2.8
1	A	187	ALA	2.8
1	A	289	THR	2.8
1	A	122	LYS	2.7
1	A	131	GLY	2.7
1	A	291	LEU	2.7
1	A	22	GLU	2.7
1	A	34	GLU	2.7
1	A	132	VAL	2.6
1	A	370	TRP	2.6
1	A	218	LYS	2.6
1	A	228	TYR	2.6
1	A	128	PHE	2.5
1	A	101	ILE	2.5
1	A	134	GLN	2.5
1	A	92	TYR	2.5
1	A	248	THR	2.5
1	A	127	VAL	2.5
1	A	296	VAL	2.5
1	A	249	MET	2.5
1	A	274	LEU	2.4
1	A	258	SER	2.4
1	A	405	ILE	2.3
1	A	277	TYR	2.3
1	A	52	GLU	2.3
1	A	259	ARG	2.3
1	A	339	PHE	2.2
1	A	63[A]	SER	2.2
1	A	250	SER	2.2
1	A	67	LEU	2.2
1	A	129	GLY	2.2
1	A	287	ARG	2.2
1	A	371	GLU	2.2
1	A	61	VAL	2.2
1	A	105	LEU	2.1
1	A	256	SER	2.1
1	A	123	ASP	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	54	ILE	2.1
1	A	262	TRP	2.1
1	A	398	ARG	2.1
1	A	51	SER	2.1
1	A	62	ALA	2.1
1	A	261	GLY	2.1
1	A	82	VAL	2.0
1	A	278	MET	2.0
1	A	64	GLY	2.0
1	A	372	GLU	2.0
1	A	356	MET	2.0
1	A	126[A]	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	MAN	A	506	12/12	0.44	0.47	6.69	69,76,77,77	0
2	FUC	A	501	10/11	0.83	0.16	0.97	32,34,35,35	0
2	BMA	A	504	10/12	0.55	0.34	-	48,50,51,53	0
2	NAG	A	500	14/15	0.86	0.13	-	30,32,35,36	0
2	MAN	A	505	11/12	0.59	0.40	-	48,50,50,51	0
2	NAG	A	503	14/15	0.84	0.23	-	38,40,44,47	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy

less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	A	555	14/15	0.75	0.21	2.40	37,42,45,46	0
4	SO4	A	703	5/5	0.96	0.18	2.29	31,31,36,39	0
4	SO4	A	702	5/5	0.90	0.18	0.71	32,36,38,40	0
7	NO3	A	707	4/4	0.93	0.11	-0.79	32,33,33,33	4
4	SO4	A	600	5/5	0.98	0.06	-3.03	24,24,27,28	0
5	ACT	A	602	4/4	0.98	0.05	-3.47	19,20,20,21	0
6	CL	A	705	1/1	0.99	0.02	-4.99	18,18,18,18	0
4	SO4	A	601	5/5	1.00	0.04	-6.04	16,17,18,20	0
4	SO4	A	701	5/5	0.79	0.25	-	67,67,68,68	0
6	CL	A	706	1/1	0.94	0.04	-	47,47,47,47	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.