



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 22, 2016 – 07:19 PM EDT

PDB ID : 5HP1  
Title : STRUCTURE OF HIV-1 REVERSE TRANSCRIPTASE In COMPLEX  
WITH A DNA aptamer and FOSCARNET, a Pyrophosphate analog  
Authors : Das, K.; Arnold, E.  
Deposited on : 2016-01-19  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027939  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027939


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	555	<div> <div>16%</div> <div>76%</div> <div>23%</div> <div>.</div> </div>
1	C	555	<div> <div>25%</div> <div>78%</div> <div>21%</div> <div>.</div> </div>
2	B	444	<div> <div>7%</div> <div>83%</div> <div>10%</div> <div>6%</div> </div>
2	D	444	<div> <div>9%</div> <div>75%</div> <div>17%</div> <div>8%</div> </div>
3	E	38	<div> <div>68%</div> <div>24%</div> <div>8%</div> </div>
3	F	38	<div> <div>63%</div> <div>29%</div> <div>8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	GOL	B	2002	-	-	-	X

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 17386 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HIV-1 REVERSE TRANSCRIPTASE P66 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	553	Total	C	N	O	S	0	0	0
			4494	2909	747	831	7			
1	C	553	Total	C	N	O	S	0	0	0
			4495	2910	748	830	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	280	SER	CYS	engineered mutation	UNP P03366
C	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 2 is a protein called HIV-1 REVERSE TRANSCRIPTASE P51 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	416	Total	C	N	O	S	0	0	0
			3430	2234	567	622	7			
2	D	410	Total	C	N	O	S	0	1	0
			3391	2212	559	614	6			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	MET	-	initiating methionine	UNP P03366
B	-14	ALA	-	expression tag	UNP P03366
B	-13	HIS	-	expression tag	UNP P03366
B	-12	HIS	-	expression tag	UNP P03366
B	-11	HIS	-	expression tag	UNP P03366
B	-10	HIS	-	expression tag	UNP P03366
B	-9	HIS	-	expression tag	UNP P03366
B	-8	HIS	-	expression tag	UNP P03366
B	-7	ALA	-	expression tag	UNP P03366
B	-6	LEU	-	expression tag	UNP P03366

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	GLU	-	expression tag	UNP P03366
B	-4	VAL	-	expression tag	UNP P03366
B	-3	LEU	-	expression tag	UNP P03366
B	-2	PHE	-	expression tag	UNP P03366
B	-1	GLN	-	expression tag	UNP P03366
B	0	GLY	-	expression tag	UNP P03366
B	280	SER	CYS	engineered mutation	UNP P03366
D	-15	MET	-	initiating methionine	UNP P03366
D	-14	ALA	-	expression tag	UNP P03366
D	-13	HIS	-	expression tag	UNP P03366
D	-12	HIS	-	expression tag	UNP P03366
D	-11	HIS	-	expression tag	UNP P03366
D	-10	HIS	-	expression tag	UNP P03366
D	-9	HIS	-	expression tag	UNP P03366
D	-8	HIS	-	expression tag	UNP P03366
D	-7	ALA	-	expression tag	UNP P03366
D	-6	LEU	-	expression tag	UNP P03366
D	-5	GLU	-	expression tag	UNP P03366
D	-4	VAL	-	expression tag	UNP P03366
D	-3	LEU	-	expression tag	UNP P03366
D	-2	PHE	-	expression tag	UNP P03366
D	-1	GLN	-	expression tag	UNP P03366
D	0	GLY	-	expression tag	UNP P03366
D	280	SER	CYS	engineered mutation	UNP P03366


- Molecule 3 is a DNA chain called DNA (38-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	35	Total	C	N	O	P	0	0	0
			720	340	130	215	35			
3	E	35	Total	C	N	O	P	0	0	0
			720	340	130	215	35			

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	Mg	0	0
			4	4		
4	C	4	Total	Mg	0	0
			4	4		

- Molecule 5 is PHOSPHONOFORMIC ACID (three-letter code: PPF) (formula: CH<sub>3</sub>O<sub>5</sub>P).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	P	0	0
			7	1	5	1		
5	C	1	Total	C	O	P	0	0
			7	1	5	1		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is SUCROSE (three-letter code: SUC) (formula:  $C_{12}H_{22}O_{11}$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			23	12	11		
7	D	1	Total	C	O	0	0
			23	12	11		

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			6	3	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is 3'-AZIDO-3'-DEOXYTHYMIDINE-5'-MONOPHOSPHATE (three-letter code: ATM) (formula: C<sub>10</sub>H<sub>14</sub>N<sub>5</sub>O<sub>7</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	F	1	Total	C	N	O	P	0	0
			22	10	5	6	1		
9	E	1	Total	C	N	O	P	0	0
			22	10	5	6	1		

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	1	Total	O	0	0
			1	1		
10	C	1	Total	O	0	0
			1	1		




### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: HIV-1 REVERSE TRANSCRIPTASE P66 SUBUNIT





#### • Molecule 1: HIV-1 REVERSE TRANSCRIPTASE P66 SUBUNIT







• Molecule 2: HIV-1 REVERSE TRANSCRIPTASE P51 SUBUNIT



• Molecule 2: HIV-1 REVERSE TRANSCRIPTASE P51 SUBUNIT



• Molecule 3: DNA (38-MER)



• Molecule 3: DNA (38-MER)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.33Å 129.07Å 131.54Å 90.00° 101.00° 90.00°	Depositor
Resolution (Å)	45.64 – 2.90 45.64 – 2.90	Depositor EDS
% Data completeness (in resolution range)	96.7 (45.64-2.90) 96.5 (45.64-2.90)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.87 (at 2.91Å)	Xtriage
Refinement program	PHENIX (1.10 _2155: ???)	Depositor
R, $R_{free}$	0.246 , 0.273 0.246 , 0.270	Depositor DCC
$R_{free}$ test set	2552 reflections (4.02%)	DCC
Wilson B-factor (Å <sup>2</sup> )	70.7	Xtriage
Anisotropy	0.413	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 65.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	17386	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	104.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OMC, GOL, MG, SUC, PPF, ATM, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/4612	0.50	0/6268
1	C	0.27	0/4613	0.54	1/6268 (0.0%)
2	B	0.28	0/3528	0.49	0/4791
2	D	0.29	0/3493	0.55	2/4747 (0.0%)
3	E	0.58	0/759	1.03	3/1170 (0.3%)
3	F	0.58	0/759	1.03	3/1170 (0.3%)
All	All	0.31	0/17764	0.59	9/24414 (0.0%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	0	DA	O4'-C4'-C3'	-6.29	101.98	104.50
2	D	245	VAL	C-N-CA	-6.16	106.31	121.70
3	E	33	DG	O4'-C4'-C3'	-6.09	102.06	104.50
2	D	308	GLU	CA-CB-CG	5.96	126.50	113.40
3	F	31	DG	C4'-C3'-C2'	-5.54	98.11	103.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4494	0	4538	90	0
1	C	4495	0	4549	78	1
2	B	3430	0	3460	37	0
2	D	3391	0	3411	48	1
3	E	720	0	396	6	0
3	F	720	0	396	8	0
4	A	4	0	0	0	0
4	C	4	0	0	0	0
5	A	7	0	0	1	0
5	C	7	0	0	1	0
6	A	5	0	0	0	0
6	C	5	0	0	0	0
7	B	23	0	22	3	0
7	D	23	0	22	1	0
8	B	12	0	16	2	0
9	E	22	0	12	1	0
9	F	22	0	12	0	0
10	A	1	0	0	0	0
10	C	1	0	0	0	0
All	All	17386	0	16834	253	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 253 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:206:ARG:NH2	2:B:230:MET:O	2.02	0.94
2:B:70:LYS:HG3	2:B:226:PRO:HD2	1.53	0.90
1:C:134:SER:HG	1:C:139:THR:HG1	1.14	0.86
1:C:122:GLU:HA	1:C:125:ARG:HE	1.41	0.83
1:C:330:GLN:HE22	1:C:340:GLN:HE22	1.24	0.83

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:197:GLN:NE2	2:D:91:GLN:OE1[2_657]	2.06	0.14

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	551/555 (99%)	526 (96%)	24 (4%)	1 (0%)	52	84
1	C	551/555 (99%)	526 (96%)	23 (4%)	2 (0%)	39	74
2	B	412/444 (93%)	399 (97%)	13 (3%)	0	100	100
2	D	407/444 (92%)	396 (97%)	11 (3%)	0	100	100
All	All	1921/1998 (96%)	1847 (96%)	71 (4%)	3 (0%)	52	84

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	313	PRO
1	C	313	PRO
1	C	135	ILE

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	492/495 (99%)	469 (95%)	23 (5%)	32	68
1	C	493/495 (100%)	477 (97%)	16 (3%)	46	81
2	B	376/403 (93%)	373 (99%)	3 (1%)	86	96
2	D	371/403 (92%)	368 (99%)	3 (1%)	86	96
All	All	1732/1796 (96%)	1687 (97%)	45 (3%)	54	85

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	450	THR
2	B	413	GLU
1	C	516	GLU
1	A	516	GLU
1	C	7	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	340	GLN
1	C	545	ASN
1	C	428	GLN
1	C	151	GLN
1	C	509	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	OMC	E	2	3	15,22,23	3.12	6 (40%)	20,31,34	1.55	1 (5%)
3	OMC	E	4	3	15,22,23	3.05	6 (40%)	20,31,34	1.61	1 (5%)
3	OMC	F	2	3	15,22,23	3.08	6 (40%)	20,31,34	1.63	1 (5%)
3	OMC	F	4	3	15,22,23	3.10	6 (40%)	20,31,34	1.68	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OMC	E	2	3	-	0/5/27/28	0/2/2/2
3	OMC	E	4	3	-	0/5/27/28	0/2/2/2
3	OMC	F	2	3	-	0/5/27/28	0/2/2/2
3	OMC	F	4	3	-	0/5/27/28	0/2/2/2

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	4	OMC	C4-N4	2.30	1.41	1.35
3	E	2	OMC	C4-N4	2.31	1.41	1.35
3	F	4	OMC	C4-N4	2.32	1.41	1.35
3	F	2	OMC	C4-N4	2.35	1.41	1.35
3	E	4	OMC	C5-C4	2.56	1.47	1.41

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	2	OMC	C6-C5-C4	5.85	119.73	117.44
3	F	2	OMC	C6-C5-C4	6.36	119.93	117.44
3	E	4	OMC	C6-C5-C4	6.42	119.95	117.44
3	F	4	OMC	C6-C5-C4	6.79	120.10	117.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	4	OMC	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 8 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul



statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	PPF	A	602	4	3,6,6	1.17	0	5,9,9	0.66	0
6	SO4	A	606	-	4,4,4	0.23	0	6,6,6	0.08	0
7	SUC	B	2001	-	24,24,24	0.42	0	36,36,36	0.64	0
8	GOL	B	2002	-	5,5,5	0.35	0	5,5,5	0.16	0
8	GOL	B	2003	-	5,5,5	0.38	0	5,5,5	0.30	0
5	PPF	C	601	4	3,6,6	1.06	0	5,9,9	0.79	0
6	SO4	C	605	-	4,4,4	0.21	0	6,6,6	0.06	0
7	SUC	D	501	-	24,24,24	0.43	0	36,36,36	0.66	0
9	ATM	E	101	3,4	13,23,24	1.35	1 (7%)	17,32,35	2.22	3 (17%)
9	ATM	F	101	3,4	13,23,24	1.32	1 (7%)	17,32,35	2.36	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PPF	A	602	4	-	0/0/6/6	0/0/0/0
6	SO4	A	606	-	-	0/0/0/0	0/0/0/0
7	SUC	B	2001	-	-	0/12/51/51	0/2/2/2
8	GOL	B	2002	-	-	0/4/4/4	0/0/0/0
8	GOL	B	2003	-	-	0/4/4/4	0/0/0/0
5	PPF	C	601	4	-	0/0/6/6	0/0/0/0
6	SO4	C	605	-	-	0/0/0/0	0/0/0/0
7	SUC	D	501	-	-	0/12/51/51	0/2/2/2
9	ATM	E	101	3,4	-	0/6/24/25	0/2/2/2
9	ATM	F	101	3,4	-	0/6/24/25	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	F	101	ATM	C4-N3	3.99	1.40	1.33
9	E	101	ATM	C4-N3	4.32	1.40	1.33

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	F	101	ATM	C5-C4-N3	-6.12	120.21	125.35
9	E	101	ATM	C5-C4-N3	-6.07	120.25	125.35
9	E	101	ATM	C2'-C1'-N1	-2.47	108.06	114.14
9	F	101	ATM	C3'-N3'-N4'	2.11	121.86	115.83
9	F	101	ATM	O4'-C1'-N1	3.25	113.40	107.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	602	PPF	1	0
7	B	2001	SUC	3	0
8	B	2003	GOL	2	0
5	C	601	PPF	1	0
7	D	501	SUC	1	0
9	E	101	ATM	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	553/555 (99%)	0.87	90 (16%) <b>2</b> <b>1</b>	35, 103, 171, 223	0
1	C	553/555 (99%)	1.34	141 (25%) <b>1</b> <b>0</b>	35, 121, 215, 263	0
2	B	416/444 (93%)	0.49	33 (7%) <b>15</b> <b>10</b>	42, 82, 149, 195	0
2	D	410/444 (92%)	0.74	42 (10%) <b>9</b> <b>5</b>	37, 98, 167, 238	0
3	E	33/38 (86%)	0.37	0 <b>100</b> <b>100</b>	80, 118, 143, 210	0
3	F	33/38 (86%)	-0.07	0 <b>100</b> <b>100</b>	71, 103, 134, 175	0
All	All	1998/2074 (96%)	0.87	306 (15%) <b>3</b> <b>1</b>	35, 99, 191, 263	0

The worst 5 of 306 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	358	ARG	14.9
1	A	73	LYS	13.3
1	C	131	THR	11.2
1	C	26	LEU	11.0
1	A	71	TRP	9.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	OMC	F	4	21/22	0.94	0.20	-	64,73,78,83	0
3	OMC	E	4	21/22	0.89	0.29	-	86,101,112,115	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	OMC	E	2	21/22	0.86	0.26	-	104,117,125,132	0
3	OMC	F	2	21/22	0.95	0.18	-	88,97,105,106	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
8	GOL	B	2002	6/6	0.84	0.27	4.41	71,79,80,81	0
4	MG	C	602	1/1	0.64	0.17	1.03	62,62,62,62	0
8	GOL	B	2003	6/6	0.92	0.26	0.96	60,64,68,73	0
9	ATM	F	101	22/23	0.88	0.30	-0.21	141,148,164,171	0
4	MG	A	604	1/1	0.50	0.18	-0.33	130,130,130,130	0
7	SUC	D	501	23/23	0.90	0.20	-0.34	81,92,98,101	0
9	ATM	E	101	22/23	0.82	0.30	-0.57	121,134,156,168	0
5	PPF	A	602	7/7	0.72	0.20	-0.62	124,142,159,169	0
5	PPF	C	601	7/7	0.70	0.16	-1.33	150,162,172,189	0
4	MG	C	604	1/1	0.51	0.09	-1.50	166,166,166,166	0
4	MG	A	601	1/1	0.75	0.13	-2.06	80,80,80,80	0
7	SUC	B	2001	23/23	0.92	0.17	-2.56	64,81,90,94	0
6	SO4	C	605	5/5	0.90	0.11	-2.94	102,108,113,123	0
4	MG	A	605	1/1	0.87	0.16	-	80,80,80,80	0
4	MG	A	603	1/1	0.90	0.14	-	133,133,133,133	1
6	SO4	A	606	5/5	0.87	0.15	-	95,108,116,127	0
4	MG	C	603	1/1	0.92	0.11	-	156,156,156,156	0
4	MG	C	606	1/1	0.85	0.12	-	63,63,63,63	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.