



Full wwPDB NMR Structure Validation Report ⓘ

Apr 27, 2016 – 05:44 AM BST

PDB ID : 5HPD
Title : Solution Structure of TAZ2-p53TAD
Authors : Krois, A.S.; Ferreon, J.C.; Martinez-Yamout, M.A.; Dyson, H.J.; Wright, P.E.
Deposited on : 2016-01-20

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR


The overall completeness of chemical shifts assignment is 82%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	158	 63% 6% 31%

2 Ensemble composition and analysis

This entry contains 20 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1767-A:1853, A:3020-A:3026, A:3041-A:3055 (109)	0.20	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 2, 5, 7, 8, 9, 11, 12, 14, 15, 16, 17, 18, 19, 20
2	4, 10
Single-model clusters	3; 6; 13

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2431 atoms, of which 1200 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called CREB-binding protein, Cellular tumor antigen p53 fusion protein.

Mol	Chain	Residues	Atoms						Trace
1	A	158	Total	C	H	N	O	S	0
			2428	753	1200	224	235	16	

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2001	GLY	-	linker	UNP P45481
A	2002	SER	-	linker	UNP P45481
A	2003	GLY	-	linker	UNP P45481
A	2004	SER	-	linker	UNP P45481
A	2005	GLY	-	linker	UNP P45481
A	2006	SER	-	linker	UNP P45481

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
2	A	3	Total	Zn
			3	3

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein



4.2.2 Score per residue for model 2


- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein






4.2.3 Score per residue for model 3

- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein




4.2.4 Score per residue for model 4

- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein




4.2.5 Score per residue for model 5

- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein




4.2.6 Score per residue for model 6

- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein




4.2.7 Score per residue for model 7

- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein




4.2.8 Score per residue for model 8 (medoid)


- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein



4.2.9 Score per residue for model 9


- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein






4.2.10 Score per residue for model 10

- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein




4.2.11 Score per residue for model 11

- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein



4.2.12 Score per residue for model 12

- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein




4.2.13 Score per residue for model 13

- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein

- Molecule 1: CREB-binding protein, Cellular tumor antigen p53 fusion protein


- Molecule 1: CREB-binding protein, Cellular tumor antigen p53 fusion protein

- Molecule 1: CREB-binding protein, Cellular tumor antigen p53 fusion protein




4.2.17 Score per residue for model 17

- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein




4.2.18 Score per residue for model 18

- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein




4.2.19 Score per residue for model 19

- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein



4.2.20 Score per residue for model 20

- Molecule 1: CREB-binding protein,Cellular tumor antigen p53 fusion protein



L3043	S3046	T3055	E3056	D3057	P3058	G3059	P3060	D3061
-------	-------	-------	-------	-------	-------	-------	-------	-------

5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing, molecular dynamics*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
AMBER	refinement	12

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	5hpd_cs.str
Number of chemical shift lists	1
Total number of shifts	1735
Number of shifts mapped to atoms	1735
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	82%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.62±0.00	0±0/888 (0.0±0.0%)	1.06±0.02	5±1/1195 (0.5±0.1%)
All	All	0.62	0/17760 (0.0%)	1.06	109/23900 (0.5%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	1770	ARG	NE-CZ-NH2	7.60	124.10	120.30	6	17
1	A	1811	ARG	NE-CZ-NH2	7.41	124.01	120.30	16	6
1	A	1769	ARG	NE-CZ-NH2	7.25	123.93	120.30	4	18
1	A	1775	ARG	NE-CZ-NH2	6.89	123.74	120.30	5	11
1	A	1787	ARG	NE-CZ-NH2	6.78	123.69	120.30	11	18
1	A	1801	ARG	NE-CZ-NH2	6.68	123.64	120.30	13	9
1	A	1770	ARG	NE-CZ-NH1	6.63	123.61	120.30	20	2
1	A	1811	ARG	NE-CZ-NH1	6.56	123.58	120.30	3	9
1	A	1801	ARG	NE-CZ-NH1	6.52	123.56	120.30	10	1
1	A	1852	ARG	NE-CZ-NH2	6.47	123.54	120.30	9	6
1	A	1769	ARG	NE-CZ-NH1	6.27	123.44	120.30	13	1
1	A	1852	ARG	NE-CZ-NH1	5.88	123.24	120.30	5	8
1	A	1775	ARG	NE-CZ-NH1	5.55	123.08	120.30	6	1
1	A	1787	ARG	NE-CZ-NH1	5.53	123.06	120.30	2	1
1	A	3053	TRP	CA-CB-CG	5.28	123.73	113.70	6	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	17480	17680	17640	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	109/158 (69%)	105±1 (96±1%)	4±1 (4±1%)	0±0 (0±0%)	100	100
All	All	2180/3160 (69%)	2099 (96%)	81 (4%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	102/146 (70%)	97±1 (95±1%)	5±1 (5±1%)	38	81
All	All	2040/2920 (70%)	1946 (95%)	94 (5%)	38	81

All 21 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	3023	TRP	20
1	A	3043	LEU	18
1	A	3055	THR	16
1	A	1837	ASN	10
1	A	1801	ARG	7
1	A	1835	GLN	3
1	A	3046	SER	3
1	A	1806	THR	2
1	A	1852	ARG	2
1	A	3042	ASP	2
1	A	1775	ARG	1
1	A	1811	ARG	1
1	A	3041	ASP	1
1	A	3022	LEU	1
1	A	3021	ASP	1
1	A	1812	LYS	1
1	A	1839	CYS	1
1	A	1836	GLU	1
1	A	1838	LYS	1
1	A	1810	LYS	1
1	A	3024	LYS	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 82% for the well-defined parts and 82% for the entire structure.

7.1 Chemical shift list 1

File name: 5hpd_cs.str

Chemical shift list name: *taz2p53_corrected_assignments.str*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1735
Number of shifts mapped to atoms	1735
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	152	-0.26 ± 0.08	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	145	0.28 ± 0.09	None needed (< 0.5 ppm)
$^{13}\text{C}'$	0	—	—
^{15}N	130	-0.50 ± 0.23	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 82%, i.e. 1152 atoms were assigned a chemical shift out of a possible 1411. 0 out of 17 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	401/535 (75%)	202/213 (95%)	106/218 (49%)	93/104 (89%)
Sidechain	676/789 (86%)	428/471 (91%)	235/272 (86%)	13/46 (28%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	75/87 (86%)	41/46 (89%)	32/32 (100%)	2/9 (22%)
Overall	1152/1411 (82%)	671/730 (92%)	373/522 (71%)	108/159 (68%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 82%, i.e. 1605 atoms were assigned a chemical shift out of a possible 1946. 0 out of 22 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	569/760 (75%)	287/301 (95%)	152/316 (48%)	130/143 (91%)
Sidechain	952/1090 (87%)	607/654 (93%)	324/382 (85%)	21/54 (39%)
Aromatic	84/96 (88%)	46/51 (90%)	36/36 (100%)	2/9 (22%)
Overall	1605/1946 (82%)	940/1006 (93%)	512/734 (70%)	153/206 (74%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

