



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:15 AM GMT

PDB ID : 3HRH
Title : Crystal Structure of Antigen 85C and Glycerol
Authors : Boucau, J.; Sanki, A.K; Umesiri, F.E.; Sucheck, S.J.; Ronning, D.R.
Deposited on : 2009-06-09
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	 2% 77% 12% • 9%
1	B	303	 82% 8% • 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	874	-	-	-	X
2	GOL	A	878	-	-	-	X
2	GOL	B	865	-	-	-	X
2	GOL	B	866	-	-	-	X
2	GOL	B	871	-	-	-	X
2	GOL	B	872	-	-	-	X
2	GOL	B	873	-	-	-	X
2	GOL	B	876	-	-	-	X
2	GOL	B	880	-	-	X	X
2	GOL	B	882	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4527 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antigen 85-C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	276	Total	C	N	O	S	0	0	0
			2148	1368	363	408	9			
1	B	277	Total	C	N	O	S	0	0	0
			2156	1374	364	409	9			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	EXPRESSION TAG	UNP P0A4V4
A	295	LEU	-	EXPRESSION TAG	UNP P0A4V4
A	296	GLU	-	EXPRESSION TAG	UNP P0A4V4
A	297	HIS	-	EXPRESSION TAG	UNP P0A4V4
A	298	HIS	-	EXPRESSION TAG	UNP P0A4V4
A	299	HIS	-	EXPRESSION TAG	UNP P0A4V4
A	300	HIS	-	EXPRESSION TAG	UNP P0A4V4
A	301	HIS	-	EXPRESSION TAG	UNP P0A4V4
A	302	HIS	-	EXPRESSION TAG	UNP P0A4V4
B	500	MET	-	EXPRESSION TAG	UNP P0A4V4
B	795	LEU	-	EXPRESSION TAG	UNP P0A4V4
B	796	GLU	-	EXPRESSION TAG	UNP P0A4V4
B	797	HIS	-	EXPRESSION TAG	UNP P0A4V4
B	798	HIS	-	EXPRESSION TAG	UNP P0A4V4
B	799	HIS	-	EXPRESSION TAG	UNP P0A4V4
B	800	HIS	-	EXPRESSION TAG	UNP P0A4V4
B	801	HIS	-	EXPRESSION TAG	UNP P0A4V4
B	802	HIS	-	EXPRESSION TAG	UNP P0A4V4

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	42	Total 42	O 42	0	0
3	B	61	Total 61	O 61	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Antigen 85-C



• Molecule 1: Antigen 85-C



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.78 Å 80.28 Å 136.84 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.44 – 2.30 48.44 – 2.20	Depositor EDS
% Data completeness (in resolution range)	95.5 (48.44-2.30) 94.4 (48.44-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.69 (at 2.20 Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.190 , 0.221 0.189 , 0.221	Depositor DCC
R_{free} test set	1634 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	24.1	Xtriage
Anisotropy	0.288	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 42.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 38584 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4527	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/2219	0.55	0/3034
1	B	0.35	0/2227	0.57	0/3046
All	All	0.34	0/4446	0.56	0/6080

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2148	0	2004	30	0
1	B	2156	0	2014	20	0
2	A	24	0	32	1	0
2	B	96	0	128	9	0
3	A	42	0	0	0	0
3	B	61	0	0	0	0
All	All	4527	0	4178	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:LYS:HE3	1:B:668:ASP:OD1	1.73	0.89
1:A:233:ARG:HB3	1:A:233:ARG:HH11	1.41	0.85
1:A:222:ILE:HG23	1:B:668:ASP:OD2	1.90	0.72
1:A:233:ARG:HB3	1:A:233:ARG:NH1	2.11	0.65
1:B:725:LYS:NZ	2:B:880:GOL:H11	2.11	0.65
1:A:222:ILE:H	2:B:879:GOL:H31	1.62	0.65
1:B:752:PHE:HB2	2:B:876:GOL:H32	1.79	0.65
1:A:227:LEU:HD23	1:A:260:HIS:CD2	2.33	0.63
1:B:507:PRO:HA	2:B:868:GOL:H2	1.84	0.59
1:B:721:ASN:O	1:B:725:LYS:HG3	2.03	0.59
1:B:722:ILE:HB	1:B:723:PRO:HD3	1.84	0.59
1:B:725:LYS:HZ2	2:B:880:GOL:H11	1.70	0.56
1:A:25:GLN:HB2	1:A:68:MET:HB2	1.86	0.56
1:A:10:TYR:OH	1:A:23:LYS:HD3	2.05	0.56
1:A:233:ARG:HH11	1:A:233:ARG:CB	2.16	0.55
1:B:704:ARG:HG3	1:B:779:VAL:HG21	1.90	0.54
1:B:758:GLY:HA2	1:B:764:TYR:CE1	2.47	0.49
1:A:258:GLY:HA2	1:A:264:TYR:CE2	2.48	0.49
1:A:100:THR:O	1:A:101:ARG:HD2	2.11	0.49
1:A:23:LYS:HB2	1:A:70:VAL:HB	1.98	0.46
1:A:116:THR:HA	1:A:140:GLN:HA	1.98	0.46
1:A:214:PRO:HD3	1:A:225:LYS:HD3	1.97	0.46
1:A:173:ASN:ND2	1:A:175:ASN:H	2.14	0.46
1:B:725:LYS:HZ1	2:B:880:GOL:H11	1.82	0.45
1:B:549:TRP:CZ3	1:B:769:LEU:HD22	2.52	0.45
1:A:222:ILE:HD12	1:B:541:ARG:HD2	1.98	0.44
1:A:152:ASN:N	1:A:153:PRO:CD	2.80	0.44
1:A:167:ASN:O	2:B:880:GOL:H12	2.17	0.44
1:A:119:ALA:HB2	1:A:280:LEU:HD11	2.00	0.44
1:B:585:GLN:O	1:B:586:SER:HB3	2.18	0.44
1:A:134:ALA:HB2	1:A:144:ALA:HB2	1.99	0.44
1:A:222:ILE:HB	2:B:879:GOL:H31	2.01	0.43
1:A:90:ASN:C	1:A:90:ASN:HD22	2.21	0.43
1:B:708:TYR:CG	1:B:709:CYS:N	2.86	0.43
1:B:628:GLY:HA3	2:B:871:GOL:H2	2.00	0.43
1:B:614:SER:HA	1:B:615:PRO:HD3	1.79	0.43
1:A:92:THR:CG2	1:A:94:LYS:HE2	2.49	0.43
1:A:78:THR:HB	2:A:874:GOL:H31	2.01	0.42
1:B:582:GLN:HB2	1:B:583:PRO:CD	2.49	0.42
1:A:97:THR:O	1:A:101:ARG:HB2	2.20	0.42
1:B:623:LEU:C	1:B:623:LEU:HD22	2.41	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:567:ILE:HG21	1:B:603:MET:CE	2.50	0.41
1:A:278:HIS:O	1:A:282:GLY:HA2	2.21	0.41
1:A:85:GLN:O	1:A:86:SER:HB3	2.21	0.40
1:A:173:ASN:HD22	1:A:174:ALA:N	2.19	0.40
1:A:188:ARG:HD3	1:A:188:ARG:O	2.20	0.40
1:A:49:TRP:CZ3	1:A:269:LEU:HD22	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/303 (90%)	262 (96%)	10 (4%)	2 (1%)	26	31
1	B	275/303 (91%)	262 (95%)	12 (4%)	1 (0%)	39	48
All	All	549/606 (91%)	524 (95%)	22 (4%)	3 (0%)	34	41

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	86	SER
1	B	586	SER
1	A	29	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/242 (92%)	215 (96%)	8 (4%)	42	57
1	B	224/242 (93%)	220 (98%)	4 (2%)	66	82
All	All	447/484 (92%)	435 (97%)	12 (3%)	52	70

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90	ASN
1	A	101	ARG
1	A	123	LEU
1	A	152	ASN
1	A	173	ASN
1	A	204	ARG
1	A	216	ASP
1	A	233	ARG
1	B	590	ASN
1	B	623	LEU
1	B	652	ASN
1	B	704	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	GLN
1	A	32	HIS
1	A	73	GLN
1	A	90	ASN
1	A	152	ASN
1	A	173	ASN
1	A	175	ASN
1	A	281	ASN
1	B	512	GLN
1	B	527	GLN
1	B	573	GLN
1	B	590	ASN
1	B	652	ASN
1	B	781	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	862	-	5,5,5	0.55	0	5,5,5	1.12	0
2	GOL	A	874	-	5,5,5	0.62	0	5,5,5	1.01	0
2	GOL	A	875	-	5,5,5	0.66	0	5,5,5	1.02	0
2	GOL	A	878	-	5,5,5	0.57	0	5,5,5	1.01	0
2	GOL	B	861	-	5,5,5	0.60	0	5,5,5	1.09	0
2	GOL	B	863	-	5,5,5	0.46	0	5,5,5	0.99	0
2	GOL	B	865	-	5,5,5	0.56	0	5,5,5	1.04	0
2	GOL	B	866	-	5,5,5	0.68	0	5,5,5	1.05	0
2	GOL	B	868	-	5,5,5	0.61	0	5,5,5	1.05	0
2	GOL	B	869	-	5,5,5	0.62	0	5,5,5	1.05	0
2	GOL	B	870	-	5,5,5	0.78	0	5,5,5	1.29	0
2	GOL	B	871	-	5,5,5	0.54	0	5,5,5	1.00	0
2	GOL	B	872	-	5,5,5	0.52	0	5,5,5	1.02	0
2	GOL	B	873	-	5,5,5	0.61	0	5,5,5	1.07	0
2	GOL	B	876	-	5,5,5	0.57	0	5,5,5	1.08	0
2	GOL	B	877	-	5,5,5	0.63	0	5,5,5	1.05	0
2	GOL	B	879	-	5,5,5	0.47	0	5,5,5	1.08	0
2	GOL	B	880	-	5,5,5	0.58	0	5,5,5	1.02	0
2	GOL	B	881	-	5,5,5	0.55	0	5,5,5	1.20	0
2	GOL	B	882	-	5,5,5	0.62	0	5,5,5	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	862	-	-	0/4/4/4	0/0/0/0
2	GOL	A	874	-	-	0/4/4/4	0/0/0/0
2	GOL	A	875	-	-	0/4/4/4	0/0/0/0
2	GOL	A	878	-	-	0/4/4/4	0/0/0/0
2	GOL	B	861	-	-	0/4/4/4	0/0/0/0
2	GOL	B	863	-	-	0/4/4/4	0/0/0/0
2	GOL	B	865	-	-	0/4/4/4	0/0/0/0
2	GOL	B	866	-	-	0/4/4/4	0/0/0/0
2	GOL	B	868	-	-	0/4/4/4	0/0/0/0
2	GOL	B	869	-	-	0/4/4/4	0/0/0/0
2	GOL	B	870	-	-	0/4/4/4	0/0/0/0
2	GOL	B	871	-	-	0/4/4/4	0/0/0/0
2	GOL	B	872	-	-	0/4/4/4	0/0/0/0
2	GOL	B	873	-	-	0/4/4/4	0/0/0/0
2	GOL	B	876	-	-	0/4/4/4	0/0/0/0
2	GOL	B	877	-	-	0/4/4/4	0/0/0/0
2	GOL	B	879	-	-	0/4/4/4	0/0/0/0
2	GOL	B	880	-	-	0/4/4/4	0/0/0/0
2	GOL	B	881	-	-	0/4/4/4	0/0/0/0
2	GOL	B	882	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	874	GOL	1	0
2	B	868	GOL	1	0
2	B	871	GOL	1	0
2	B	876	GOL	1	0
2	B	879	GOL	2	0
2	B	880	GOL	4	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/303 (91%)	-0.01	5 (1%) 71 78	18, 29, 44, 53	0
1	B	277/303 (91%)	-0.22	1 (0%) 93 95	12, 20, 34, 50	0
All	All	553/606 (91%)	-0.11	6 (1%) 82 86	12, 25, 39, 53	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	782	GLY	3.5
1	A	86	SER	2.7
1	A	90	ASN	2.6
1	A	101	ARG	2.3
1	A	60	TYR	2.1
1	A	85	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	GOL	B	872	6/6	0.89	0.22	10.91	30,33,35,39	0
2	GOL	B	871	6/6	0.81	0.30	10.19	47,49,51,52	0
2	GOL	B	882	6/6	0.66	0.41	9.17	63,67,68,69	0
2	GOL	B	876	6/6	0.66	0.29	5.04	59,61,61,62	0
2	GOL	B	880	6/6	0.58	0.26	4.69	62,63,63,65	0
2	GOL	B	865	6/6	0.85	0.29	4.55	60,60,61,62	0
2	GOL	B	873	6/6	0.36	0.34	4.54	60,62,63,63	0
2	GOL	A	874	6/6	0.96	0.23	4.35	38,39,40,44	0
2	GOL	B	866	6/6	0.80	0.18	3.98	37,48,50,53	0
2	GOL	A	878	6/6	0.74	0.15	2.14	39,41,43,43	0
2	GOL	B	863	6/6	0.87	0.18	1.99	31,32,35,38	0
2	GOL	A	875	6/6	0.87	0.17	1.81	65,65,66,66	0
2	GOL	B	861	6/6	0.89	0.16	1.07	29,33,37,41	0
2	GOL	B	869	6/6	0.91	0.19	0.85	56,57,59,59	0
2	GOL	A	862	6/6	0.93	0.13	0.72	35,36,38,42	0
2	GOL	B	868	6/6	0.89	0.17	0.57	52,54,54,55	0
2	GOL	B	881	6/6	0.79	0.18	-	62,64,64,65	0
2	GOL	B	870	6/6	-0.32	1.42	-	85,86,86,86	0
2	GOL	B	877	6/6	0.76	0.27	-	64,65,65,67	0
2	GOL	B	879	6/6	0.66	0.34	-	68,69,70,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.