



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 13, 2017 – 07:45 PM EST

PDB ID : 5HR1
Title : Crystal structure of thioredoxin L107A mutant
Authors : Noguera, M.E.; Vazquez, D.S.; Howard, E.I.; Cousido-Siah, A.; Mitschler, A.; Podjarny, A.; Santos, J.
Deposited on : 2016-01-22
Resolution : 2.14 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1693 (2.16-2.12)
Clashscore	102246	1824 (2.16-2.12)
Ramachandran outliers	100387	1798 (2.16-2.12)
Sidechain outliers	100360	1798 (2.16-2.12)
RSRZ outliers	91569	1699 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	109	<div> <div style="width: 95%;"></div> <div>95%</div> </div>
1	B	109	<div> <div style="width: 93%;"></div> <div>93%</div> </div>
1	C	109	<div> <div style="width: 94%;"></div> <div>94%</div> </div>
1	D	109	<div> <div style="width: 92%;"></div> <div>92%</div> </div>
1	E	109	<div> <div style="width: 94%;"></div> <div>94%</div> </div>
1	F	109	<div> <div style="width: 90%;"></div> <div>90%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	109	 A horizontal bar chart showing the quality of chain G. The bar is divided into three segments: a red segment on the left labeled '48%', a green segment in the middle labeled '68%', and a grey segment on the right labeled '30%'. A small yellow dot is located at the end of the green segment, and a small black dot is located at the end of the grey segment.

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11028 atoms, of which 5487 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thioredoxin-1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	107	Total	C	H	N	O	S	0	0	0
			1631	522	818	131	157	3			
1	B	107	Total	C	H	N	O	S	0	0	0
			1631	522	818	131	157	3			
1	C	106	Total	C	H	N	O	S	0	0	0
			1619	519	812	130	155	3			
1	D	107	Total	C	H	N	O	S	0	0	0
			1631	522	818	131	157	3			
1	E	106	Total	C	H	N	O	S	0	0	0
			1619	519	812	130	155	3			
1	F	106	Total	C	H	N	O	S	0	0	0
			1619	519	812	130	155	3			
1	G	76	Total	C	H	N	O	S	0	0	0
			1176	376	597	95	105	3			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	107	ALA	LEU	engineered mutation	UNP P0AA27
B	107	ALA	LEU	engineered mutation	UNP P0AA27
C	107	ALA	LEU	engineered mutation	UNP P0AA27
D	107	ALA	LEU	engineered mutation	UNP P0AA27
E	107	ALA	LEU	engineered mutation	UNP P0AA27
F	107	ALA	LEU	engineered mutation	UNP P0AA27
G	107	ALA	LEU	engineered mutation	UNP P0AA27

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Cu	0	0
			1	1		
2	A	1	Total	Cu	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	37	Total 37	O 37	0	0
3	B	37	Total 37	O 37	0	0
3	C	10	Total 10	O 10	0	0
3	D	3	Total 3	O 3	0	0
3	E	12	Total 12	O 12	0	0
3	G	1	Total 1	O 1	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Thioredoxin-1



- Molecule 1: Thioredoxin-1




- Molecule 1: Thioredoxin-1



- Molecule 1: Thioredoxin-1



- Molecule 1: Thioredoxin-1




- Molecule 1: Thioredoxin-1





● Molecule 1: Thioredoxin-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.41Å 88.69Å 115.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.48 – 2.14 47.83 – 2.14	Depositor EDS
% Data completeness (in resolution range)	99.0 (38.48-2.14) 98.8 (47.83-2.14)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.71 (at 2.14Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.206 , 0.235 0.204 , 0.233	Depositor DCC
R_{free} test set	2442 reflections (5.07%)	DCC
Wilson B-factor (Å ²)	37.7	Xtriage
Anisotropy	0.369	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 49.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.020 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11028	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.26	0/828	0.47	0/1123
1	B	0.26	0/828	0.45	0/1123
1	C	0.25	0/822	0.45	0/1115
1	D	0.25	0/828	0.43	0/1123
1	E	0.25	0/822	0.44	0/1115
1	F	0.26	0/822	0.45	0/1115
1	G	0.24	0/591	0.42	0/800
All	All	0.26	0/5541	0.45	0/7514

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	813	818	820	2	0
1	B	813	818	820	4	0
1	C	807	812	812	2	0
1	D	813	818	820	4	0
1	E	807	812	812	2	0
1	F	807	812	812	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	579	597	597	1	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	37	0	0	0	0
3	B	37	0	0	1	0
3	C	10	0	0	0	0
3	D	3	0	0	0	0
3	E	12	0	0	0	0
3	G	1	0	0	0	0
All	All	5541	5487	5493	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 20 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:30:GLU:O	1:F:36:LYS:NZ	2.07	0.86
1:D:82:LYS:NZ	1:D:106:ASN:O	2.21	0.69
1:D:30:GLU:O	1:D:36:LYS:NZ	2.25	0.66
1:E:2:ASP:N	1:E:2:ASP:OD2	2.28	0.66
1:F:2:ASP:OD1	1:F:2:ASP:N	2.41	0.54

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	105/109 (96%)	105 (100%)	0	0	100	100
1	B	105/109 (96%)	105 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	104/109 (95%)	104 (100%)	0	0	100	100
1	D	105/109 (96%)	105 (100%)	0	0	100	100
1	E	104/109 (95%)	103 (99%)	1 (1%)	0	100	100
1	F	104/109 (95%)	104 (100%)	0	0	100	100
1	G	74/109 (68%)	74 (100%)	0	0	100	100
All	All	701/763 (92%)	700 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/87 (99%)	86 (100%)	0	100	100
1	B	86/87 (99%)	86 (100%)	0	100	100
1	C	85/87 (98%)	85 (100%)	0	100	100
1	D	86/87 (99%)	86 (100%)	0	100	100
1	E	85/87 (98%)	84 (99%)	1 (1%)	78	81
1	F	85/87 (98%)	85 (100%)	0	100	100
1	G	60/87 (69%)	60 (100%)	0	100	100
All	All	573/609 (94%)	572 (100%)	1 (0%)	95	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	2	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	107/109 (98%)	0.33	1 (0%) 85 89	24, 37, 72, 94	0
1	B	107/109 (98%)	0.54	1 (0%) 85 89	22, 39, 71, 98	0
1	C	106/109 (97%)	0.52	6 (5%) 27 35	33, 58, 90, 107	0
1	D	107/109 (98%)	0.33	3 (2%) 56 66	41, 57, 89, 110	0
1	E	106/109 (97%)	0.74	8 (7%) 17 22	35, 58, 102, 119	0
1	F	106/109 (97%)	0.59	7 (6%) 22 28	45, 66, 95, 116	0
1	G	76/109 (69%)	4.14	52 (68%) 0 0	53, 112, 149, 171	0
All	All	715/763 (93%)	0.89	78 (10%) 7 11	22, 57, 113, 171	0

The worst 5 of 78 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	81	PHE	13.4
1	G	70	TYR	11.5
1	G	58	LEU	11.1
1	G	69	LYS	10.9
1	G	86	VAL	9.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	CU	A	201	1/1	0.91	0.06	-	73,73,73,73	0
2	CU	B	201	1/1	0.92	0.12	-	79,79,79,79	1

6.5 Other polymers [i](#)

There are no such residues in this entry.