



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 26, 2016 – 05:35 PM EDT

PDB ID : 5HSQ  
Title : The surface engineered photosensory module (PAS-GAF-PHY) of the bacterial phytochrome Agp1 (AtBphP1) in the Pr form, chromophore modelled with an endocyclic double bond in pyrrole ring A.  
Authors : Nagano, S.; Scheerer, P.; Zubow, K.; Lamparter, T.; Krauss, N.  
Deposited on : 2016-01-26  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027939  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027939


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	503	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	603	-	-	-	X
5	IMD	A	604	-	-	X	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4238 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bacteriophytochrome protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	488	3830	2433	666	716	15	0	18	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	86	ALA	GLU	engineered mutation	UNP Q7CY45
A	87	ALA	GLU	engineered mutation	UNP Q7CY45
A	336	ALA	GLU	engineered mutation	UNP Q7CY45
A	337	ALA	LYS	engineered mutation	UNP Q7CY45
A	505	LEU	-	expression tag	UNP Q7CY45
A	506	GLU	-	expression tag	UNP Q7CY45
A	507	HIS	-	expression tag	UNP Q7CY45
A	508	HIS	-	expression tag	UNP Q7CY45
A	509	HIS	-	expression tag	UNP Q7CY45
A	510	HIS	-	expression tag	UNP Q7CY45
A	511	HIS	-	expression tag	UNP Q7CY45
A	512	HIS	-	expression tag	UNP Q7CY45

- Molecule 2 is BILIVERDINE IX ALPHA (three-letter code: BLA) (formula: C<sub>33</sub>H<sub>34</sub>N<sub>4</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			43	33	4	6		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $\text{C}_3\text{H}_8\text{O}_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is IMIDAZOLE (three-letter code: IMD) (formula:  $C_3H_5N_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	N	0	0
			5	3	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	349	Total	O	0	4
			353	353		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bacteriophytochrome protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.35Å 69.35Å 236.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.31 – 1.85 31.75 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.0 (34.31-1.85) 98.0 (31.75-1.85)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.04 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.183 , 0.224 0.191 , 0.230	Depositor DCC
$R_{free}$ test set	2527 reflections (5.38%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.3	Xtriage
Anisotropy	0.387	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 43.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4238	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, BLA, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.00	3/3963 (0.1%)	1.04	14/5381 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	286	SER	CB-OG	-5.89	1.34	1.42
1	A	115	GLU	CD-OE1	5.43	1.31	1.25
1	A	23	GLU	CD-OE1	5.20	1.31	1.25

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	192	ARG	NE-CZ-NH1	11.24	125.92	120.30
1	A	488	ARG	NE-CZ-NH1	10.40	125.50	120.30
1	A	488	ARG	NE-CZ-NH2	-8.90	115.85	120.30
1	A	304[A]	MET	CG-SD-CE	7.47	112.16	100.20
1	A	304[B]	MET	CG-SD-CE	7.47	112.16	100.20
1	A	203	ARG	NE-CZ-NH1	6.84	123.72	120.30
1	A	167	ASP	CB-CG-OD1	6.18	123.86	118.30
1	A	292	ARG	NE-CZ-NH1	5.88	123.24	120.30
1	A	458	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	A	192	ARG	CD-NE-CZ	5.77	131.68	123.60
1	A	448	ASP	CB-CG-OD2	-5.74	113.14	118.30
1	A	170	LEU	CB-CG-CD1	5.48	120.32	111.00
1	A	423	ARG	NE-CZ-NH1	5.36	122.98	120.30
1	A	423	ARG	NE-CZ-NH2	-5.20	117.70	120.30

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3830	0	3843	26	0
2	A	43	0	31	6	0
3	A	1	0	0	0	0
4	A	6	0	8	1	0
5	A	5	0	5	4	0
6	A	353	0	0	1	0
All	All	4238	0	3887	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225[A]:ARG:CG	1:A:225[A]:ARG:HH11	1.73	1.01
2:A:601:BLA:HMC1	2:A:601:BLA:HBC1	1.52	0.90
1:A:408:THR:HG23	6:A:943:HOH:O	1.73	0.85
1:A:225[A]:ARG:HG3	1:A:225[A]:ARG:HH11	1.44	0.81
1:A:225[A]:ARG:HG2	1:A:225[A]:ARG:HH11	1.43	0.80
1:A:339:VAL:HG23	1:A:363:LEU:HD21	1.78	0.65
1:A:139:SER:OG	1:A:140:ALA:N	2.33	0.62
1:A:225[A]:ARG:HG3	1:A:225[A]:ARG:NH1	2.11	0.59
2:A:601:BLA:HMC1	2:A:601:BLA:CBC	2.31	0.56
1:A:213:MET:H	5:A:604:IMD:C5	2.19	0.56
1:A:192:ARG:HG2	1:A:445:TRP:CZ3	2.43	0.54
1:A:183:GLY:HA2	4:A:603:GOL:H12	1.92	0.52
1:A:140:ALA:O	1:A:142:SER:N	2.43	0.52
1:A:225[A]:ARG:HG2	1:A:225[A]:ARG:NH1	2.16	0.52
1:A:192:ARG:HG3	1:A:193:PHE:N	2.26	0.51
1:A:105:ARG:HG2	1:A:105:ARG:HH11	1.76	0.50
1:A:140:ALA:O	1:A:141:PRO:C	2.49	0.49
1:A:300:GLN:O	1:A:304[A]:MET:HG3	2.12	0.49
1:A:135[A]:LYS:HE3	1:A:135[A]:LYS:HB2	1.52	0.47
1:A:212:ARG:HA	5:A:604:IMD:C4	2.46	0.46
1:A:247:SER:OG	1:A:249:VAL:HG12	2.15	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175[A]:HIS:CE1	1:A:192:ARG:HD3	2.53	0.44
2:A:601:BLA:C1B	2:A:601:BLA:HMA1	2.47	0.44
1:A:53:ASN:O	1:A:56:ASN:HB3	2.18	0.44
1:A:339:VAL:HG23	1:A:363:LEU:CD2	2.47	0.42
2:A:601:BLA:NB	2:A:601:BLA:HMA1	2.34	0.41
1:A:242:GLN:O	5:A:604:IMD:C5	2.68	0.41
2:A:601:BLA:HAD1	2:A:601:BLA:HMD1	1.89	0.41
1:A:213:MET:H	5:A:604:IMD:H5	1.85	0.41
1:A:104:HIS:CE1	1:A:111:ILE:HB	2.56	0.41
1:A:55:ALA:O	1:A:59:GLY:N	2.51	0.41
2:A:601:BLA:HAC	2:A:601:BLA:HHD	1.95	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	505/503 (100%)	495 (98%)	9 (2%)	1 (0%)	52 36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	141	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	404/402 (100%)	389 (96%)	15 (4%)	41	20

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	LEU
1	A	105	ARG
1	A	135[A]	LYS
1	A	135[B]	LYS
1	A	135[C]	LYS
1	A	143	LEU
1	A	170	LEU
1	A	192	ARG
1	A	225[A]	ARG
1	A	225[B]	ARG
1	A	227[A]	GLU
1	A	227[B]	GLU
1	A	311	SER
1	A	455	GLU
1	A	499	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	BLA	A	601	1	36,46,46	2.80	14 (38%)	45,67,67	1.82	11 (24%)
4	GOL	A	603	-	5,5,5	0.49	0	5,5,5	1.27	0
5	IMD	A	604	-	3,5,5	0.61	0	4,5,5	1.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BLA	A	601	1	-	0/22/74/74	0/4/4/4
4	GOL	A	603	-	-	0/4/4/4	0/0/0/0
5	IMD	A	604	-	-	0/0/0/0	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	BLA	C4D-C3D	-4.57	1.37	1.45
2	A	601	BLA	C1D-C2D	-3.63	1.37	1.45
2	A	601	BLA	C1B-C2B	-3.21	1.39	1.45
2	A	601	BLA	C1C-C2C	-3.17	1.38	1.47
2	A	601	BLA	CAB-C3B	-3.03	1.38	1.47
2	A	601	BLA	CAD-C3D	-2.99	1.43	1.51
2	A	601	BLA	C3C-C4C	-2.92	1.41	1.45
2	A	601	BLA	C3B-C4B	-2.68	1.39	1.47
2	A	601	BLA	C4B-NB	-2.17	1.33	1.37
2	A	601	BLA	CMB-C2B	2.16	1.55	1.50
2	A	601	BLA	CBB-CAB	3.43	1.47	1.30
2	A	601	BLA	CBC-CAC	4.20	1.50	1.30
2	A	601	BLA	CAA-C2A	4.98	1.60	1.52
2	A	601	BLA	CHA-C4D	10.51	1.44	1.35

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	A	601	BLA	C1B-NB-C4B	-4.15	104.72	110.73
2	A	601	BLA	CHD-C4C-C3C	-2.86	120.49	128.02
2	A	601	BLA	C3D-C4D-ND	-2.75	105.79	109.98
2	A	601	BLA	C2D-C1D-ND	-2.59	105.00	110.65
2	A	601	BLA	CMB-C2B-C3B	-2.04	123.69	128.58
2	A	601	BLA	CMC-C2C-C1C	2.02	126.58	121.40
2	A	601	BLA	C4D-ND-C1D	2.61	111.47	106.50
2	A	601	BLA	CMB-C2B-C1B	2.75	127.79	124.20
2	A	601	BLA	C1D-C2D-C3D	3.14	110.06	106.43
2	A	601	BLA	CAB-C3B-C4B	3.14	132.63	123.58
2	A	601	BLA	CHD-C1D-ND	3.94	132.24	124.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	BLA	6	0
4	A	603	GOL	1	0
5	A	604	IMD	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	488/503 (97%)	-0.06	16 (3%)	50 47	24, 34, 61, 96	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	17	LEU	5.1
1	A	141	PRO	3.9
1	A	140	ALA	3.9
1	A	394	ALA	3.6
1	A	61	ASP	3.6
1	A	55	ALA	3.3
1	A	390[A]	GLU	3.0
1	A	392	GLU	3.0
1	A	277	ILE	2.9
1	A	395	ASP	2.7
1	A	86	ALA	2.6
1	A	276	LEU	2.3
1	A	63	PRO	2.2
1	A	59	GLY	2.2
1	A	265	VAL	2.1
1	A	403	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	A	603	6/6	0.77	0.31	17.65	41,56,64,64	0
5	IMD	A	604	5/5	0.89	0.17	1.27	29,38,42,42	0
2	BLA	A	601	43/43	0.94	0.14	0.55	24,27,32,35	0
3	CA	A	602	1/1	0.99	0.03	-	37,37,37,37	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.