



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 2, 2016 – 09:14 PM EDT

PDB ID : 1HT1  
Title : Nucleotide-Dependent Conformational Changes in a Protease-Associated ATPase HslU  
Authors : Wang, J.; Song, J.J.; Seong, I.S.; Franklin, M.C.; Kamtekar, S.; Eom, S.H.; Chung, C.H.  
Deposited on : 2000-12-27  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	175	<div> <div>2%</div> <div> <div></div> <div>54%</div> <div>42%</div> <div>••</div> </div> </div>
1	B	175	<div> <div>7%</div> <div> <div></div> <div>38%</div> <div>54%</div> <div>7%</div> <div>•</div> </div> </div>
1	C	175	<div> <div>8%</div> <div> <div></div> <div>37%</div> <div>53%</div> <div>9%</div> <div>•</div> </div> </div>
1	D	175	<div> <div>4%</div> <div> <div></div> <div>53%</div> <div>39%</div> <div>7%</div> <div>•</div> </div> </div>
1	V	175	<div> <div>7%</div> <div> <div></div> <div>34%</div> <div>56%</div> <div>9%</div> <div>•</div> </div> </div>
1	X	175	<div> <div>2%</div> <div> <div></div> <div>52%</div> <div>40%</div> <div>7%</div> <div>•</div> </div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	Y	175	<div> <div>2%</div> <div>42%</div> <div>49%</div> <div>7%</div> <div>..</div> </div>
1	Z	175	<div> <div>2%</div> <div>51%</div> <div>44%</div> <div></div> <div>..</div> </div>
2	E	449	<div> <div>8%</div> <div>44%</div> <div>40%</div> <div>6%</div> <div>•</div> <div>9%</div> </div>
2	F	449	<div> <div>12%</div> <div>40%</div> <div>41%</div> <div>8%</div> <div>•</div> <div>9%</div> </div>
2	G	449	<div> <div>8%</div> <div>44%</div> <div>39%</div> <div>7%</div> <div>•</div> <div>9%</div> </div>
2	I	449	<div> <div>12%</div> <div>43%</div> <div>39%</div> <div>8%</div> <div>•</div> <div>9%</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 23636 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEAT SHOCK LOCUS HSLV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	174	Total	C	N	O	S	0	0	0
			1328	834	237	253	4			
1	D	174	Total	C	N	O	S	0	0	0
			1328	834	237	253	4			
1	V	174	Total	C	N	O	S	0	0	0
			1328	834	237	253	4			
1	X	174	Total	C	N	O	S	0	0	0
			1328	834	237	253	4			
1	A	174	Total	C	N	O	S	0	0	0
			1328	834	237	253	4			
1	B	174	Total	C	N	O	S	0	0	0
			1328	834	237	253	4			
1	Z	174	Total	C	N	O	S	0	0	0
			1328	834	237	253	4			
1	Y	174	Total	C	N	O	S	0	0	0
			1328	834	237	253	4			

- Molecule 2 is a protein called HEAT SHOCK LOCUS HSLU.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	408	Total	C	N	O	S	0	0	0
			3226	2014	577	625	10			
2	F	408	Total	C	N	O	S	0	0	0
			3226	2014	577	625	10			
2	G	408	Total	C	N	O	S	0	0	0
			3226	2014	577	625	10			
2	I	408	Total	C	N	O	S	0	0	0
			3226	2014	577	625	10			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	-5	HIS	-	EXPRESSION TAG	UNP P0A6H5

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	-4	HIS	-	EXPRESSION TAG	UNP P0A6H5
E	-3	HIS	-	EXPRESSION TAG	UNP P0A6H5
E	-2	HIS	-	EXPRESSION TAG	UNP P0A6H5
E	-1	HIS	-	EXPRESSION TAG	UNP P0A6H5
E	0	HIS	-	EXPRESSION TAG	UNP P0A6H5
E	1	HIS	-	EXPRESSION TAG	UNP P0A6H5
F	-5	HIS	-	EXPRESSION TAG	UNP P0A6H5
F	-4	HIS	-	EXPRESSION TAG	UNP P0A6H5
F	-3	HIS	-	EXPRESSION TAG	UNP P0A6H5
F	-2	HIS	-	EXPRESSION TAG	UNP P0A6H5
F	-1	HIS	-	EXPRESSION TAG	UNP P0A6H5
F	0	HIS	-	EXPRESSION TAG	UNP P0A6H5
F	1	HIS	-	EXPRESSION TAG	UNP P0A6H5
G	-5	HIS	-	EXPRESSION TAG	UNP P0A6H5
G	-4	HIS	-	EXPRESSION TAG	UNP P0A6H5
G	-3	HIS	-	EXPRESSION TAG	UNP P0A6H5
G	-2	HIS	-	EXPRESSION TAG	UNP P0A6H5
G	-1	HIS	-	EXPRESSION TAG	UNP P0A6H5
G	0	HIS	-	EXPRESSION TAG	UNP P0A6H5
G	1	HIS	-	EXPRESSION TAG	UNP P0A6H5
I	-5	HIS	-	EXPRESSION TAG	UNP P0A6H5
I	-4	HIS	-	EXPRESSION TAG	UNP P0A6H5
I	-3	HIS	-	EXPRESSION TAG	UNP P0A6H5
I	-2	HIS	-	EXPRESSION TAG	UNP P0A6H5
I	-1	HIS	-	EXPRESSION TAG	UNP P0A6H5
I	0	HIS	-	EXPRESSION TAG	UNP P0A6H5
I	1	HIS	-	EXPRESSION TAG	UNP P0A6H5

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	E	1	Total 27	C 10	N 5	O 10	P 2	0	0
3	F	1	Total 27	C 10	N 5	O 10	P 2	0	0
3	G	1	Total 27	C 10	N 5	O 10	P 2	0	0
3	I	1	Total 27	C 10	N 5	O 10	P 2	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: HEAT SHOCK LOCUS HSLV



#### • Molecule 1: HEAT SHOCK LOCUS HSLV




#### • Molecule 1: HEAT SHOCK LOCUS HSLV







• Molecule 1: HEAT SHOCK LOCUS HSLV




• Molecule 1: HEAT SHOCK LOCUS HSLV




• Molecule 1: HEAT SHOCK LOCUS HSLV




• Molecule 1: HEAT SHOCK LOCUS HSLV









● Molecule 2: HEAT SHOCK LOCUS HSLU



● Molecule 2: HEAT SHOCK LOCUS HSLU





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	172.02Å 172.02Å 276.57Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.62 – 2.80 29.62 – 2.80	Depositor EDS
% Data completeness (in resolution range)	92.5 (29.62-2.80) 92.5 (29.62-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.49 (at 2.80Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.261 , 0.309 0.256 , 0.305	Depositor DCC
$R_{free}$ test set	10933 reflections (11.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	67.6	Xtriage
Anisotropy	0.378	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 34.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.499 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	23636	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 60.98 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.4024e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.46	0/1345	0.72	0/1817
1	B	0.41	0/1345	0.66	0/1817
1	C	0.44	0/1345	0.66	0/1817
1	D	0.37	0/1345	0.65	0/1817
1	V	0.45	0/1345	0.64	0/1817
1	X	0.36	0/1345	0.64	0/1817
1	Y	0.41	0/1345	0.67	0/1817
1	Z	0.46	0/1345	0.72	0/1817
2	E	0.42	0/3266	0.69	3/4400 (0.1%)
2	F	0.45	1/3266 (0.0%)	0.69	2/4400 (0.0%)
2	G	0.43	1/3266 (0.0%)	0.69	4/4400 (0.1%)
2	I	0.42	0/3266	0.68	2/4400 (0.0%)
All	All	0.43	2/23824 (0.0%)	0.68	11/32136 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	219	LYS	C-N	5.58	1.46	1.34
2	F	152	PRO	CA-C	-5.43	1.42	1.52

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	318	LEU	CA-CB-CG	6.46	130.16	115.30
2	I	220	ASP	CB-CA-C	-6.40	97.59	110.40
2	G	318	LEU	CA-CB-CG	6.35	129.90	115.30
2	G	152	PRO	C-N-CA	-5.99	106.72	121.70
2	F	152	PRO	CA-N-CD	-5.91	103.23	111.50

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1328	0	1348	96	0
1	B	1328	0	1348	126	0
1	C	1328	0	1348	140	0
1	D	1328	0	1348	87	0
1	V	1328	0	1348	143	0
1	X	1328	0	1348	98	0
1	Y	1328	0	1348	121	0
1	Z	1328	0	1348	101	0
2	E	3226	0	3294	260	1
2	F	3226	0	3293	265	1
2	G	3226	0	3294	280	0
2	I	3226	0	3293	303	0
3	E	27	0	12	3	0
3	F	27	0	12	3	0
3	G	27	0	12	3	0
3	I	27	0	12	4	0
All	All	23636	0	24006	1871	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

The worst 5 of 1871 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:152:PRO:O	2:E:154:ALA:CA	1.83	1.25
2:E:152:PRO:C	2:E:154:ALA:H	1.27	1.15
2:E:152:PRO:C	2:E:154:ALA:N	1.80	1.12
1:B:135:ALA:HB1	1:Z:136:LEU:HD13	1.21	1.12
2:F:216:LEU:HG	2:F:221:ALA:HB2	1.25	1.11

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:7:ARG:NH2	2:F:409:ASP:OD2[2_665]	2.06	0.14

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	172/175 (98%)	151 (88%)	17 (10%)	4 (2%)	8	26
1	B	172/175 (98%)	138 (80%)	27 (16%)	7 (4%)	3	11
1	C	172/175 (98%)	140 (81%)	27 (16%)	5 (3%)	6	19
1	D	172/175 (98%)	146 (85%)	23 (13%)	3 (2%)	11	36
1	V	172/175 (98%)	144 (84%)	25 (14%)	3 (2%)	11	36
1	X	172/175 (98%)	148 (86%)	20 (12%)	4 (2%)	8	26
1	Y	172/175 (98%)	136 (79%)	26 (15%)	10 (6%)	2	5
1	Z	172/175 (98%)	150 (87%)	18 (10%)	4 (2%)	8	26
2	E	404/449 (90%)	349 (86%)	46 (11%)	9 (2%)	8	28
2	F	404/449 (90%)	343 (85%)	45 (11%)	16 (4%)	4	12
2	G	404/449 (90%)	344 (85%)	45 (11%)	15 (4%)	4	14
2	I	404/449 (90%)	347 (86%)	43 (11%)	14 (4%)	4	15
All	All	2992/3196 (94%)	2536 (85%)	362 (12%)	94 (3%)	5	17

5 of 94 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	116	GLU
1	V	116	GLU
1	Y	54	PHE
1	Y	68	GLN
1	Y	71	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	136/136 (100%)	129 (95%)	7 (5%)	29	63
1	B	136/136 (100%)	125 (92%)	11 (8%)	15	39
1	C	136/136 (100%)	121 (89%)	15 (11%)	8	23
1	D	136/136 (100%)	124 (91%)	12 (9%)	12	35
1	V	136/136 (100%)	119 (88%)	17 (12%)	6	17
1	X	136/136 (100%)	124 (91%)	12 (9%)	12	35
1	Y	136/136 (100%)	127 (93%)	9 (7%)	21	51
1	Z	136/136 (100%)	128 (94%)	8 (6%)	24	57
2	E	350/383 (91%)	308 (88%)	42 (12%)	6	19
2	F	350/383 (91%)	301 (86%)	49 (14%)	4	13
2	G	350/383 (91%)	307 (88%)	43 (12%)	6	18
2	I	350/383 (91%)	301 (86%)	49 (14%)	4	13
All	All	2488/2620 (95%)	2214 (89%)	274 (11%)	8	23

5 of 274 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	337	THR
2	F	152	PRO
2	I	232	LYS
2	E	355	TYR
2	F	68	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 92 such sidechains are listed below:

Mol	Chain	Res	Type
2	E	348	ASN
2	F	141	ASN
2	I	150	GLN
2	E	382	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	E	428	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ADP	E	450	-	24,29,29	1.51	7 (29%)	23,45,45	1.98	3 (13%)
3	ADP	F	1450	-	24,29,29	1.48	5 (20%)	23,45,45	2.00	3 (13%)
3	ADP	G	2450	-	24,29,29	1.53	7 (29%)	23,45,45	2.05	4 (17%)
3	ADP	I	3450	-	24,29,29	1.46	5 (20%)	23,45,45	2.02	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	E	450	-	-	0/12/32/32	0/3/3/3
3	ADP	F	1450	-	-	0/12/32/32	0/3/3/3
3	ADP	G	2450	-	-	0/12/32/32	0/3/3/3
3	ADP	I	3450	-	-	0/12/32/32	0/3/3/3

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	2450	ADP	C8-N7	-3.88	1.27	1.34
3	E	450	ADP	C8-N7	-3.82	1.27	1.34
3	F	1450	ADP	C8-N7	-3.81	1.27	1.34
3	I	3450	ADP	C8-N7	-3.73	1.27	1.34
3	E	450	ADP	PA-O1A	-2.25	1.43	1.51

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	1450	ADP	N3-C2-N1	-5.09	124.87	128.87
3	G	2450	ADP	N3-C2-N1	-5.03	124.92	128.87
3	I	3450	ADP	N3-C2-N1	-4.90	125.02	128.87
3	E	450	ADP	N3-C2-N1	-4.66	125.21	128.87
3	G	2450	ADP	C2'-C1'-N9	-2.13	107.76	113.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	450	ADP	3	0
3	F	1450	ADP	3	0
3	G	2450	ADP	3	0
3	I	3450	ADP	4	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	174/175 (99%)	0.45	3 (1%) 73 63	30, 54, 89, 98	0
1	B	174/175 (99%)	0.56	12 (6%) 20 11	36, 68, 101, 102	0
1	C	174/175 (99%)	0.58	14 (8%) 15 7	54, 81, 100, 102	0
1	D	174/175 (99%)	0.45	7 (4%) 42 30	49, 71, 97, 100	0
1	V	174/175 (99%)	0.53	12 (6%) 20 11	54, 80, 98, 102	0
1	X	174/175 (99%)	0.39	3 (1%) 73 63	51, 70, 98, 101	0
1	Y	174/175 (99%)	0.50	4 (2%) 64 52	36, 68, 100, 102	0
1	Z	174/175 (99%)	0.44	3 (1%) 73 63	30, 55, 90, 97	0
2	E	408/449 (90%)	0.59	37 (9%) 11 6	32, 59, 102, 102	0
2	F	408/449 (90%)	0.85	54 (13%) 4 2	33, 61, 102, 102	0
2	G	408/449 (90%)	0.62	34 (8%) 14 7	32, 59, 102, 102	0
2	I	408/449 (90%)	0.82	53 (12%) 5 2	32, 60, 102, 102	0
All	All	3024/3196 (94%)	0.61	236 (7%) 16 8	30, 63, 102, 102	0

The worst 5 of 236 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	G	90	GLY	9.0
2	G	144	GLY	8.7
2	F	90	GLY	8.1
2	I	141	ASN	7.5
2	F	141	ASN	7.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ADP	G	2450	27/27	0.96	0.26	0.90	38,46,58,59	0
3	ADP	F	1450	27/27	0.95	0.26	0.85	47,51,56,58	0
3	ADP	E	450	27/27	0.96	0.25	0.65	40,46,58,58	0
3	ADP	I	3450	27/27	0.96	0.23	0.26	45,49,54,55	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.