



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:17 PM GMT

PDB ID : 4HUT
Title : Structure of ATP:co(I)rrinoid adenosyltransferase (CobA) from *Salmonella enterica* in complex with four and five-coordinate cob(II)alamin and ATP
Authors : Moore, T.C.; Newmister, S.A.; Rayment, I.; Escalante-Semerena, J.C.
Deposited on : 2012-11-04
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	191	
1	B	191	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	A	204	-	-	-	X
5	EDO	A	205	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3320 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cob(I)yrinic acid a,c-diamide adenosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	191	Total	C	N	O	S	0	1	0
			1482	931	269	274	8			
1	B	169	Total	C	N	O	S	0	0	0
			1288	815	228	237	8			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 4 is COBALAMIN (three-letter code: B12) (formula: $C_{62}H_{89}CoN_{13}O_{14}P$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total 91	C 62	Co 1	N 13	O 14	P 1	0	0
4	B	1	Total 64	C 45	Co 1	N 11	O 7		0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	186	Total	O	0	0
			186	186		
6	B	137	Total	O	0	0
			137	137		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cob(I)yrinic acid a,c-diamide adenosyltransferase



- Molecule 1: Cob(I)yrinic acid a,c-diamide adenosyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.86 Å 74.22 Å 92.40 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.87 – 1.95 29.85 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.7 (29.87-1.95) 97.8 (29.85-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.12 (at 1.95 Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.175 , 0.232 0.180 , 0.232	Depositor DCC
R_{free} test set	1514 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	29.6	Xtriage
Anisotropy	0.171	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 29962 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3320	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, B12, EDO, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.00	0/1513	1.17	14/2048 (0.7%)
1	B	1.09	0/1314	0.94	2/1782 (0.1%)
All	All	1.04	0/2827	1.07	16/3830 (0.4%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	114	ARG	NE-CZ-NH2	-17.58	111.51	120.30
1	A	114	ARG	NE-CZ-NH1	13.91	127.25	120.30
1	B	150	ARG	NE-CZ-NH2	-7.63	116.49	120.30
1	A	98	ARG	NE-CZ-NH2	-7.00	116.80	120.30
1	A	108	VAL	CG1-CB-CG2	6.88	121.91	110.90
1	B	150	ARG	NE-CZ-NH1	6.48	123.54	120.30
1	A	118	ASP	CB-CG-OD1	6.14	123.83	118.30
1	A	114	ARG	CD-NE-CZ	6.04	132.05	123.60
1	A	14	LYS	CD-CE-NZ	-5.79	98.37	111.70
1	A	28	ARG	NE-CZ-NH1	5.78	123.19	120.30
1	A	108	VAL	CB-CA-C	-5.51	100.93	111.40
1	A	114	ARG	CG-CD-NE	-5.49	100.27	111.80
1	A	129	LEU	CB-CG-CD1	-5.46	101.72	111.00
1	A	15	ASP	CB-CG-OD1	5.43	123.19	118.30
1	A	98	ARG	CG-CD-NE	-5.29	100.69	111.80
1	A	15	ASP	CB-CG-OD2	-5.12	113.69	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1482	0	1465	25	0
1	B	1288	0	1274	21	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	31	0	12	0	0
3	B	31	0	12	0	0
4	A	91	0	88	10	0
4	B	64	0	64	8	0
5	A	8	0	12	3	0
6	A	186	0	0	6	0
6	B	137	0	0	3	0
All	All	3320	0	2927	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20[B]:ARG:CG	1:A:20[B]:ARG:HH11	1.41	1.24
4:A:203:B12:H362	4:A:203:B12:H351	1.40	1.03
1:A:20[B]:ARG:NH1	1:A:20[B]:ARG:HG2	1.42	0.95
4:B:203:B12:H351	4:B:203:B12:H362	1.50	0.93
1:B:176:GLU:OE2	6:B:436:HOH:O	1.94	0.85
1:A:80:HIS:HD2	6:A:468:HOH:O	1.64	0.81
1:A:80:HIS:HE1	1:B:80:HIS:HE1	1.30	0.78
1:A:80:HIS:HE1	1:B:80:HIS:CE1	2.07	0.72
4:B:203:B12:H531	4:B:203:B12:H552	1.73	0.70
1:A:6:TYR:CG	1:B:98:ARG:NH1	2.59	0.70
1:A:80:HIS:CE1	1:B:80:HIS:HE1	2.10	0.68
1:B:92:THR:H	4:B:203:B12:H522	1.40	0.66
4:A:203:B12:C36	4:A:203:B12:H351	2.21	0.66
1:B:98:ARG:HG3	1:B:98:ARG:HH11	1.61	0.66
4:B:203:B12:C36	4:B:203:B12:H351	2.23	0.65
1:A:178:ARG:HH11	1:A:178:ARG:HB2	1.62	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20[B]:ARG:HG2	1:A:20[B]:ARG:HH11	0.57	0.65
4:A:203:B12:H291	4:A:203:B12:H3	1.62	0.64
1:A:16:ARG:HD3	1:A:20[A]:ARG:HH21	1.64	0.62
1:A:20[B]:ARG:CG	1:A:20[B]:ARG:NH1	2.18	0.62
1:A:14:LYS:HD3	1:A:18:ASP:OD2	2.00	0.61
1:A:196:TYR:OXT	1:B:55:HIS:HE1	1.84	0.61
1:A:55:HIS:HE1	1:B:196:TYR:OXT	1.84	0.60
5:A:204:EDO:H11	6:A:349:HOH:O	2.02	0.59
1:A:92:THR:O	1:A:93:TRP:HB2	2.02	0.59
1:A:6:TYR:CD1	1:B:98:ARG:NH1	2.72	0.58
1:A:11:GLN:NE2	1:B:96:GLN:HE22	2.03	0.56
4:B:203:B12:H531	4:B:203:B12:C55	2.35	0.56
1:B:64:PHE:CD2	1:B:132:MET:HG3	2.44	0.53
1:B:55:HIS:HD2	6:B:433:HOH:O	1.93	0.51
1:B:80:HIS:HD2	6:B:434:HOH:O	1.94	0.51
4:A:203:B12:N29	4:A:203:B12:H3	2.25	0.51
4:A:203:B12:H531	4:A:203:B12:H552	1.93	0.50
1:A:10:GLN:HE21	1:B:93:TRP:HA	1.75	0.50
4:A:203:B12:C3	4:A:203:B12:H291	2.25	0.49
5:A:204:EDO:H12	6:A:437:HOH:O	2.11	0.48
1:A:14:LYS:CD	1:A:18:ASP:OD2	2.60	0.48
4:B:203:B12:H262	4:B:203:B12:H601	1.95	0.48
1:A:65:ILE:HG21	4:A:203:B12:H462	1.94	0.48
1:B:98:ARG:HG3	1:B:98:ARG:NH1	2.30	0.47
4:B:203:B12:H3	4:B:203:B12:H291	1.81	0.46
1:A:14:LYS:HG2	1:B:93:TRP:HD1	1.80	0.46
4:A:203:B12:O51	6:A:462:HOH:O	2.21	0.46
4:A:203:B12:H262	4:A:203:B12:H91	1.83	0.46
1:A:6:TYR:CD2	1:B:98:ARG:NH1	2.84	0.45
4:B:203:B12:H3	4:B:203:B12:N29	2.32	0.45
1:A:20[A]:ARG:CZ	6:A:388:HOH:O	2.65	0.44
4:A:203:B12:H362	4:A:203:B12:C35	2.30	0.44
1:B:123:MET:HA	1:B:155:THR:O	2.18	0.43
5:A:204:EDO:C1	6:A:349:HOH:O	2.65	0.42
1:A:37:ASN:HD22	1:A:161:ARG:HG2	1.84	0.41
1:B:85:GLN:HG3	1:B:111:HIS:CG	2.56	0.41
1:A:43:THR:O	1:B:47:GLY:HA3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	190/191 (100%)	187 (98%)	3 (2%)	0	100	100
1	B	167/191 (87%)	165 (99%)	2 (1%)	0	100	100
All	All	357/382 (94%)	352 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	151/152 (99%)	146 (97%)	5 (3%)	45	32
1	B	131/152 (86%)	129 (98%)	2 (2%)	72	68
All	All	282/304 (93%)	275 (98%)	7 (2%)	55	45

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LYS
1	A	16	ARG
1	A	27	GLU
1	A	108	VAL
1	A	178	ARG
1	B	108	VAL
1	B	165	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	GLN
1	A	11	GLN
1	A	25	GLN
1	A	37	ASN
1	A	55	HIS
1	A	80	HIS
1	B	55	HIS
1	B	80	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ATP	A	202	2	24,33,33	1.38	4 (16%)	31,52,52	1.94	4 (12%)
4	B12	A	203	-	74,101,101	0.94	5 (6%)	111,166,166	2.07	18 (16%)
5	EDO	A	204	-	3,3,3	0.38	0	2,2,2	0.60	0
5	EDO	A	205	-	3,3,3	0.47	0	2,2,2	0.71	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ATP	B	202	2	24,33,33	1.89	4 (16%)	31,52,52	2.42	10 (32%)
4	B12	B	203	-	47,71,101	0.54	0	73,120,166	1.12	5 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	A	202	2	-	0/18/38/38	0/3/3/3
4	B12	A	203	-	-	1/51/223/223	0/3/11/11
5	EDO	A	204	-	-	0/1/1/1	0/0/0/0
5	EDO	A	205	-	-	0/1/1/1	0/0/0/0
3	ATP	B	202	2	-	0/18/38/38	0/3/3/3
4	B12	B	203	-	-	0/34/186/223	0/0/8/11

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	202	ATP	C2'-C3'	-2.10	1.47	1.53
4	A	203	B12	P-O5	2.38	1.54	1.48
4	A	203	B12	P-O2	2.44	1.66	1.59
3	A	202	ATP	C5-C4	2.47	1.46	1.40
3	A	202	ATP	C8-N7	2.53	1.39	1.34
4	A	203	B12	C11-C10	2.62	1.45	1.41
3	B	202	ATP	C5'-C4'	2.71	1.60	1.51
4	A	203	B12	C1P-C2P	3.11	1.59	1.51
3	A	202	ATP	C2-N3	3.53	1.38	1.32
4	A	203	B12	P-O3	3.63	1.69	1.59
3	B	202	ATP	C5-C4	3.76	1.49	1.40
3	B	202	ATP	O4'-C1'	3.99	1.46	1.41
3	B	202	ATP	C2-N3	4.48	1.40	1.32

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	202	ATP	N3-C2-N1	-8.59	122.32	128.89
3	A	202	ATP	C1'-N9-C4	-5.35	118.87	126.94
3	A	202	ATP	N3-C2-N1	-5.25	124.87	128.89
3	B	202	ATP	C1'-N9-C4	-4.36	120.37	126.94
4	A	203	B12	O5-P-O3	-3.78	97.93	109.36
4	B	203	B12	C9-C10-C11	-3.66	123.11	132.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	203	B12	C9-C10-C11	-3.48	123.56	132.28
4	A	203	B12	C35-C5-C4	-3.33	112.29	118.25
4	A	203	B12	C13-C14-C15	-3.27	120.84	131.88
4	B	203	B12	C25-C2-C1	-3.12	108.87	113.79
3	B	202	ATP	PB-O3B-PG	-3.05	122.42	132.67
4	A	203	B12	C2P-C1P-N59	-2.86	108.69	112.92
3	A	202	ATP	C4-C5-N7	-2.84	106.86	109.48
4	A	203	B12	C26-C2-C3	-2.74	102.15	107.58
4	B	203	B12	C1-C19-N24	-2.70	102.94	106.20
4	A	203	B12	O3-C2P-C1P	-2.59	101.37	106.79
4	B	203	B12	C13-C14-C15	-2.57	123.19	131.88
3	A	202	ATP	O4'-C1'-N9	-2.50	102.87	108.10
4	B	203	B12	C60-C18-C17	-2.46	110.90	115.68
3	B	202	ATP	O4'-C1'-N9	-2.46	102.95	108.10
4	A	203	B12	C20-C1-C19	-2.44	106.99	109.38
4	A	203	B12	C30-C3-C2	-2.40	114.36	119.11
4	A	203	B12	C3-C4-C5	-2.35	123.95	131.88
3	B	202	ATP	O5'-PA-O1A	-2.12	101.39	109.62
4	A	203	B12	C41-C42-C43	-2.00	106.19	112.53
4	A	203	B12	C53-C15-C16	2.22	122.22	118.25
4	A	203	B12	O3-P-O4	2.24	116.38	108.69
3	B	202	ATP	O5'-C5'-C4'	2.51	118.37	109.12
4	A	203	B12	C37-C7-C8	2.60	115.52	108.27
3	B	202	ATP	O2'-C2'-C3'	2.75	120.77	111.83
3	B	202	ATP	N6-C6-N1	2.77	125.14	119.20
4	A	203	B12	O2-C3R-C4R	2.83	121.11	109.99
3	B	202	ATP	C2-N1-C6	2.92	123.99	118.77
3	B	202	ATP	O2G-PG-O1G	3.42	121.60	110.58
4	A	203	B12	C35-C5-C6	4.74	126.74	118.25
4	A	203	B12	O2-P-O3	10.15	109.72	100.07
4	A	203	B12	P-O3-C2P	13.69	138.88	120.92

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	203	B12	P-O3-C2P-C3P

There are no ring outliers.

3 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	203	B12	10	0
5	A	204	EDO	3	0
4	B	203	B12	8	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	191/191 (100%)	0.21	16 (8%) 14 22	22, 31, 52, 67	0
1	B	169/191 (88%)	-0.17	3 (1%) 71 80	21, 30, 41, 57	1 (0%)
All	All	360/382 (94%)	0.03	19 (5%) 30 41	21, 31, 49, 67	1 (0%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	157	ILE	4.3
1	A	34	PHE	3.3
1	A	125	VAL	3.2
1	A	6	TYR	3.1
1	B	44	ALA	3.0
1	A	44	ALA	2.8
1	A	45	ALA	2.6
1	A	12	LYS	2.5
1	A	189	LYS	2.5
1	B	45	ALA	2.4
1	A	69	TRP	2.3
1	A	126	LEU	2.3
1	A	187	GLY	2.3
1	A	46	PHE	2.2
1	A	43	THR	2.2
1	A	159	THR	2.2
1	B	46	PHE	2.1
1	A	8	GLN	2.1
1	A	16	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	EDO	A	204	4/4	0.80	0.40	15.90	53,54,55,59	0
5	EDO	A	205	4/4	0.84	0.19	2.04	54,58,59,65	0
4	B12	A	203	91/91	0.95	0.15	0.20	29,40,72,82	0
3	ATP	B	202	31/31	0.98	0.12	0.00	23,31,34,38	0
4	B12	B	203	64/91	0.97	0.11	-0.10	22,27,37,44	0
3	ATP	A	202	31/31	0.98	0.09	-0.84	23,28,31,35	0
2	MG	B	201	1/1	0.99	0.10	-3.53	25,25,25,25	0
2	MG	A	201	1/1	0.98	0.14	-	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.