



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:37 AM GMT

PDB ID : 2HV7
Title : Crystal structure of phosphotyrosyl phosphatase activator bound to ATPgammaS
Authors : Chao, Y.; Jeffrey, J.D.; Shi, Y.
Deposited on : 2006-07-27
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	323	
1	B	323	
1	C	323	
1	D	323	
1	E	323	
1	F	323	
1	G	323	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	H	323	<div><div></div><div>74%</div><div>18%</div><div>• 7%</div></div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 19612 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein phosphatase 2A, regulatory subunit B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	301	Total	C	N	O	S	0	0	0
			2438	1586	406	435	11			
1	B	301	Total	C	N	O	S	0	0	0
			2438	1586	406	435	11			
1	C	301	Total	C	N	O	S	0	0	0
			2438	1586	406	435	11			
1	D	301	Total	C	N	O	S	0	0	0
			2438	1586	406	435	11			
1	E	301	Total	C	N	O	S	0	0	0
			2438	1586	406	435	11			
1	F	301	Total	C	N	O	S	0	0	0
			2438	1586	406	435	11			
1	G	301	Total	C	N	O	S	0	0	0
			2438	1586	406	435	11			
1	H	301	Total	C	N	O	S	0	0	0
			2438	1586	406	435	11			

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total 27	C 10	N 5	O 10	P 2	0	0
2	C	1	Total 27	C 10	N 5	O 10	P 2	0	0
2	E	1	Total 27	C 10	N 5	O 10	P 2	0	0
2	G	1	Total 27	C 10	N 5	O 10	P 2	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: Protein phosphatase 2A, regulatory subunit B




- Molecule 1: Protein phosphatase 2A, regulatory subunit B




- Molecule 1: Protein phosphatase 2A, regulatory subunit B




- Molecule 1: Protein phosphatase 2A, regulatory subunit B




- Molecule 1: Protein phosphatase 2A, regulatory subunit B




- Molecule 1: Protein phosphatase 2A, regulatory subunit B



- Molecule 1: Protein phosphatase 2A, regulatory subunit B



● Molecule 1: Protein phosphatase 2A, regulatory subunit B



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	80.04Å 86.10Å 95.10Å 73.91° 89.97° 73.31°	Depositor
Resolution (Å)	50.00 – 2.50	Depositor
% Data completeness (in resolution range)	87.7 (50.00-2.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.249 , 0.312	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	19612	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/2506	0.61	0/3401
1	B	0.55	0/2506	0.61	0/3401
1	C	0.55	1/2506 (0.0%)	0.59	0/3401
1	D	0.50	1/2506 (0.0%)	0.58	0/3401
1	E	0.51	0/2506	0.59	0/3401
1	F	0.73	4/2506 (0.2%)	0.64	2/3401 (0.1%)
1	G	0.51	0/2506	0.57	0/3401
1	H	0.53	0/2506	0.62	1/3401 (0.0%)
All	All	0.56	6/20048 (0.0%)	0.60	3/27208 (0.0%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	229	ASP	CG-OD1	18.58	1.68	1.25
1	F	229	ASP	CG-OD2	14.23	1.58	1.25
1	D	176	ASP	CG-OD1	7.17	1.41	1.25
1	F	53	LEU	CG-CD2	6.77	1.76	1.51
1	F	53	LEU	CG-CD1	5.71	1.73	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	229	ASP	CB-CG-OD1	-7.56	111.50	118.30
1	F	229	ASP	CB-CG-OD2	-5.39	113.45	118.30
1	H	98	SER	N-CA-C	5.34	125.42	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2438	0	2423	52	0
1	B	2438	0	2423	48	0
1	C	2438	0	2423	43	0
1	D	2438	0	2423	55	0
1	E	2438	0	2423	54	0
1	F	2438	0	2423	59	0
1	G	2438	0	2423	55	0
1	H	2438	0	2423	34	0
2	B	27	0	12	2	0
2	C	27	0	12	1	0
2	E	27	0	12	0	0
2	G	27	0	12	3	0
All	All	19612	0	19432	378	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 378 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:53:LEU:CD2	1:F:53:LEU:CG	1.76	1.59
1:F:229:ASP:OD1	1:F:229:ASP:CG	1.68	1.30
1:F:263:MET:HE3	1:G:97:PRO:HB2	1.38	1.00
1:E:235:PRO:HG2	1:E:281:VAL:O	1.64	0.97
1:F:293:ARG:HG3	1:F:293:ARG:HH11	1.32	0.94

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/323 (93%)	280 (94%)	16 (5%)	3 (1%)	19	34
1	B	299/323 (93%)	277 (93%)	22 (7%)	0	100	100
1	C	299/323 (93%)	278 (93%)	20 (7%)	1 (0%)	46	68
1	D	299/323 (93%)	278 (93%)	20 (7%)	1 (0%)	46	68
1	E	299/323 (93%)	267 (89%)	30 (10%)	2 (1%)	26	46
1	F	299/323 (93%)	276 (92%)	21 (7%)	2 (1%)	26	46
1	G	299/323 (93%)	270 (90%)	27 (9%)	2 (1%)	26	46
1	H	299/323 (93%)	272 (91%)	26 (9%)	1 (0%)	46	68
All	All	2392/2584 (93%)	2198 (92%)	182 (8%)	12 (0%)	34	55

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	98	SER
1	A	97	PRO
1	A	98	SER
1	C	208	GLY
1	D	209	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/280 (94%)	250 (95%)	13 (5%)	31	55
1	B	263/280 (94%)	248 (94%)	15 (6%)	25	46
1	C	263/280 (94%)	250 (95%)	13 (5%)	31	55
1	D	263/280 (94%)	244 (93%)	19 (7%)	18	33
1	E	263/280 (94%)	253 (96%)	10 (4%)	40	67
1	F	263/280 (94%)	243 (92%)	20 (8%)	16	30

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	263/280 (94%)	243 (92%)	20 (8%)	16	30
1	H	263/280 (94%)	248 (94%)	15 (6%)	25	46
All	All	2104/2240 (94%)	1979 (94%)	125 (6%)	24	44

5 of 125 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	274	GLN
1	E	321	THR
1	H	94	VAL
1	D	285	SER
1	E	53	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 53 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	277	ASN
1	E	237	HIS
1	H	186	ASN
1	D	318	HIS
1	E	177	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ADP	B	324	-	22,29,29	1.12	3 (13%)	27,45,45	1.89	5 (18%)
2	ADP	C	324	-	22,29,29	1.17	3 (13%)	27,45,45	1.73	2 (7%)
2	ADP	E	324	-	22,29,29	1.17	2 (9%)	27,45,45	1.87	5 (18%)
2	ADP	G	324	-	22,29,29	1.21	2 (9%)	27,45,45	1.73	3 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	B	324	-	-	0/12/32/32	0/3/3/3
2	ADP	C	324	-	-	0/12/32/32	0/3/3/3
2	ADP	E	324	-	-	0/12/32/32	0/3/3/3
2	ADP	G	324	-	-	0/12/32/32	0/3/3/3

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	324	ADP	C2-N3	2.02	1.35	1.32
2	B	324	ADP	C2-N3	2.03	1.35	1.32
2	B	324	ADP	O4'-C1'	2.35	1.44	1.41
2	C	324	ADP	O4'-C1'	2.64	1.44	1.41
2	E	324	ADP	O4'-C1'	2.75	1.44	1.41

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	324	ADP	N3-C2-N1	-7.18	123.40	128.89
2	C	324	ADP	N3-C2-N1	-6.86	123.64	128.89
2	G	324	ADP	N3-C2-N1	-6.80	123.69	128.89
2	B	324	ADP	N3-C2-N1	-6.68	123.78	128.89
2	B	324	ADP	C4-C5-N7	-4.23	105.59	109.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	324	ADP	2	0
2	C	324	ADP	1	0
2	G	324	ADP	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.