



# Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 05:19 PM GMT

PDB ID : 4HY1

Title : Pyrrolopyrimidine inhibitors of dna gyrase b and topoisomerase iv, part i: structure guided discovery and optimization of dual targeting agents with potent, broad-spectrum enzymatic activity.

Authors : Bensen, D.C.; Creighton, C.J.; Kwan, B; Tari, L.W.

Deposited on : 2012-11-12

Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.7 (RC4), CSD as536be (2015)

Xtriage (Phenix) : 1.9-1692

EDS : rb-20026688

Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)

Refmac : 5.8.0135

CCP4 : 6.5.0

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 5292 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Topoisomerase IV, subunit B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	360	Total	C	N	O	S	0	3	0
			2855	1822	477	548	8			
1	B	228	Total	C	N	O	S	0	0	0
			1806	1156	302	343	5			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	258	PRO	LEU	SEE REMARK 999	UNP Q2A1P5
A	383	LEU	-	EXPRESSION TAG	UNP Q2A1P5
A	384	GLU	-	EXPRESSION TAG	UNP Q2A1P5
A	385	HIS	-	EXPRESSION TAG	UNP Q2A1P5
A	386	HIS	-	EXPRESSION TAG	UNP Q2A1P5
A	387	HIS	-	EXPRESSION TAG	UNP Q2A1P5
A	388	HIS	-	EXPRESSION TAG	UNP Q2A1P5
A	389	HIS	-	EXPRESSION TAG	UNP Q2A1P5
A	390	HIS	-	EXPRESSION TAG	UNP Q2A1P5
B	258	PRO	LEU	SEE REMARK 999	UNP Q2A1P5
B	383	LEU	-	EXPRESSION TAG	UNP Q2A1P5
B	384	GLU	-	EXPRESSION TAG	UNP Q2A1P5
B	385	HIS	-	EXPRESSION TAG	UNP Q2A1P5
B	386	HIS	-	EXPRESSION TAG	UNP Q2A1P5
B	387	HIS	-	EXPRESSION TAG	UNP Q2A1P5
B	388	HIS	-	EXPRESSION TAG	UNP Q2A1P5
B	389	HIS	-	EXPRESSION TAG	UNP Q2A1P5
B	390	HIS	-	EXPRESSION TAG	UNP Q2A1P5

- Molecule 2 is 6-({4-[{(3R)-3-AMINOPYRROLIDIN-1-YL]-5-CHLORO-6-ETHYL-7H-PYRROLO[2,3-D]PYRIMIDIN-2-YL}SULFANYL)-1H-ISOINDOL-1-ONE (three-letter code: 19X) (formula: C<sub>20</sub>H<sub>19</sub>ClN<sub>6</sub>OS).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	Cl	N	O	S		
2	A	1	29	20	1	6	1	1	0	0
2	B	1	29	20	1	6	1	1	0	0



- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	401	Total O 401 401		0	0
3	B	172	Total O 172 172		0	0



### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Topoisomerase IV, subunit B



- Molecule 1: Topoisomerase IV, subunit B



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	44.16 Å    130.28 Å    80.27 Å 90.00°    98.12°    90.00°	Depositor
Resolution (Å)	38.96 – 1.90 38.95 – 1.89	Depositor EDS
% Data completeness (in resolution range)	99.6 (38.96-1.90) 98.6 (38.95-1.89)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.93 (at 1.89 Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
$R$ , $R_{free}$	0.241 , 0.299 0.252 , 0.307	Depositor DCC
$R_{free}$ test set	3538 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.7	Xtriage
Anisotropy	0.356	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 46.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Outliers	0 of 70303 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5292	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 19X

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.77	1/2916 (0.0%)	0.83	3/3939 (0.1%)
1	B	0.61	1/1837 (0.1%)	0.72	1/2473 (0.0%)
All	All	0.71	2/4753 (0.0%)	0.79	4/6412 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	167	TRP	CD2-CE2	6.44	1.49	1.41
1	B	212	TRP	CD2-CE2	5.44	1.47	1.41

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	228	ALA	N-CA-C	5.63	126.20	111.00
1	B	231	LEU	CA-CB-CG	5.47	127.89	115.30
1	A	201	TYR	CA-CB-CG	5.20	123.27	113.40
1	A	12	LEU	CA-CB-CG	5.14	127.13	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2855	0	2883	49	0
1	B	1806	0	1839	31	0
2	A	29	0	19	2	0
2	B	29	0	19	3	0
3	A	401	0	0	11	0
3	B	172	0	0	4	0
All	All	5292	0	4760	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:365:ARG:HD3	3:A:1823:HOH:O	1.56	1.03
1:A:343:ASN:HD22	1:A:343:ASN:H	1.05	0.98
1:A:227:GLU:HA	1:A:228:ALA:O	1.69	0.92
1:A:343:ASN:H	1:A:343:ASN:ND2	1.69	0.90
1:A:73:MET:HE1	1:A:86:ILE:HD12	1.54	0.89
1:A:73:MET:CE	1:A:86:ILE:HD12	2.04	0.87
1:A:256:GLU:O	1:A:258:PRO:HD3	1.77	0.84
1:A:73:MET:CE	1:A:86:ILE:CD1	2.58	0.81
1:B:21:ARG:HG2	1:B:21:ARG:HH11	1.52	0.74
1:A:73:MET:HE2	1:A:86:ILE:CD1	2.17	0.74
1:B:155:VAL:HG13	1:B:159:ASN:HB3	1.70	0.72
1:A:21:ARG:HB3	1:A:24[B]:MET:HG3	1.71	0.71
1:A:186:LEU:HD12	3:A:1574:HOH:O	1.92	0.69
1:A:262:ILE:H	1:A:316:ASN:HD22	1.41	0.68
1:A:73:MET:HE2	1:A:86:ILE:HD13	1.79	0.64
1:A:200:LYS:HE2	3:A:1677:HOH:O	1.96	0.64
1:A:343:ASN:HD22	1:A:343:ASN:N	1.86	0.63
1:A:73:MET:HG2	2:A:1401:19X:C8	2.29	0.62
1:B:213:HIS:HE1	1:B:215:GLU:OE1	1.83	0.62
1:B:78:HIS:HD2	3:B:1634:HOH:O	1.83	0.61
1:A:73:MET:CE	1:A:86:ILE:HD13	2.33	0.59
1:A:195:LYS:HE2	3:A:1709:HOH:O	2.02	0.59
1:A:262:ILE:H	1:A:316:ASN:ND2	2.01	0.58
1:A:73:MET:HE2	1:A:86:ILE:HD12	1.82	0.58
1:A:36[B]:GLN:HE21	1:A:190:LYS:NZ	2.02	0.57
1:A:73:MET:HE1	1:A:86:ILE:CD1	2.28	0.56
1:A:261:SER:HA	1:A:316:ASN:HD21	1.71	0.55
1:B:216:THR:HB	1:B:219:LYS:HB3	1.87	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:79:PRO:O	1:B:80:GLU:HB2	2.06	0.55
1:A:304:ILE:N	3:A:1663:HOH:O	2.41	0.54
1:A:36[B]:GLN:NE2	1:A:190:LYS:NZ	2.56	0.54
2:B:1401:19X:H6	2:B:1401:19X:CL1	2.45	0.54
1:B:18:VAL:HG12	1:B:121:LEU:CD1	2.39	0.53
1:A:327:GLN:HG2	3:A:1653:HOH:O	2.08	0.53
1:A:377:LYS:HG3	3:A:1833:HOH:O	2.09	0.52
1:A:41:ASN:ND2	1:A:115:VAL:HG13	2.25	0.52
1:B:18:VAL:HG12	1:B:121:LEU:HD13	1.92	0.52
1:A:226:LEU:O	1:A:227:GLU:C	2.48	0.51
1:B:21:ARG:HB3	1:B:24:MET:HG2	1.93	0.51
1:A:273:PRO:HD2	1:A:334:GLU:HB3	1.90	0.51
1:B:91:THR:O	1:B:91:THR:HG22	2.12	0.50
1:A:36[A]:GLN:HG2	1:A:186:LEU:HD22	1.94	0.49
1:B:63:SER:HB3	1:B:167:TRP:CD1	2.48	0.49
1:A:115:VAL:HG11	2:A:1401:19X:CL1	2.50	0.48
1:A:256:GLU:C	1:A:258:PRO:HD3	2.34	0.47
1:A:77:ILE:HD11	3:A:1609:HOH:O	2.13	0.47
1:B:89:ILE:HD13	2:B:1401:19X:H14	1.98	0.46
1:B:195:LYS:H	1:B:195:LYS:HD3	1.81	0.46
1:A:36[B]:GLN:HE21	1:A:190:LYS:HZ1	1.63	0.45
1:B:220:GLY:HA2	1:B:223:ASP:OD1	2.16	0.45
1:B:82:LYS:HG2	3:B:1522:HOH:O	2.16	0.45
1:B:141:GLU:HB2	1:B:146:THR:HG21	1.99	0.44
1:B:79:PRO:O	1:B:80:GLU:CB	2.66	0.44
1:B:221:TYR:N	3:B:1571:HOH:O	2.51	0.44
1:A:78:HIS:HA	1:A:79:PRO:HD3	1.87	0.44
1:A:147[A]:LYS:HE3	1:A:147[A]:LYS:HB2	1.47	0.43
1:A:261:SER:HA	1:A:316:ASN:ND2	2.34	0.43
1:A:73:MET:HG3	1:A:162:THR:HG21	2.01	0.43
1:B:201:TYR:CZ	1:B:203:ASN:HB2	2.53	0.43
1:B:73:MET:SD	1:B:86:ILE:HD12	2.58	0.43
1:A:227:GLU:HA	1:A:228:ALA:C	2.38	0.43
1:A:343:ASN:HB3	3:A:1600:HOH:O	2.19	0.43
1:A:256:GLU:HG2	3:A:1691:HOH:O	2.19	0.43
1:B:21:ARG:HG2	1:B:21:ARG:NH1	2.24	0.43
1:A:147[B]:LYS:HB3	1:A:147[B]:LYS:HE3	1.55	0.42
1:B:134:ASN:HB3	1:B:151:ILE:HG23	2.01	0.42
1:B:30:ASN:HA	1:B:175:ASP:O	2.20	0.42
1:A:232:PRO:HB3	1:A:356:TRP:CZ2	2.55	0.42
1:B:18:VAL:HG23	1:B:93:LEU:HD11	2.00	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:126:GLU:HG2	1:B:139:VAL:HG22	2.01	0.42
1:A:12:LEU:HD21	1:A:94:HIS:CD2	2.55	0.42
1:B:228:ALA:HB3	3:B:1563:HOH:O	2.20	0.41
1:B:197:LEU:O	1:B:213:HIS:HD2	2.03	0.41
1:A:200:LYS:CE	3:A:1677:HOH:O	2.64	0.41
1:A:305:LYS:O	1:A:305:LYS:HD3	2.21	0.41
1:B:20:LYS:HG2	1:B:21:ARG:HG3	2.02	0.40
1:B:41:ASN:HB3	2:B:1401:19X:CL1	2.58	0.40
1:A:36[B]:GLN:NE2	1:A:190:LYS:HZ3	2.18	0.40
1:B:90:MET:HE2	1:B:115:VAL:HG23	2.03	0.40
1:B:139:VAL:HB	1:B:147:LYS:HB3	2.04	0.40
1:A:223:ASP:O	1:A:227:GLU:HG2	2.22	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	357/390 (92%)	344 (96%)	11 (3%)	2 (1%)	30 17
1	B	220/390 (56%)	207 (94%)	11 (5%)	2 (1%)	21 9
All	All	577/780 (74%)	551 (96%)	22 (4%)	4 (1%)	26 14

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	228	ALA
1	B	80	GLU
1	B	259	SER
1	A	227	GLU

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	319/344 (93%)	302 (95%)	17 (5%)	28 16
1	B	202/344 (59%)	188 (93%)	14 (7%)	19 8
All	All	521/688 (76%)	490 (94%)	31 (6%)	25 12

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	LEU
1	A	58	LEU
1	A	110	LEU
1	A	147[A]	LYS
1	A	147[B]	LYS
1	A	171	LYS
1	A	186	LEU
1	A	222	LEU
1	A	226	LEU
1	A	227	GLU
1	A	234	GLU
1	A	278	HIS
1	A	296	LYS
1	A	305	LYS
1	A	315	LEU
1	A	343	ASN
1	A	377	LYS
1	B	8	SER
1	B	12	LEU
1	B	20	LYS
1	B	58	LEU
1	B	83	MET
1	B	95	SER
1	B	121	LEU
1	B	155	VAL
1	B	192	ILE
1	B	195	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	216	THR
1	B	219	LYS
1	B	223	ASP
1	B	224	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	94	HIS
1	A	134	ASN
1	A	154	ASN
1	A	240	ASN
1	A	316	ASN
1	A	343	ASN
1	B	36	GLN
1	B	213	HIS

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	19X	A	1401	-	31,33,33	2.18	10 (32%)	33,49,49	2.72	17 (51%)
2	19X	B	1401	-	31,33,33	2.36	10 (32%)	33,49,49	2.75	17 (51%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	19X	A	1401	-	-	0/9/28/28	0/5/5/5
2	19X	B	1401	-	-	0/9/28/28	0/5/5/5

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1401	19X	C10-S19	-5.24	1.68	1.76
2	A	1401	19X	C20-S19	-3.86	1.70	1.77
2	B	1401	19X	C20-S19	-3.74	1.70	1.77
2	A	1401	19X	C10-S19	-3.68	1.70	1.76
2	A	1401	19X	C22-C28	-2.24	1.47	1.50
2	A	1401	19X	C3-C2	2.09	1.48	1.43
2	B	1401	19X	C8-N9	2.18	1.35	1.32
2	B	1401	19X	C3-C2	2.35	1.49	1.43
2	B	1401	19X	O29-C28	2.76	1.29	1.23
2	A	1401	19X	C8-C3	2.80	1.49	1.43
2	B	1401	19X	C23-C26	3.00	1.46	1.42
2	A	1401	19X	C23-C26	3.06	1.46	1.42
2	A	1401	19X	O29-C28	3.21	1.29	1.23
2	B	1401	19X	C8-C3	3.92	1.51	1.43
2	A	1401	19X	C22-C23	4.11	1.47	1.41
2	B	1401	19X	C22-C23	4.26	1.47	1.41
2	A	1401	19X	C4-C5	4.76	1.46	1.38
2	B	1401	19X	C26-N27	4.79	1.49	1.33
2	A	1401	19X	C26-N27	4.82	1.49	1.33
2	B	1401	19X	C4-C5	5.19	1.47	1.38

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1401	19X	C22-C21-C20	-5.23	115.84	121.04
2	B	1401	19X	C22-C21-C20	-5.03	116.05	121.04
2	A	1401	19X	C4-C3-C2	-4.97	97.60	106.63

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1401	19X	C4-C3-C2	-4.61	98.25	106.63
2	A	1401	19X	C5-C4-CL1	-4.07	120.17	126.59
2	A	1401	19X	O29-C28-C22	-3.56	118.99	122.67
2	B	1401	19X	C23-C22-C28	-3.53	101.04	107.30
2	B	1401	19X	C25-C24-C23	-3.49	116.95	121.67
2	A	1401	19X	C25-C24-C23	-3.48	116.95	121.67
2	A	1401	19X	C23-C22-C28	-3.46	101.18	107.30
2	B	1401	19X	C15-C14-N13	-3.16	99.24	103.35
2	A	1401	19X	C23-C26-N27	-2.98	103.78	112.39
2	B	1401	19X	C23-C26-N27	-2.89	104.01	112.39
2	B	1401	19X	C5-C4-CL1	-2.48	122.68	126.59
2	A	1401	19X	C6-C5-C4	-2.36	126.43	130.41
2	B	1401	19X	C24-C23-C26	-2.21	128.06	135.42
2	B	1401	19X	C7-C6-C5	-2.19	109.37	115.03
2	A	1401	19X	C24-C23-C26	-2.03	128.66	135.42
2	A	1401	19X	C24-C25-C20	2.07	123.26	120.58
2	B	1401	19X	C15-C16-C17	2.14	105.18	102.03
2	A	1401	19X	C22-C28-N27	2.27	114.22	109.37
2	B	1401	19X	C22-C28-N27	2.28	114.26	109.37
2	B	1401	19X	C21-C22-C23	2.42	123.01	120.43
2	A	1401	19X	C3-C8-N13	2.49	127.61	121.44
2	B	1401	19X	C3-C8-N13	2.52	127.67	121.44
2	B	1401	19X	C14-C15-C16	2.69	109.06	103.66
2	B	1401	19X	C10-N9-C8	3.06	122.03	113.12
2	A	1401	19X	C10-N9-C8	3.12	122.21	113.12
2	A	1401	19X	C21-C22-C23	3.34	124.00	120.43
2	A	1401	19X	C14-C15-C16	3.48	110.65	103.66
2	A	1401	19X	C10-S19-C20	4.97	110.92	103.39
2	A	1401	19X	C21-C22-C28	5.65	134.79	128.87
2	B	1401	19X	C10-S19-C20	6.43	113.14	103.39
2	B	1401	19X	C21-C22-C28	6.75	135.94	128.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1401	19X	2	0
2	B	1401	19X	3	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	360/390 (92%)	0.23	17 (4%) 35 38	23, 33, 55, 77	0
1	B	228/390 (58%)	1.19	43 (18%) 2 2	30, 51, 83, 113	0
All	All	588/780 (75%)	0.60	60 (10%) 9 10	23, 38, 73, 113	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	224	HIS	8.3
1	B	267	VAL	6.5
1	B	233	ALA	6.4
1	B	230	THR	5.8
1	B	225	LYS	5.7
1	B	220	GLY	5.7
1	B	234	GLU	5.5
1	B	113	VAL	5.4
1	B	221	TYR	4.6
1	B	231	LEU	4.5
1	B	263	LYS	4.5
1	B	260	GLU	4.4
1	B	262	ILE	4.4
1	B	265	SER	4.3
1	B	24	MET	4.3
1	B	217	GLY	4.1
1	B	228	ALA	4.0
1	B	99	PHE	3.9
1	B	229	GLU	3.8
1	B	218	LEU	3.7
1	B	226	LEU	3.7
1	B	223	ASP	3.7
1	B	232	PRO	3.6
1	A	227	GLU	3.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	94	HIS	3.6
1	B	79	PRO	3.5
1	B	222	LEU	3.4
1	A	306	ILE	3.3
1	B	264	ASN	3.3
1	B	93	LEU	3.1
1	A	305	LYS	3.1
1	B	25	TYR	3.1
1	A	38	ILE	2.9
1	B	100	SER	2.8
1	B	261	SER	2.8
1	A	258	PRO	2.7
1	A	64	ILE	2.6
1	B	214	PHE	2.6
1	B	15	LEU	2.6
1	B	98	LYS	2.5
1	A	34	LEU	2.5
1	B	82	LYS	2.5
1	B	216	THR	2.5
1	B	219	LYS	2.4
1	B	96	GLY	2.4
1	B	148	ASP	2.3
1	A	35	ILE	2.3
1	B	192	ILE	2.3
1	A	36[A]	GLN	2.3
1	B	144	PHE	2.3
1	A	376	GLN	2.3
1	A	260	GLU	2.2
1	B	258	PRO	2.2
1	A	118	VAL	2.2
1	B	38	ILE	2.1
1	A	187	LEU	2.1
1	A	307	THR	2.1
1	A	374	VAL	2.1
1	A	39	ILE	2.0
1	A	43	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	19X	A	1401	29/29	0.93	0.12	-0.50	22,29,46,47	0
2	19X	B	1401	29/29	0.83	0.13	-0.51	39,54,64,68	0

### 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.