



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:39 AM GMT

PDB ID : 2I0R
Title : Crystal structure of aromatic amine dehydrogenase TTQ-formamide adduct
Authors : Roujeinikova, A.; Leys, D.
Deposited on : 2006-08-11
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1199 (1.40-1.40)
Clashscore	102246	1295 (1.40-1.40)
Ramachandran outliers	100387	1259 (1.40-1.40)
Sidechain outliers	100360	1258 (1.40-1.40)
RSRZ outliers	91569	1198 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	124	<div> <div>5%</div> <div> <div></div> <div>86%</div> <div>14%</div> </div> </div>
1	H	124	<div> <div>2%</div> <div> <div></div> <div>90%</div> <div>8%</div> <div>...</div> </div> </div>
2	A	361	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>9%</div> <div>.</div> </div> </div>
2	B	361	<div> <div>%</div> <div> <div></div> <div>92%</div> <div>8%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9178 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aromatic Amine Dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	124	Total	C	N	O	S	0	0	0
			944	574	166	189	15			
1	H	122	Total	C	N	O	S	0	1	0
			941	573	165	188	15			

- Molecule 2 is a protein called Aromatic Amine Dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	360	Total	C	N	O	S	0	2	0
			2809	1771	489	535	14			
2	B	360	Total	C	N	O	S	0	1	0
			2784	1756	481	534	13			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	199	Total	O	0	0
			199	199		
3	H	195	Total	O	0	0
			195	195		
3	A	672	Total	O	0	0
			672	672		
3	B	634	Total	O	0	0
			634	634		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Aromatic Amine Dehydrogenase



- Molecule 1: Aromatic Amine Dehydrogenase



- Molecule 2: Aromatic Amine Dehydrogenase



- Molecule 2: Aromatic Amine Dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.66Å 88.42Å 80.03Å 90.00° 90.45° 90.00°	Depositor
Resolution (Å)	15.00 – 1.40 14.98 – 1.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (15.00-1.40) 99.8 (14.98-1.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.51 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.139 , 0.165 0.141 , 0.166	Depositor DCC
R_{free} test set	9682 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	8.6	Xtriage
Anisotropy	0.308	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 57.6	EDS
Estimated twinning fraction	0.018 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 192630 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9178	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1TQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	D	0.60	0/948	0.70	0/1289
1	H	0.62	0/950	0.73	0/1295
2	A	0.63	0/2880	0.78	1/3905 (0.0%)
2	B	0.61	0/2853	0.76	3/3871 (0.1%)
All	All	0.62	0/7631	0.75	4/10360 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	263	ARG	NE-CZ-NH2	-7.71	116.45	120.30
2	B	255	ASP	CB-CG-OD1	5.14	122.93	118.30
2	B	218	ASP	CB-CG-OD2	-5.10	113.71	118.30
2	B	148	ARG	NE-CZ-NH2	-5.02	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	944	0	826	13	0
1	H	941	0	827	10	0
2	A	2809	0	2743	36	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	2784	0	2695	24	0
3	A	672	0	0	8	0
3	B	634	0	0	5	0
3	D	199	0	0	3	0
3	H	195	0	0	4	0
All	All	9178	0	7091	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:105:SER:HA	1:H:164:ASN:HD21	1.51	0.75
2:B:104:ARG:HH11	2:B:106:HIS:HE1	1.33	0.75
2:A:104:ARG:HH11	2:A:106:HIS:HE1	1.31	0.75
2:B:124:ASN:HD21	2:B:178:GLY:H	1.34	0.73
1:D:105:SER:HA	1:D:164:ASN:HD21	1.55	0.71
2:B:74:GLU:N	3:B:1011:HOH:O	2.21	0.71
2:A:124:ASN:HD21	2:A:178:GLY:H	1.36	0.71
1:D:70:ASP:C	1:D:71:GLU:CA	2.59	0.70
2:B:91:TYR:OH	2:B:428:HIS:HD2	1.73	0.70
3:H:246:HOH:O	2:B:118:MET:SD	2.52	0.68
2:A:91:TYR:OH	2:A:428:HIS:HD2	1.77	0.67
2:A:73:ARG:N	3:A:1075:HOH:O	2.27	0.67
2:A:261:GLN:NE2	3:A:594:HOH:O	2.21	0.66
2:B:197:GLN:NE2	2:B:227:SER:H	1.94	0.65
2:A:335:ARG:H	2:A:386:GLN:HE22	1.44	0.64
2:A:197:GLN:NE2	2:A:227:SER:H	1.97	0.63
2:A:263:ARG:HG3	3:A:737:HOH:O	2.01	0.61
2:A:335:ARG:H	2:A:386:GLN:NE2	2.00	0.59
3:D:204:HOH:O	2:B:143:HIS:HE1	1.87	0.57
2:A:91:TYR:OH	2:A:428:HIS:CD2	2.58	0.57
2:A:143:HIS:HD2	3:A:685:HOH:O	1.86	0.57
3:H:221:HOH:O	2:A:143:HIS:HE1	1.88	0.56
1:D:176:LEU:HD23	2:A:73:ARG:HD3	1.86	0.56
1:D:68:ASN:HD21	1:D:70:ASP:HB2	1.69	0.56
2:B:335:ARG:H	2:B:386:GLN:NE2	2.03	0.56
2:B:216:VAL:O	2:B:217:GLU:HB3	2.06	0.56
1:H:156:ASN:HD22	1:H:156:ASN:C	2.09	0.56
2:B:106:HIS:HD2	2:B:418:ALA:O	1.89	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:197:GLN:HE21	2:B:227:SER:H	1.54	0.55
2:A:197:GLN:HE21	2:A:227:SER:H	1.53	0.55
2:B:335:ARG:H	2:B:386:GLN:HE22	1.53	0.55
3:D:302:HOH:O	2:A:118[A]:MET:SD	2.58	0.55
2:A:406[B]:GLN:HA	2:A:406[B]:GLN:HE21	1.71	0.54
2:A:106:HIS:HD2	2:A:418:ALA:O	1.92	0.53
1:D:167:SER:HB2	2:B:177:GLN:HE22	1.71	0.53
2:A:133:GLY:O	2:A:428:HIS:HE1	1.92	0.53
2:A:398:ASN:HD22	2:A:416:GLU:HA	1.75	0.52
2:B:398:ASN:HD22	2:B:417:GLY:H	1.59	0.51
1:H:159:ASN:HD22	2:A:177:GLN:NE2	2.08	0.51
1:H:167:SER:HB2	2:A:177:GLN:HE22	1.76	0.50
1:H:109:1TQ:HB2	1:H:160:TRP:NE1	2.27	0.50
1:D:159:ASN:HD22	2:B:177:GLN:NE2	2.09	0.50
2:A:128:GLN:HE22	2:A:186:ARG:C	2.15	0.49
1:D:109:1TQ:HB2	1:D:160:TRP:NE1	2.28	0.48
1:D:155:HIS:HE1	3:B:446:HOH:O	1.97	0.48
2:B:398:ASN:HD22	2:B:416:GLU:HA	1.79	0.47
1:D:176:LEU:HD23	2:A:73:ARG:CD	2.45	0.47
2:A:131:ASN:H	2:A:187:GLN:HE22	1.64	0.45
2:A:275:ILE:HA	2:A:290:SER:HA	1.99	0.45
2:B:398:ASN:ND2	2:B:417:GLY:H	2.14	0.45
2:B:143:HIS:HD2	3:B:610:HOH:O	1.99	0.44
1:H:109:1TQ:O1	1:H:128:ASP:OD2	2.35	0.44
1:D:109:1TQ:O1	1:D:128:ASP:OD2	2.36	0.44
2:B:275:ILE:HA	2:B:290:SER:HA	2.00	0.44
2:A:186:ARG:HH22	2:A:426:GLN:NE2	2.16	0.44
1:H:155:HIS:HE1	3:A:453:HOH:O	1.99	0.44
2:A:406[A]:GLN:H	2:A:406[A]:GLN:CD	2.21	0.43
3:H:221:HOH:O	2:A:143:HIS:CE1	2.68	0.43
1:H:146:GLU:CD	2:A:354:LYS:HZ1	2.21	0.43
2:B:265:LYS:CG	3:B:849:HOH:O	2.66	0.43
1:H:117:HIS:HE1	3:H:209:HOH:O	2.00	0.43
1:H:109:1TQ:HB2	1:H:160:TRP:HE1	1.84	0.43
1:D:112:THR:HA	1:D:122:TYR:O	2.20	0.42
2:A:308:ASP:C	2:A:308:ASP:OD1	2.57	0.41
2:A:398:ASN:HD22	2:A:417:GLY:H	1.68	0.41
2:A:297:SER:OG	3:A:627:HOH:O	2.21	0.41
1:D:143:GLN:HE22	2:B:354:LYS:HZ1	1.68	0.41
2:A:406[B]:GLN:NE2	3:A:887:HOH:O	2.53	0.41
2:A:114:LYS:NZ	3:A:1030:HOH:O	2.52	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:406[B]:GLN:HA	2:A:406[B]:GLN:NE2	2.36	0.41
2:A:131:ASN:H	2:A:187:GLN:NE2	2.19	0.41
2:B:91:TYR:OH	2:B:428:HIS:CD2	2.62	0.40
1:D:146:GLU:OE2	1:D:155:HIS:HD2	2.05	0.40
3:D:204:HOH:O	2:B:143:HIS:CE1	2.68	0.40
2:B:321:LYS:HE3	3:B:718:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	119/124 (96%)	116 (98%)	3 (2%)	0	100	100
1	H	120/124 (97%)	117 (98%)	3 (2%)	0	100	100
2	A	360/361 (100%)	348 (97%)	11 (3%)	1 (0%)	46	19
2	B	358/361 (99%)	344 (96%)	13 (4%)	1 (0%)	46	19
All	All	957/970 (99%)	925 (97%)	30 (3%)	2 (0%)	52	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	200	SER
2	B	200	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	105/108 (97%)	104 (99%)	1 (1%)	82	59
1	H	107/108 (99%)	106 (99%)	1 (1%)	84	63
2	A	304/304 (100%)	301 (99%)	3 (1%)	82	59
2	B	300/304 (99%)	299 (100%)	1 (0%)	94	83
All	All	816/824 (99%)	810 (99%)	6 (1%)	90	70

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	182	ASN
1	H	156	ASN
2	A	104	ARG
2	A	406[A]	GLN
2	A	406[B]	GLN
2	B	104	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (44) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	68	ASN
1	D	139	GLN
1	D	143	GLN
1	D	155	HIS
1	D	164	ASN
1	H	117	HIS
1	H	139	GLN
1	H	143	GLN
1	H	155	HIS
1	H	156	ASN
1	H	164	ASN
2	A	86	GLN
2	A	106	HIS
2	A	124	ASN
2	A	128	GLN
2	A	143	HIS
2	A	177	GLN
2	A	180	ASN
2	A	187	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	A	197	GLN
2	A	231	GLN
2	A	322	ASN
2	A	386	GLN
2	A	398	ASN
2	A	424	GLN
2	A	426	GLN
2	A	428	HIS
2	B	86	GLN
2	B	106	HIS
2	B	124	ASN
2	B	128	GLN
2	B	143	HIS
2	B	177	GLN
2	B	180	ASN
2	B	187	GLN
2	B	197	GLN
2	B	231	GLN
2	B	266	GLN
2	B	386	GLN
2	B	398	ASN
2	B	406	GLN
2	B	424	GLN
2	B	426	GLN
2	B	428	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	1TQ	D	109	1	16,19,20	3.41	5 (31%)	11,26,28	1.89	2 (18%)
1	1TQ	H	109	1	16,19,20	3.28	5 (31%)	11,26,28	2.02	3 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	1TQ	D	109	1	-	0/6/9/11	0/2/2/2
1	1TQ	H	109	1	-	0/6/9/11	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	109	1TQ	CZ2-CE2	-3.21	1.36	1.42
1	H	109	1TQ	CZ2-CE2	-3.12	1.36	1.42
1	H	109	1TQ	CD1-CG	-2.38	1.34	1.38
1	D	109	1TQ	CD1-CG	-2.31	1.34	1.38
1	H	109	1TQ	O2-CZ2	2.34	1.44	1.35
1	D	109	1TQ	O2-CZ2	2.46	1.44	1.35
1	H	109	1TQ	CH3-N2	7.72	1.45	1.34
1	D	109	1TQ	CH2-N2	7.99	1.54	1.41
1	H	109	1TQ	CH2-N2	9.17	1.56	1.41
1	D	109	1TQ	CH3-N2	9.52	1.47	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	109	1TQ	O1-CH3-N2	-4.43	120.37	125.84
1	D	109	1TQ	O1-CH3-N2	-4.13	120.74	125.84
1	D	109	1TQ	CB-CG-CD1	-2.87	124.42	127.97
1	H	109	1TQ	CB-CG-CD1	-2.86	124.43	127.97
1	H	109	1TQ	O-C-CA	-2.24	119.66	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	109	1TQ	2	0
1	H	109	1TQ	3	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	D	123/124 (99%)	-0.17	6 (4%)	33 30	6, 10, 27, 34	0
1	H	121/124 (97%)	-0.33	2 (1%)	73 71	6, 10, 23, 29	0
2	A	360/361 (99%)	-0.61	2 (0%)	90 89	4, 8, 16, 27	0
2	B	360/361 (99%)	-0.45	2 (0%)	90 89	5, 10, 20, 29	0
All	All	964/970 (99%)	-0.46	12 (1%)	81 80	4, 9, 20, 34	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	66	LEU	5.3
2	A	73	ARG	3.8
2	B	303	ASP	3.4
1	D	70	ASP	3.4
1	D	60	ILE	3.3
2	A	303	ASP	3.1
1	D	68	ASN	2.9
1	D	63	ASN	2.8
1	H	68	ASN	2.6
1	H	60	ILE	2.4
2	B	216	VAL	2.3
1	D	67	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	1TQ	D	109	18/19	0.97	0.06	-	7,8,16,17	0
1	1TQ	H	109	18/19	0.97	0.07	-	5,8,14,14	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.