



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:20 PM GMT

PDB ID : 4I0E  
Title : Design and Synthesis of Thiophene Dihydroisoquinolins as Novel BACE-1 Inhibitors  
Authors : Yao, N.; Brecht, E.  
Deposited on : 2012-11-16  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3352 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-secretase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	393	Total	C	N	O	S	0	0	0
			3099	1977	523	585	14			


There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	56	MET	-	EXPRESSION TAG	UNP P56817
A	454	ARG	-	EXPRESSION TAG	UNP P56817
A	455	SER	-	EXPRESSION TAG	UNP P56817
A	456	HIS	-	EXPRESSION TAG	UNP P56817
A	457	HIS	-	EXPRESSION TAG	UNP P56817
A	458	HIS	-	EXPRESSION TAG	UNP P56817
A	459	HIS	-	EXPRESSION TAG	UNP P56817
A	460	HIS	-	EXPRESSION TAG	UNP P56817
A	461	HIS	-	EXPRESSION TAG	UNP P56817

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Zn	0	0
			3	3		

- Molecule 3 is 3-[2-BROMO-4-(1H-PYRAZOL-4-YL)THIOPHEN-3-YL]-N-(6-CHLORO-3,3-DIMETHYL-3,4-DIHYDROISOQUINOLIN-1-YL)-L-ALANINE (three-letter code: 1B8) (formula: C<sub>21</sub>H<sub>20</sub>BrClN<sub>4</sub>O<sub>2</sub>S).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
3	A	1	Total	Br	C	Cl	N	O	S	0	0
			30	1	21	1	4	2	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	220	Total	O	0	0
			220	220		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Beta-secretase 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.06Å 103.33Å 100.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	60.00 – 1.70 29.29 – 1.63	Depositor EDS
% Data completeness (in resolution range)	84.8 (60.00-1.70) 84.8 (29.29-1.63)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.34 (at 1.63Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.230 , 0.300 0.219 , 0.290	Depositor DCC
$R_{free}$ test set	2194 reflections (6.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.0	Xtriage
Anisotropy	0.374	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 43.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 41475 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3352	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1B8, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.07	1/3180 (0.0%)	0.99	5/4322 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	330	CYS	CB-SG	-5.76	1.72	1.81

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	180	LEU	CB-CG-CD2	-8.25	96.97	111.00
1	A	210	LEU	CA-CB-CG	7.80	133.24	115.30
1	A	284	ASP	CB-CG-OD2	6.35	124.01	118.30
1	A	277	ASP	CB-CG-OD1	5.98	123.68	118.30
1	A	273	ASP	CB-CG-OD1	5.39	123.16	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	258	TRP	Peptide

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	A	294	ASN	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3099	0	2990	38	0
2	A	3	0	0	0	0
3	A	30	0	18	1	0
4	A	220	0	0	8	0
All	All	3352	0	3008	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:368:ARG:HG2	1:A:368:ARG:HH11	1.15	1.11
1:A:421:HIS:HA	4:A:777:HOH:O	1.81	0.78
1:A:458:HIS:HB3	1:A:460:HIS:NE2	2.01	0.76
1:A:279:LYS:NZ	1:A:442:ASP:O	2.18	0.76
1:A:368:ARG:HG2	1:A:368:ARG:NH1	1.96	0.75
1:A:368:ARG:HH11	1:A:368:ARG:CG	1.99	0.75
1:A:59:SER:O	1:A:63:MET:HG3	1.88	0.73
1:A:59:SER:N	1:A:62:GLU:HB2	2.05	0.71
1:A:59:SER:H	1:A:62:GLU:HB2	1.54	0.71
1:A:382:LYS:HE2	4:A:718:HOH:O	1.93	0.67
1:A:368:ARG:HG3	1:A:384:ALA:HB2	1.87	0.57
1:A:176:TRP:HZ2	3:A:504:1B8:H17	1.70	0.56
1:A:108:PHE:CE2	1:A:172:ASN:OD1	2.58	0.56
1:A:458:HIS:HB3	1:A:460:HIS:CD2	2.42	0.55
1:A:317:LYS:HG2	4:A:814:HOH:O	2.09	0.52
1:A:353:THR:O	1:A:354:ASN:HB2	2.11	0.51
1:A:59:SER:HB3	1:A:62:GLU:H	1.76	0.51
1:A:210:LEU:HD12	1:A:211:PHE:N	2.25	0.51

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:426:PHE:N	1:A:426:PHE:CD2	2.81	0.48
1:A:361:ILE:HG22	1:A:365:GLN:HE22	1.78	0.48
1:A:58:GLY:HA3	1:A:151:GLY:O	2.14	0.48
1:A:250:TRP:O	1:A:414:GLY:HA2	2.14	0.48
1:A:368:ARG:NH1	1:A:368:ARG:CG	2.67	0.47
1:A:154:VAL:HG21	1:A:205:THR:HG21	1.97	0.46
1:A:174:SER:OG	1:A:175:ASN:N	2.48	0.45
1:A:283:TYR:HA	1:A:284:ASP:HA	1.69	0.45
1:A:263:ILE:HB	1:A:349:MET:HE2	1.98	0.44
1:A:376:SER:HB3	4:A:639:HOH:O	2.18	0.43
1:A:285:LYS:HE3	4:A:793:HOH:O	2.18	0.43
1:A:438:LEU:O	1:A:439:ASP:HB3	2.19	0.42
1:A:333:ALA:HB2	1:A:377:GLN:O	2.19	0.42
1:A:428:THR:HG22	4:A:737:HOH:O	2.19	0.42
1:A:186:GLU:OE2	1:A:256:ARG:NH1	2.50	0.41
1:A:245:TYR:CD1	1:A:403:TYR:HD2	2.38	0.41
1:A:114:GLN:HB3	1:A:117:LEU:HG	2.03	0.41
1:A:332:GLN:HB3	4:A:790:HOH:O	2.20	0.40
1:A:285:LYS:HE3	4:A:606:HOH:O	2.22	0.40
1:A:275:LYS:HB3	1:A:275:LYS:HE2	1.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	389/406 (96%)	368 (95%)	19 (5%)	2 (0%)	34 15

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	421	HIS

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	174	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	337/348 (97%)	323 (96%)	14 (4%)	36 14

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	VAL
1	A	111	ARG
1	A	136	LYS
1	A	147	SER
1	A	210	LEU
1	A	234	SER
1	A	249	LEU
1	A	258	TRP
1	A	368	ARG
1	A	373	VAL
1	A	375	THR
1	A	421	HIS
1	A	426	PHE
1	A	438	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	159	ASN
1	A	355	GLN
1	A	365	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	1B8	A	504	-	25,33,33	3.03	7 (28%)	28,49,49	4.89	11 (39%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1B8	A	504	-	-	0/8/30/30	0/4/4/4

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	504	1B8	N22-N23	-2.49	1.32	1.37
3	A	504	1B8	C14-C15	-2.05	1.49	1.51
3	A	504	1B8	C2-CL1	2.27	1.79	1.74
3	A	504	1B8	C18-C19	3.40	1.39	1.37

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	504	1B8	C5-C6	3.60	1.45	1.40
3	A	504	1B8	C5-C8	4.16	1.52	1.46
3	A	504	1B8	C19-C15	12.09	1.50	1.40

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	504	1B8	C19-C18-S17	-10.44	109.61	112.53
3	A	504	1B8	C21-C20-C19	-6.58	119.47	127.84
3	A	504	1B8	C18-C19-C20	-5.65	117.14	125.65
3	A	504	1B8	C7-C2-CL1	-2.94	115.50	119.14
3	A	504	1B8	C6-C5-C8	-2.68	115.62	118.51
3	A	504	1B8	C14-C15-C19	3.24	133.64	124.78
3	A	504	1B8	C14-C13-C25	3.38	118.81	111.55
3	A	504	1B8	N12-C8-N9	3.63	124.58	119.58
3	A	504	1B8	C21-C20-C24	4.06	113.19	104.60
3	A	504	1B8	C14-C13-N12	5.01	123.08	108.23
3	A	504	1B8	C18-C19-C15	19.45	117.05	111.69

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	504	1B8	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	393/406 (96%)	0.66	56 (14%) 4 4	20, 34, 57, 73	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	374	ALA	12.6
1	A	373	VAL	9.4
1	A	375	THR	7.2
1	A	59	SER	6.8
1	A	376	SER	6.6
1	A	426	PHE	6.0
1	A	206	HIS	5.5
1	A	438	LEU	5.2
1	A	425	GLU	4.9
1	A	439	ASP	4.7
1	A	422	VAL	4.5
1	A	421	HIS	4.0
1	A	134	GLN	4.0
1	A	61	VAL	3.9
1	A	361	ILE	3.9
1	A	171	ILE	3.8
1	A	398	ILE	3.8
1	A	60	PHE	3.7
1	A	371	GLU	3.6
1	A	108	PHE	3.6
1	A	377	GLN	3.6
1	A	359	ILE	3.6
1	A	231	VAL	3.6
1	A	172	ASN	3.2
1	A	228	LEU	3.1
1	A	174	SER	3.1
1	A	460	HIS	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	344	ILE	2.7
1	A	133	THR	2.7
1	A	334	GLY	2.7
1	A	404	VAL	2.6
1	A	395	GLY	2.6
1	A	173	GLY	2.6
1	A	62	GLU	2.6
1	A	71	SER	2.6
1	A	346	LEU	2.5
1	A	317	LYS	2.5
1	A	372	ASP	2.5
1	A	397	VAL	2.5
1	A	125	ARG	2.5
1	A	107	PRO	2.4
1	A	399	MET	2.4
1	A	58	GLY	2.4
1	A	368	ARG	2.3
1	A	402	PHE	2.3
1	A	207	VAL	2.3
1	A	415	PHE	2.3
1	A	423	HIS	2.2
1	A	90	ILE	2.2
1	A	440	MET	2.2
1	A	295	LEU	2.1
1	A	243	SER	2.1
1	A	396	ALA	2.1
1	A	288	VAL	2.1
1	A	129	TYR	2.0
1	A	332	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	1B8	A	504	30/30	0.96	0.09	-0.79	24,31,40,41	0
2	ZN	A	501	1/1	1.00	0.04	-2.58	38,38,38,38	0
2	ZN	A	502	1/1	0.96	0.07	-	67,67,67,67	0
2	ZN	A	503	1/1	0.96	0.03	-	49,49,49,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.