



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:20 PM GMT

PDB ID : 4I0G
Title : Design and Synthesis of Thiophene Dihydroisoquinolins as Novel BACE-1 Inhibitors
Authors : Yao, N.; Brecht, E.
Deposited on : 2012-11-16
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6655 (1.80-1.76)
Clashscore	102246	7658 (1.80-1.76)
Ramachandran outliers	100387	7570 (1.80-1.76)
Sidechain outliers	100360	7569 (1.80-1.76)
RSRZ outliers	91569	6671 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	<div> <div>21%</div> <div>79%</div> <div>17%</div> <div>..</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3356 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-secretase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	393	Total	C	N	O	S	0	0	0
			3099	1977	523	585	14			


There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	56	MET	-	EXPRESSION TAG	UNP P56817
A	454	ARG	-	EXPRESSION TAG	UNP P56817
A	455	SER	-	EXPRESSION TAG	UNP P56817
A	456	HIS	-	EXPRESSION TAG	UNP P56817
A	457	HIS	-	EXPRESSION TAG	UNP P56817
A	458	HIS	-	EXPRESSION TAG	UNP P56817
A	459	HIS	-	EXPRESSION TAG	UNP P56817
A	460	HIS	-	EXPRESSION TAG	UNP P56817
A	461	HIS	-	EXPRESSION TAG	UNP P56817

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Zn	0	0
			3	3		

- Molecule 3 is 3-(4-BROMOTHIOPHEN-3-YL)-N-(6-CHLORO-3,3-DIMETHYL-3,4-DIHYDROISOQUINOLIN-1-YL)-L-ALANINE (three-letter code: 1B9) (formula: C₁₈H₁₈BrClN₂O₂S).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
3	A	1	Total	Br	C	Cl	N	O	S	0	0
			25	1	18	1	2	2	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	229	Total	O	0	0
			229	229		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-secretase 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	74.77Å 103.36Å 99.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	60.00 – 1.78 22.96 – 1.78	Depositor EDS
% Data completeness (in resolution range)	95.5 (60.00-1.78) 95.5 (22.96-1.78)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.36 (at 1.78Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.232 , 0.300 0.234 , 0.298	Depositor DCC
R_{free} test set	1855 reflections (5.51%)	DCC
Wilson B-factor (Å ²)	35.8	Xtriage
Anisotropy	0.203	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	1 of 35469 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3356	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1B9, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.07	1/3180 (0.0%)	0.96	4/4322 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	281	TYR	CD2-CE2	5.46	1.47	1.39

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	277	ASP	CB-CG-OD1	6.89	124.50	118.30
1	A	157	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	A	151	GLY	N-CA-C	-5.26	99.95	113.10
1	A	295	LEU	CB-CG-CD2	-5.05	102.42	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	258	TRP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3099	0	2990	45	0
2	A	3	0	0	0	0
3	A	25	0	17	0	0
4	A	229	0	0	9	1
All	All	3356	0	3007	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59:SER:H	1:A:62:GLU:HG3	1.28	0.95
1:A:165:GLU:OE1	4:A:654:HOH:O	1.88	0.90
1:A:458:HIS:ND1	4:A:719:HOH:O	2.13	0.80
1:A:167:ASP:OD1	4:A:654:HOH:O	2.01	0.79
1:A:372:ASP:HB3	1:A:375:THR:OG1	1.88	0.73
1:A:59:SER:O	1:A:63:MET:HG3	1.89	0.72
1:A:192:ASP:OD2	4:A:719:HOH:O	2.08	0.71
1:A:136:LYS:NZ	1:A:165:GLU:OE2	2.29	0.66
1:A:101:VAL:O	1:A:101:VAL:HG23	1.94	0.65
1:A:264:ILE:HD11	1:A:394:MET:HE2	1.78	0.65
1:A:114:GLN:HG2	1:A:116:GLN:HE22	1.63	0.64
1:A:253:PRO:HG2	1:A:349:MET:CE	2.30	0.62
1:A:310:LYS:HE2	1:A:323:TRP:CD1	2.36	0.60
1:A:132:TYR:OH	4:A:624:HOH:O	2.14	0.59
1:A:58:GLY:HA2	1:A:62:GLU:HB2	1.85	0.57
1:A:253:PRO:HG2	1:A:349:MET:HE3	1.86	0.57
1:A:130:VAL:HG21	1:A:137:TRP:CZ2	2.40	0.55
1:A:114:GLN:HB3	1:A:117:LEU:HG	1.89	0.55
1:A:101:VAL:O	1:A:101:VAL:CG2	2.56	0.53
1:A:452:GLU:O	1:A:456:HIS:HD2	1.91	0.53
1:A:59:SER:H	1:A:62:GLU:CG	2.13	0.52
1:A:460:HIS:HE1	4:A:719:HOH:O	1.93	0.51
1:A:210:LEU:HD23	1:A:211:PHE:N	2.26	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:210:LEU:C	1:A:210:LEU:HD23	2.32	0.50
1:A:93:ASP:OD1	1:A:179:ILE:HD11	2.12	0.49
1:A:289:ASP:O	1:A:395:GLY:HA2	2.12	0.49
1:A:165:GLU:CD	4:A:738:HOH:O	2.51	0.48
1:A:460:HIS:CE1	4:A:719:HOH:O	2.66	0.47
1:A:97:SER:OG	1:A:187:ILE:HG13	2.15	0.46
1:A:458:HIS:O	1:A:460:HIS:N	2.45	0.45
1:A:136:LYS:HE3	1:A:167:ASP:OD2	2.17	0.45
1:A:203:LYS:HE2	1:A:203:LYS:HB3	1.85	0.45
1:A:83:SER:HA	1:A:84:PRO:C	2.37	0.44
1:A:452:GLU:O	1:A:456:HIS:CD2	2.71	0.44
1:A:363:PRO:HA	1:A:366:TYR:CE2	2.53	0.44
1:A:283:TYR:HA	1:A:284:ASP:HA	1.67	0.44
1:A:92:VAL:HG13	1:A:182:LEU:CD1	2.48	0.43
1:A:316:GLU:HA	4:A:661:HOH:O	2.17	0.43
1:A:92:VAL:HG13	1:A:182:LEU:HD11	2.00	0.43
1:A:310:LYS:HG2	1:A:323:TRP:CZ2	2.54	0.42
1:A:60:PHE:O	1:A:64:VAL:HG23	2.20	0.42
1:A:111:ARG:O	1:A:177:GLU:HG2	2.20	0.41
1:A:299:LYS:O	1:A:303:GLU:HG3	2.20	0.41
1:A:345:SER:HA	1:A:359:ILE:O	2.20	0.41
1:A:310:LYS:HE2	1:A:323:TRP:NE1	2.35	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:622:HOH:O	4:A:751:HOH:O[3_655]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	389/406 (96%)	371 (95%)	17 (4%)	1 (0%)	46 28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	440	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	337/348 (97%)	320 (95%)	17 (5%)	30 11

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	VAL
1	A	83	SER
1	A	125	ARG
1	A	134	GLN
1	A	168	LYS
1	A	171	ILE
1	A	199	ASP
1	A	249	LEU
1	A	258	TRP
1	A	375	THR
1	A	421	HIS
1	A	422	VAL
1	A	423	HIS
1	A	426	PHE
1	A	438	LEU
1	A	448	PRO
1	A	453	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	116	GLN
1	A	159	ASN
1	A	355	GLN
1	A	456	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	1B9	A	504	-	22,27,27	2.55	6 (27%)	26,40,40	11.44	8 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1B9	A	504	-	-	0/7/26/26	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	504	1B9	C20-S21	2.58	1.74	1.70
3	A	504	1B9	C6-CL1	3.14	1.81	1.74
3	A	504	1B9	BR2-C23	4.10	1.99	1.89
3	A	504	1B9	C3-C10	4.27	1.52	1.46
3	A	504	1B9	C3-C2	4.70	1.47	1.40
3	A	504	1B9	C20-C19	7.33	1.41	1.37

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	504	1B9	C23-C22-S21	-47.13	99.38	112.53
3	A	504	1B9	C19-C20-S21	-32.35	103.50	112.53
3	A	504	1B9	BR2-C23-C22	-7.32	112.89	124.40
3	A	504	1B9	C11-C8-C7	-2.54	103.19	110.78
3	A	504	1B9	C4-C5-C6	2.52	122.03	119.23
3	A	504	1B9	C1-C6-CL1	2.56	122.32	119.14
3	A	504	1B9	N13-C10-N9	2.76	123.39	119.58
3	A	504	1B9	C22-S21-C20	6.27	105.15	92.37

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	393/406 (96%)	1.12	84 (21%) 1 1	28, 42, 65, 80	0

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	373	VAL	11.1
1	A	422	VAL	10.9
1	A	374	ALA	10.0
1	A	228	LEU	8.1
1	A	59	SER	7.0
1	A	440	MET	6.4
1	A	206	HIS	6.4
1	A	61	VAL	6.2
1	A	438	LEU	5.8
1	A	426	PHE	5.6
1	A	377	GLN	5.1
1	A	229	ALA	5.0
1	A	421	HIS	5.0
1	A	182	LEU	4.8
1	A	180	LEU	4.7
1	A	60	PHE	4.7
1	A	107	PRO	4.5
1	A	77	VAL	4.4
1	A	153	ASN	4.1
1	A	172	ASN	4.0
1	A	334	GLY	4.0
1	A	404	VAL	3.9
1	A	99	PHE	3.9
1	A	376	SER	3.9
1	A	439	ASP	3.8
1	A	460	HIS	3.8
1	A	375	THR	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	425	GLU	3.6
1	A	58	GLY	3.6
1	A	213	LEU	3.4
1	A	175	ASN	3.4
1	A	171	ILE	3.3
1	A	62	GLU	3.3
1	A	217	GLY	3.2
1	A	423	HIS	3.2
1	A	385	ILE	3.2
1	A	371	GLU	3.1
1	A	133	THR	3.1
1	A	179	ILE	3.0
1	A	230	SER	3.0
1	A	211	PHE	2.9
1	A	329	VAL	2.8
1	A	344	ILE	2.8
1	A	90	ILE	2.8
1	A	92	VAL	2.8
1	A	419	ALA	2.7
1	A	110	HIS	2.7
1	A	134	GLN	2.7
1	A	348	LEU	2.6
1	A	155	THR	2.6
1	A	458	HIS	2.6
1	A	91	LEU	2.6
1	A	406	PHE	2.6
1	A	235	MET	2.6
1	A	359	ILE	2.6
1	A	398	ILE	2.5
1	A	361	ILE	2.5
1	A	94	THR	2.4
1	A	112	TYR	2.4
1	A	108	PHE	2.3
1	A	333	ALA	2.3
1	A	207	VAL	2.3
1	A	288	VAL	2.3
1	A	295	LEU	2.3
1	A	381	TYR	2.2
1	A	415	PHE	2.2
1	A	260	TYR	2.2
1	A	372	ASP	2.2
1	A	173	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	393	VAL	2.2
1	A	395	GLY	2.2
1	A	231	VAL	2.2
1	A	125	ARG	2.1
1	A	69	GLY	2.1
1	A	424	ASP	2.1
1	A	176	TRP	2.1
1	A	297	LEU	2.1
1	A	394	MET	2.1
1	A	366	TYR	2.1
1	A	246	THR	2.1
1	A	383	PHE	2.1
1	A	275	LYS	2.1
1	A	441	GLU	2.0
1	A	100	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	1B9	A	504	25/25	0.95	0.09	-1.03	35,41,55,61	0
2	ZN	A	501	1/1	0.99	0.02	-1.54	44,44,44,44	0
2	ZN	A	502	1/1	0.94	0.13	-	78,78,78,78	0
2	ZN	A	503	1/1	0.98	0.04	-	55,55,55,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.