



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 07:57 PM GMT

PDB ID : 1I1D
Title : CRYSTAL STRUCTURE OF YEAST GNA1 BOUND TO COA AND GLNAC-6P
Authors : Peneff, C.; Mengin-Lecreulx, D.; Bourne, Y.
Deposited on : 2001-02-01
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	161	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>10%</div> <div>••</div> </div> </div>
1	B	161	<div> <div>3%</div> <div> <div></div> <div>84%</div> <div>11%</div> <div>••</div> </div> </div>
1	C	161	<div> <div>4%</div> <div> <div></div> <div>74%</div> <div>21%</div> <div>••</div> </div> </div>
1	D	161	<div> <div>4%</div> <div> <div></div> <div>86%</div> <div>13%</div> <div>•</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	16G	B	906	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5713 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GLUCOSAMINE-PHOSPHATE N-ACETYLTRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	156	Total	C	N	O	S	0	4	0
			1257	804	206	237	10			
1	B	154	Total	C	N	O	S	0	4	0
			1240	792	204	234	10			
1	C	155	Total	C	N	O	S	0	1	0
			1236	789	202	234	11			
1	D	161	Total	C	N	O	S	0	2	0
			1283	814	210	248	11			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	CLONING ARTIFACT	UNP P43577
A	0	SER	-	CLONING ARTIFACT	UNP P43577
A	39	CYS	SER	ENGINEERED	UNP P43577
B	-1	GLY	-	CLONING ARTIFACT	UNP P43577
B	0	SER	-	CLONING ARTIFACT	UNP P43577
B	39	CYS	SER	ENGINEERED	UNP P43577
C	-1	GLY	-	CLONING ARTIFACT	UNP P43577
C	0	SER	-	CLONING ARTIFACT	UNP P43577
C	39	CYS	SER	ENGINEERED	UNP P43577
D	-1	GLY	-	CLONING ARTIFACT	UNP P43577
D	0	SER	-	CLONING ARTIFACT	UNP P43577
D	39	CYS	SER	ENGINEERED	UNP P43577

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE-6-PHOSPHATE) (three-letter code: 16G) (formula: C₈H₁₆NO₉P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			19	8	1	9	1		
2	B	1	Total	C	N	O	P	0	0
			19	8	1	9	1		
2	D	1	Total	C	N	O	P	0	0
			19	8	1	9	1		

- Molecule 3 is COENZYME A (three-letter code: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			45	19	7	16	3		

- Molecule 4 is IMIDAZOLE (three-letter code: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			5	3	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	152	Total	O	0	0
			152	152		
5	B	170	Total	O	0	0
			170	170		
5	C	117	Total	O	0	0
			117	117		
5	D	103	Total	O	0	0
			103	103		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: GLUCOSAMINE-PHOSPHATE N-ACETYLTRANSFERASE





- Molecule 1: GLUCOSAMINE-PHOSPHATE N-ACETYLTRANSFERASE



- Molecule 1: GLUCOSAMINE-PHOSPHATE N-ACETYLTRANSFERASE



- Molecule 1: GLUCOSAMINE-PHOSPHATE N-ACETYLTRANSFERASE



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	156.01Å 51.62Å 91.66Å 90.00° 107.98° 90.00°	Depositor
Resolution (Å)	24.12 – 1.80 35.98 – 1.80	Depositor EDS
% Data completeness (in resolution range)	96.9 (24.12-1.80) 97.0 (35.98-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	4.80	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.27 (at 1.81Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.183 , 0.223 0.184 , 0.224	Depositor DCC
R_{free} test set	1123 reflections (1.79%)	DCC
Wilson B-factor (Å ²)	21.2	Xtriage
Anisotropy	0.291	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 59.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 62765 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5713	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: COA, 16G, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	0/1298	0.88	1/1744 (0.1%)
1	B	0.82	0/1282	0.84	0/1724
1	C	0.80	1/1260 (0.1%)	0.84	0/1695
1	D	0.73	0/1313	0.83	2/1767 (0.1%)
All	All	0.80	1/5153 (0.0%)	0.85	3/6930 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	84	GLU	CB-CG	5.63	1.62	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	85	ARG	NE-CZ-NH1	5.65	123.12	120.30
1	D	158	ARG	NE-CZ-NH2	-5.62	117.49	120.30
1	A	85	ARG	NE-CZ-NH2	-5.43	117.58	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	62	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1257	0	1279	19	0
1	B	1240	0	1249	20	0
1	C	1236	0	1253	32	0
1	D	1283	0	1291	15	0
2	A	19	0	14	0	0
2	B	19	0	14	0	0
2	D	19	0	14	1	0
3	A	48	0	31	0	0
3	B	45	0	26	0	0
4	A	5	0	5	0	0
5	A	152	0	0	3	1
5	B	170	0	0	6	0
5	C	117	0	0	5	1
5	D	103	0	0	3	0
All	All	5713	0	5176	80	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:GLN:NE2	1:D:61:GLN:OE1	1.95	0.98
1:B:3:LEU:HB3	1:B:4:PRO:HD2	1.52	0.90
1:C:19:GLN:HE22	1:C:74:THR:HA	1.38	0.88
1:A:66[B]:VAL:HG13	1:A:75:VAL:HG13	1.56	0.88
1:C:19:GLN:HE22	1:C:74:THR:CA	1.92	0.83
1:B:3:LEU:HB3	1:B:4:PRO:CD	2.09	0.82
1:D:105:LYS:HG2	5:D:940:HOH:O	1.84	0.77
1:C:28:THR:HG22	1:C:29:THR:H	1.49	0.76
1:A:66[B]:VAL:HG11	1:A:75:VAL:HG22	1.67	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:130:LYS:HE2	5:C:262:HOH:O	1.87	0.74
1:B:39[B]:CYS:SG	5:B:1037:HOH:O	2.46	0.73
1:D:61:GLN:NE2	5:D:963:HOH:O	2.22	0.72
1:C:14:GLU:OE1	1:C:39[A]:CYS:SG	2.46	0.72
1:A:66[B]:VAL:CG1	1:A:75:VAL:HG13	2.20	0.70
1:A:89:HIS:HE1	2:D:908:16G:O3P	1.74	0.70
1:A:14:GLU:OE2	1:A:39:CYS:HB2	1.92	0.70
1:C:19:GLN:NE2	1:C:75:VAL:H	1.92	0.67
1:C:19:GLN:NE2	1:C:74:THR:HA	2.11	0.65
1:A:43[B]:LYS:HG2	5:A:1052:HOH:O	1.98	0.62
1:A:9:ILE:HD13	1:A:118[A]:GLN:HG2	1.81	0.62
1:B:58:LYS:HE2	1:B:60[B]:MET:CE	2.31	0.60
1:D:54:ASN:ND2	1:D:57:LYS:HD2	2.17	0.60
1:B:105:LYS:NZ	5:B:979:HOH:O	2.35	0.59
1:B:51:TRP:CZ2	1:B:53:ASP:HA	2.37	0.59
1:C:72:THR:OG1	1:C:74:THR:HB	2.03	0.58
1:C:43:LYS:O	1:C:47:GLU:HG3	2.04	0.58
1:B:89[A]:HIS:CE1	1:C:28:THR:HG21	2.40	0.57
1:A:43[B]:LYS:HE3	5:A:1052:HOH:O	2.04	0.57
1:C:28:THR:HG22	1:C:29:THR:N	2.17	0.57
1:B:14:GLU:HG3	5:B:1044:HOH:O	2.05	0.56
1:D:141:LYS:O	1:D:145:LYS:HG2	2.07	0.55
1:D:141:LYS:HD3	1:D:144:GLU:OE2	2.06	0.55
1:B:53:ASP:HB2	5:B:1038:HOH:O	2.07	0.53
1:C:86:LYS:O	1:C:91:LEU:HA	2.10	0.52
1:B:14:GLU:OE2	1:B:39[B]:CYS:SG	2.48	0.51
1:A:9:ILE:HG12	1:A:65[A]:MET:CE	2.41	0.51
1:C:50:VAL:HG13	1:C:51:TRP:N	2.25	0.51
1:C:34:THR:HB	1:C:36:GLU:OE1	2.11	0.51
1:C:140:VAL:O	1:C:144:GLU:HG3	2.11	0.51
1:D:48:ALA:HB1	1:D:61:GLN:NE2	2.27	0.49
1:B:62:TYR:OH	5:B:1073:HOH:O	2.20	0.49
1:A:26:VAL:HG12	1:A:26:VAL:O	2.13	0.49
1:C:50:VAL:CG1	1:C:51:TRP:N	2.76	0.48
1:C:83:ILE:HG22	1:C:93:LEU:HD22	1.96	0.48
1:D:36:GLU:HG3	5:D:949:HOH:O	2.12	0.48
1:B:3:LEU:C	1:B:4:PRO:O	2.50	0.47
1:A:9:ILE:HG12	1:A:65[A]:MET:HE3	1.95	0.47
1:D:67:ILE:HG21	1:D:111:LEU:HD11	1.97	0.47
1:C:150:ASN:HD22	1:C:150:ASN:C	2.17	0.46
1:D:14:GLU:OE2	1:D:39[A]:CYS:SG	2.74	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:GLU:OE2	1:A:39:CYS:CB	2.63	0.46
1:C:85:ARG:HB2	5:C:253:HOH:O	2.16	0.46
1:C:25:LYS:NZ	5:C:261:HOH:O	2.49	0.46
1:C:58:LYS:HE2	1:C:60:MET:SD	2.56	0.45
1:B:90:GLU:HG3	1:C:30:VAL:O	2.16	0.45
1:C:36:GLU:CD	1:C:36:GLU:H	2.20	0.45
1:C:7:PHE:CE1	1:C:111:LEU:HD21	2.52	0.45
1:A:86:LYS:O	1:A:91:LEU:HA	2.18	0.44
1:D:16:ASP:HB3	1:D:20:VAL:HG23	2.00	0.43
1:D:54:ASN:HD21	1:D:57:LYS:HD2	1.84	0.43
1:C:19:GLN:HE22	1:C:75:VAL:N	2.17	0.43
1:D:4:PRO:HD2	1:D:114:LEU:HD21	2.01	0.43
1:C:19:GLN:NE2	1:C:75:VAL:N	2.63	0.42
1:B:41:LEU:HD13	1:C:85:ARG:HG2	2.02	0.42
1:B:4:PRO:HD3	1:B:114:LEU:HD13	2.01	0.42
1:A:1:MET:CE	5:A:1062:HOH:O	2.67	0.42
1:B:40:LYS:NZ	5:B:975:HOH:O	2.51	0.42
1:C:129:TYR:OH	5:C:262:HOH:O	2.20	0.42
1:A:105:LYS:NZ	1:A:105:LYS:HA	2.34	0.42
1:A:150:ASN:HD22	1:A:150:ASN:C	2.22	0.42
1:D:141:LYS:HA	1:D:141:LYS:HD3	1.65	0.42
1:A:66[B]:VAL:HG11	1:A:75:VAL:CG2	2.43	0.41
1:B:97:ILE:HG23	1:B:100:ILE:HD11	2.01	0.41
1:B:89[A]:HIS:NE2	1:C:28:THR:HG21	2.36	0.41
1:C:16:ASP:HB3	1:C:20:VAL:HG23	2.03	0.41
1:C:85:ARG:CB	5:C:253:HOH:O	2.69	0.41
1:B:43:LYS:O	1:B:47:GLU:HG3	2.21	0.40
1:C:150:ASN:HD22	1:C:151:ALA:N	2.19	0.40
1:A:156:GLN:HG2	1:D:151:ALA:HB2	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1013:HOH:O	5:C:205:HOH:O[2_656]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	156/161 (97%)	154 (99%)	2 (1%)	0	100	100
1	B	154/161 (96%)	152 (99%)	2 (1%)	0	100	100
1	C	152/161 (94%)	150 (99%)	2 (1%)	0	100	100
1	D	161/161 (100%)	159 (99%)	2 (1%)	0	100	100
All	All	623/644 (97%)	615 (99%)	8 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/142 (100%)	140 (99%)	2 (1%)	74	65
1	B	140/142 (99%)	139 (99%)	1 (1%)	88	86
1	C	138/142 (97%)	135 (98%)	3 (2%)	60	45
1	D	144/142 (101%)	140 (97%)	4 (3%)	51	35
All	All	564/568 (99%)	554 (98%)	10 (2%)	70	54

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105	LYS
1	A	150	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	58	LYS
1	C	18	GLU
1	C	50	VAL
1	C	150	ASN
1	D	39[A]	CYS
1	D	39[B]	CYS
1	D	52	ASN
1	D	56	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	ASN
1	A	63	ASN
1	A	89	HIS
1	A	107	GLN
1	A	150	ASN
1	B	61	GLN
1	B	107	GLN
1	B	109	GLN
1	C	19	GLN
1	C	61	GLN
1	C	63	ASN
1	C	107	GLN
1	C	150	ASN
1	D	52	ASN
1	D	63	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	COA	A	900	-	40,50,50	1.30	4 (10%)	50,75,75	2.46	9 (18%)
2	16G	A	905	-	19,19,19	1.45	3 (15%)	23,28,28	1.53	4 (17%)
4	IMD	A	910	-	3,5,5	0.82	0	4,5,5	2.98	3 (75%)
3	COA	B	901	-	37,47,50	1.14	3 (8%)	48,72,75	1.31	5 (10%)
2	16G	B	906	-	19,19,19	1.56	3 (15%)	23,28,28	1.25	3 (13%)
2	16G	D	908	-	19,19,19	1.79	3 (15%)	23,28,28	1.64	7 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	COA	A	900	-	-	0/44/64/64	0/3/3/3
2	16G	A	905	-	-	0/10/30/30	0/1/1/1
4	IMD	A	910	-	-	0/0/0/0	0/1/1/1
3	COA	B	901	-	-	0/40/60/64	0/3/3/3
2	16G	B	906	-	-	0/10/30/30	0/1/1/1
2	16G	D	908	-	-	0/10/30/30	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	908	16G	C4-C5	-5.20	1.42	1.53
2	B	906	16G	C4-C5	-4.24	1.44	1.53
2	A	905	16G	C4-C5	-3.84	1.44	1.53
3	A	900	COA	P3B-O9A	-2.99	1.44	1.54
3	B	901	COA	C8A-N7A	-2.54	1.29	1.34
2	D	908	16G	C2-N2	-2.27	1.42	1.45
2	B	906	16G	O7-C7	-2.24	1.18	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	905	16G	P-O2P	-2.22	1.46	1.54
2	D	908	16G	P-O2P	-2.15	1.47	1.54
2	A	905	16G	C8-C7	-2.10	1.46	1.50
2	B	906	16G	C8-C7	-2.08	1.46	1.50
3	B	901	COA	C5A-C4A	-2.05	1.35	1.40
3	A	900	COA	C4A-N3A	2.01	1.38	1.35
3	B	901	COA	C4A-N3A	2.32	1.39	1.35
3	A	900	COA	O4B-C1B	2.63	1.44	1.41
3	A	900	COA	C2A-N3A	3.70	1.38	1.32

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	900	COA	C2P-C3P-N4P	-5.67	101.18	112.37
4	A	910	IMD	C4-C5-N1	-3.10	102.16	107.74
2	A	905	16G	C2-N2-C7	-3.06	115.24	123.10
2	D	908	16G	O2P-P-O6	-2.58	99.13	106.56
2	D	908	16G	C2-N2-C7	-2.54	116.57	123.10
3	A	900	COA	N3A-C2A-N1A	-2.48	127.00	128.89
3	A	900	COA	C7P-C6P-C5P	-2.47	108.24	112.31
3	B	901	COA	N3A-C2A-N1A	-2.45	127.02	128.89
4	A	910	IMD	C4-N3-C2	-2.36	98.51	105.76
2	B	906	16G	C4-C3-C2	-2.19	107.40	110.43
3	A	900	COA	O6A-CCP-CBP	-2.13	107.13	110.55
2	A	905	16G	C4-C3-C2	-2.04	107.61	110.43
2	D	908	16G	O4-C4-C3	2.01	114.86	110.34
2	D	908	16G	C6-C5-C4	2.09	116.75	112.03
2	D	908	16G	O3P-P-O6	2.12	112.68	106.56
2	B	906	16G	C6-C5-C4	2.21	117.03	112.03
3	B	901	COA	O2A-P1A-O3A	2.21	115.14	105.09
3	A	900	COA	O9A-P3B-O8A	2.55	117.09	107.38
3	B	901	COA	C4A-C5A-N7A	2.55	111.83	109.48
3	A	900	COA	O4B-C1B-N9A	2.59	113.53	108.10
3	B	901	COA	O9A-P3B-O8A	2.67	117.56	107.38
3	A	900	COA	CDP-CBP-CAP	2.82	114.50	109.34
2	A	905	16G	C6-C5-C4	2.98	118.77	112.03
2	D	908	16G	O2P-P-O1P	3.10	120.55	110.58
2	B	906	16G	C1-O5-C5	3.12	119.24	113.47
2	A	905	16G	C1-O5-C5	3.53	120.01	113.47
2	D	908	16G	C1-O5-C5	4.03	120.93	113.47
3	A	900	COA	C4B-O4B-C1B	4.42	114.58	109.72
4	A	910	IMD	C5-C4-N3	4.45	115.74	107.74

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	B	901	COA	C4B-O4B-C1B	5.28	115.53	109.72
3	A	900	COA	C3P-N4P-C5P	13.39	149.13	122.79

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	908	16G	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	156/161 (96%)	-0.18	3 (1%) 70 66	12, 20, 36, 60	0
1	B	154/161 (95%)	-0.33	5 (3%) 51 45	13, 20, 41, 59	0
1	C	155/161 (96%)	-0.22	7 (4%) 37 31	15, 26, 46, 63	0
1	D	161/161 (100%)	-0.12	6 (3%) 45 39	15, 26, 45, 65	0
All	All	626/644 (97%)	-0.21	21 (3%) 49 43	12, 23, 45, 65	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	51	TRP	6.7
1	B	53	ASP	5.5
1	D	56	ASP	4.5
1	C	58	LYS	3.9
1	B	5	ASP	3.8
1	D	-1	GLY	3.6
1	C	50	VAL	3.6
1	A	53	ASP	3.3
1	C	32	THR	3.3
1	A	1	MET	3.1
1	D	145	LYS	3.0
1	C	0	SER	3.0
1	A	57	LYS	2.9
1	D	138	LYS	2.8
1	D	141	LYS	2.8
1	B	4	PRO	2.7
1	C	14	GLU	2.7
1	B	3	LEU	2.3
1	D	55	GLU	2.2
1	B	51	TRP	2.2
1	C	35	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	16G	B	906	19/19	0.91	0.14	3.21	27,32,35,35	0
2	16G	D	908	19/19	0.88	0.13	1.55	25,34,38,40	0
3	COA	B	901	45/48	0.98	0.08	0.09	14,18,32,35	0
3	COA	A	900	48/48	0.97	0.10	-0.00	14,19,38,52	0
4	IMD	A	910	5/5	0.94	0.12	-0.06	19,20,20,21	0
2	16G	A	905	19/19	0.97	0.07	-1.03	17,22,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.