



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:20 PM GMT

PDB ID : 4I15
Title : Crystal structure of TbrPDEB1
Authors : Wang, H.; Ke, H.
Deposited on : 2012-11-20
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1226 (1.66-1.66)
Clashscore	102246	1323 (1.66-1.66)
Ramachandran outliers	100387	1295 (1.66-1.66)
Sidechain outliers	100360	1295 (1.66-1.66)
RSRZ outliers	91569	1227 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	
1	B	343	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5680 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Class 1 phosphodiesterase PDEB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	333	Total	C	N	O	S	0	0	0
			2636	1673	445	500	18			
1	B	333	Total	C	N	O	S	0	0	0
			2636	1673	445	500	18			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	244	Total	O	0	0
			244	244		
4	B	160	Total	O	0	0
			160	160		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Class 1 phosphodiesterase PDEB1



- Molecule 1: Class 1 phosphodiesterase PDEB1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	115.01Å 115.34Å 68.51Å 90.00° 108.13° 90.00°	Depositor
Resolution (Å)	30.00 – 1.65 28.57 – 1.65	Depositor EDS
% Data completeness (in resolution range)	86.4 (30.00-1.65) 86.5 (28.57-1.65)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.56 (at 1.65Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.204 , 0.223 0.204 , 0.224	Depositor DCC
R_{free} test set	8783 reflections (9.99%)	DCC
Wilson B-factor (Å ²)	21.7	Xtriage
Anisotropy	0.751	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 43.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 92150 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5680	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/2687	0.52	0/3634
1	B	0.31	0/2687	0.51	0/3634
All	All	0.31	0/5374	0.52	0/7268

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2636	0	2598	20	0
1	B	2636	0	2598	32	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	244	0	0	0	0
4	B	160	0	0	1	0
All	All	5680	0	5196	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:643:GLN:H	1:B:643:GLN:HE21	1.15	0.89
1:B:636:LEU:HD13	1:B:652:LEU:HD11	1.54	0.87
1:B:643:GLN:NE2	1:B:643:GLN:H	1.75	0.83
1:B:860:PRO:HG3	1:B:866:LYS:HZ1	1.54	0.73
1:A:590:THR:OG1	1:A:593:GLU:HG3	1.91	0.70
1:B:643:GLN:N	1:B:643:GLN:HE21	1.89	0.69
1:A:719:SER:OG	1:A:723:LYS:HE3	1.93	0.68
1:A:860:PRO:HG3	1:A:866:LYS:NZ	2.09	0.67
1:B:590:THR:OG1	1:B:593:GLU:HG3	1.96	0.65
1:B:883:ALA:HB3	1:B:884:PRO:HD3	1.77	0.65
1:B:795:PHE:CG	1:B:816:ILE:HG13	2.33	0.64
1:A:860:PRO:HA	1:A:866:LYS:HD2	1.82	0.60
1:B:595:GLU:O	1:B:599:VAL:HG23	2.01	0.60
1:B:621:THR:HG22	1:B:622:ASP:N	2.18	0.58
1:A:605:PHE:HE2	1:A:616:ARG:HE	1.52	0.57
1:B:589:ILE:HD11	1:B:658:GLN:HG2	1.89	0.55
1:A:602:LEU:N	1:A:603:PRO:HD3	2.22	0.54
1:A:771:LEU:HD23	1:A:771:LEU:C	2.29	0.53
1:A:664:ARG:NH1	1:A:714:MET:HG2	2.24	0.52
1:A:795:PHE:CG	1:A:816:ILE:HG13	2.45	0.51
1:B:621:THR:HG22	1:B:623:LYS:H	1.76	0.50
1:A:860:PRO:HG3	1:A:866:LYS:HZ2	1.75	0.50
1:A:716:LEU:O	1:A:741:LEU:HD11	2.12	0.49
1:B:589:ILE:N	1:B:589:ILE:HD12	2.27	0.49
1:B:827:THR:O	1:B:903:ARG:HG2	2.12	0.49
1:A:860:PRO:HG3	1:A:866:LYS:HZ3	1.74	0.49
1:B:869:GLU:HB2	1:B:872:LYS:HG3	1.95	0.49
1:A:847:GLN:O	1:A:851:GLU:HG3	2.12	0.49
1:B:771:LEU:C	1:B:771:LEU:HD23	2.34	0.48
1:B:716:LEU:O	1:B:741:LEU:HD11	2.15	0.47
1:A:851:GLU:HB3	1:A:856:VAL:HG23	1.96	0.47
1:B:910:GLN:O	1:B:914:VAL:HG23	2.15	0.46
1:B:910:GLN:HA	1:B:913:ARG:HE	1.80	0.46
1:A:851:GLU:HB2	1:A:858:VAL:HG22	1.97	0.46
1:A:854:ARG:HH11	1:A:854:ARG:HG2	1.80	0.46
1:A:848:GLY:O	1:A:852:LYS:HG3	2.16	0.46
1:A:605:PHE:CZ	1:A:617:ALA:HA	2.51	0.45
1:B:603:PRO:O	1:B:604:SER:OG	2.29	0.44
1:B:657:LEU:O	1:B:661:LYS:HG3	2.17	0.44
1:B:636:LEU:CD1	1:B:652:LEU:HD11	2.39	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:589:ILE:HG23	1:B:593:GLU:HB2	2.00	0.44
1:A:909:ALA:O	1:A:913:ARG:HG3	2.18	0.44
1:B:719:SER:OG	1:B:723:LYS:HE3	2.17	0.44
1:B:591:LYS:O	1:B:595:GLU:HG2	2.17	0.44
1:A:850:MET:O	1:A:854:ARG:HG3	2.18	0.44
1:B:802:GLN:C	1:B:804:SER:H	2.21	0.43
1:B:621:THR:CG2	1:B:622:ASP:N	2.82	0.43
1:B:621:THR:HB	4:B:1144:HOH:O	2.18	0.43
1:B:910:GLN:HG3	1:B:913:ARG:NH2	2.33	0.43
1:B:884:PRO:O	1:B:888:LYS:HG3	2.19	0.42
1:B:887:GLN:HE21	1:B:891:ASP:CG	2.23	0.42
1:B:847:GLN:O	1:B:851:GLU:HG3	2.20	0.42

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	331/343 (96%)	322 (97%)	7 (2%)	2 (1%)	30	9
1	B	331/343 (96%)	317 (96%)	10 (3%)	4 (1%)	16	2
All	All	662/686 (96%)	639 (96%)	17 (3%)	6 (1%)	21	4

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	600	CYS
1	B	800	ALA
1	A	893	CYS
1	A	867	ASN
1	B	893	CYS
1	B	916	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/297 (97%)	286 (99%)	2 (1%)	88	78
1	B	288/297 (97%)	285 (99%)	3 (1%)	82	68
All	All	576/594 (97%)	571 (99%)	5 (1%)	84	72

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	665	ASN
1	A	899	TRP
1	B	643	GLN
1	B	665	ASN
1	B	899	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	665	ASN
1	A	887	GLN
1	B	643	GLN
1	B	665	ASN
1	B	887	GLN
1	B	895	GLN
1	B	910	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	333/343 (97%)	0.21	23 (6%) 20 17	19, 27, 48, 58	0
1	B	333/343 (97%)	0.49	37 (11%) 7 5	20, 30, 59, 79	0
All	All	666/686 (97%)	0.35	60 (9%) 12 10	19, 29, 54, 79	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	917	THR	10.1
1	B	918	ARG	9.3
1	B	802	GLN	8.8
1	A	586	VAL	8.2
1	B	602	LEU	6.8
1	B	801	ASP	6.7
1	B	803	SER	6.0
1	B	916	GLU	5.5
1	B	603	PRO	5.5
1	B	867	ASN	5.4
1	B	804	SER	5.4
1	B	586	VAL	5.1
1	B	868	MET	4.9
1	B	601	GLU	4.9
1	B	865	SER	4.4
1	B	587	THR	4.2
1	B	599	VAL	4.1
1	A	591	LYS	3.7
1	A	677	VAL	3.5
1	A	859	LEU	3.4
1	B	592	VAL	3.3
1	A	853	GLU	3.3
1	B	604	SER	3.2
1	A	603	PRO	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	800	ALA	3.1
1	A	858	VAL	3.1
1	B	702	LEU	3.0
1	A	857	GLU	2.9
1	B	621	THR	2.9
1	B	869	GLU	2.6
1	B	704	ILE	2.6
1	A	587	THR	2.6
1	A	674	VAL	2.5
1	B	915	LEU	2.5
1	B	590	THR	2.5
1	B	598	LEU	2.5
1	A	604	SER	2.4
1	B	588	ALA	2.4
1	B	591	LYS	2.4
1	A	854	ARG	2.4
1	A	616	ARG	2.3
1	A	706	ALA	2.3
1	A	867	ASN	2.3
1	B	708	VAL	2.3
1	B	798	SER	2.3
1	A	880	PHE	2.3
1	B	595	GLU	2.2
1	A	868	MET	2.2
1	A	678	CYS	2.2
1	B	678	CYS	2.2
1	B	723	LYS	2.2
1	A	866	LYS	2.2
1	B	619	GLU	2.2
1	A	802	GLN	2.2
1	B	765	GLU	2.1
1	A	622	ASP	2.1
1	B	914	VAL	2.1
1	A	601	GLU	2.1
1	A	619	GLU	2.1
1	B	677	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	MG	B	1002	1/1	1.00	0.12	0.47	25,25,25,25	0
3	MG	A	1002	1/1	0.98	0.12	0.43	26,26,26,26	0
2	ZN	B	1001	1/1	0.99	0.06	-2.09	26,26,26,26	0
2	ZN	A	1001	1/1	1.00	0.05	-2.33	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.