



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 20, 2016 – 04:20 PM EDT

PDB ID : 5I4V
Title : Discovery of novel, orally efficacious Liver X Receptor (LXR) beta agonists
Authors : Chen, G.; McKeever, B.M.
Deposited on : 2016-02-12
Resolution : 2.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027790
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027790


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2700 (2.64-2.60)
Clashscore	102246	3065 (2.64-2.60)
Ramachandran outliers	100387	3015 (2.64-2.60)
Sidechain outliers	100360	3015 (2.64-2.60)
RSRZ outliers	91569	2706 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	270	<div> <div></div> <div>90% 6% .</div> </div>
1	E	270	<div> <div>%</div> <div>82% 8% . 9%</div> </div>
2	B	255	<div> <div>2%</div> <div>76% 7% . 16%</div> </div>
2	F	255	<div> <div></div> <div>77% 7% . 15%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7643 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Oxysterols receptor LXR-beta,Nuclear receptor coactivator 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	0	0	0
			2087	1329	368	383	7			
1	E	245	Total	C	N	O	S	0	0	0
			1993	1277	349	360	7			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	213	HIS	GLY	engineered mutation	UNP P55055
A	214	MET	GLU	engineered mutation	UNP P55055
A	259	ALA	GLN	engineered mutation	UNP P55055
A	261	GLY	ARG	engineered mutation	UNP P55055
A	262	SER	ASP	engineered mutation	UNP P55055
A	264	SER	ARG	engineered mutation	UNP P55055
A	462	GLY	-	linker	UNP P55055
A	463	SER	-	linker	UNP P55055
A	464	GLY	-	linker	UNP P55055
A	465	SER	-	linker	UNP P55055
A	466	GLY	-	linker	UNP P55055
A	467	SER	-	linker	UNP P55055
E	213	HIS	GLY	engineered mutation	UNP P55055
E	214	MET	GLU	engineered mutation	UNP P55055
E	259	ALA	GLN	engineered mutation	UNP P55055
E	261	GLY	ARG	engineered mutation	UNP P55055
E	262	SER	ASP	engineered mutation	UNP P55055
E	264	SER	ARG	engineered mutation	UNP P55055
E	462	GLY	-	linker	UNP P55055
E	463	SER	-	linker	UNP P55055
E	464	GLY	-	linker	UNP P55055
E	465	SER	-	linker	UNP P55055
E	466	GLY	-	linker	UNP P55055
E	467	SER	-	linker	UNP P55055


- Molecule 2 is a protein called Retinoic acid receptor RXR-beta, Nuclear receptor coactivator 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	215	Total	C	N	O	S	0	0	0
			1721	1106	300	305	10			
2	F	218	Total	C	N	O	S	0	0	0
			1735	1111	305	309	10			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	293	MET	GLY	engineered mutation	UNP P28702
B	529	GLY	-	linker	UNP P28702
B	530	SER	-	linker	UNP P28702
B	531	GLY	-	linker	UNP P28702
B	532	SER	-	linker	UNP P28702
B	533	GLY	-	linker	UNP P28702
B	534	SER	-	linker	UNP P28702
F	293	MET	GLY	engineered mutation	UNP P28702
F	529	GLY	-	linker	UNP P28702
F	530	SER	-	linker	UNP P28702
F	531	GLY	-	linker	UNP P28702
F	532	SER	-	linker	UNP P28702
F	533	GLY	-	linker	UNP P28702
F	534	SER	-	linker	UNP P28702

- Molecule 3 is {2-[(2R)-4-[4-(hydroxymethyl)-3-(methylsulfonyl)phenyl]-2-(propan-2-yl)piperazin-1-yl]-4-(trifluoromethyl)pyrimidin-5-yl}methanol (three-letter code: 67S) (formula: C₂₁H₂₇F₃N₄O₄S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	S	0	0
			33	21	3	4	4	1		
3	E	1	Total	C	F	N	O	S	0	0
			33	21	3	4	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	14	Total	O	0	0
			14	14		
4	B	7	Total	O	0	0
			7	7		
4	E	12	Total	O	0	0
			12	12		
4	F	8	Total	O	0	0
			8	8		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Oxysterols receptor LXR-beta,Nuclear receptor coactivator 2

Chain A: 





- Molecule 1: Oxysterols receptor LXR-beta,Nuclear receptor coactivator 2

Chain E: 





- Molecule 2: Retinoic acid receptor RXR-beta,Nuclear receptor coactivator 2

Chain B: 



- Molecule 2: Retinoic acid receptor RXR-beta,Nuclear receptor coactivator 2

Chain F: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.64Å 101.22Å 143.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.72 – 2.61 47.72 – 2.61	Depositor EDS
% Data completeness (in resolution range)	95.3 (47.72-2.61) 95.3 (47.72-2.61)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.8.0071	Depositor
R, R_{free}	0.208 , 0.248 0.210 , 0.249	Depositor DCC
R_{free} test set	1479 reflections (5.25%)	DCC
Wilson B-factor (Å ²)	50.9	Xtriage
Anisotropy	0.119	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 30.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7643	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 67S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2129	0.58	0/2880
1	E	0.39	0/2030	0.57	0/2743
2	B	0.36	0/1750	0.61	0/2356
2	F	0.37	0/1764	0.59	0/2374
All	All	0.38	0/7673	0.58	0/10353

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2087	0	2107	11	0
1	E	1993	0	2032	16	0
2	B	1721	0	1781	11	0
2	F	1735	0	1782	12	0
3	A	33	0	0	0	0
3	E	33	0	0	1	0
4	A	14	0	0	0	0
4	B	7	0	0	0	0
4	E	12	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	F	8	0	0	0	0
All	All	7643	0	7702	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:447:PHE:O	2:B:464:ARG:NH1	2.15	0.80
1:A:423:MET:HG3	2:B:468:TYR:OH	1.91	0.69
1:E:423:MET:HG3	2:F:468:TYR:OH	1.98	0.64
2:B:389:ILE:O	2:B:429:ARG:NH1	2.33	0.61
2:F:389:ILE:O	2:F:429:ARG:NH1	2.31	0.60
1:E:294:GLN:O	1:E:391:ARG:NH2	2.36	0.59
1:E:363:LEU:HB3	1:E:365:LEU:HD13	1.86	0.57
1:A:363:LEU:HB3	1:A:365:LEU:HD13	1.87	0.55
1:A:467:SER:HB3	1:E:243:PHE:CD2	2.45	0.52
2:F:413:VAL:CG1	2:F:413:VAL:O	2.58	0.51
2:B:413:VAL:O	2:B:413:VAL:CG1	2.58	0.51
1:E:407:THR:HG21	1:E:418:PHE:HB2	1.93	0.50
1:E:356:PHE:CE1	1:E:360:MET:HE1	2.47	0.49
1:E:346:GLN:HE22	1:E:438:GLN:HA	1.79	0.48
2:B:381:ILE:HA	2:B:384:PHE:CE2	2.49	0.48
2:F:337:THR:HG22	2:F:510:PHE:CZ	2.49	0.48
2:B:518:ILE:HG22	2:B:519:ASP:N	2.29	0.47
2:B:502:LYS:O	2:B:505:GLU:HB3	2.15	0.47
2:F:512:LEU:C	2:F:513:ILE:HG13	2.36	0.46
2:B:512:LEU:C	2:B:513:ILE:HG13	2.36	0.46
2:F:381:ILE:HA	2:F:384:PHE:CE2	2.50	0.46
1:E:241:ARG:HH21	1:E:284:ASP:CG	2.19	0.46
2:F:417:PHE:CE2	2:F:421:LEU:HD11	2.51	0.46
1:E:219:GLN:HG2	1:E:220:LEU:N	2.32	0.45
1:A:453:LEU:HA	1:A:456:ILE:HG22	1.99	0.45
1:E:221:THR:HG22	1:E:223:ALA:N	2.31	0.45
1:E:221:THR:HG22	1:E:223:ALA:H	1.81	0.45
1:A:221:THR:HG22	1:A:223:ALA:N	2.32	0.45
2:B:520:THR:HG23	2:B:521:PHE:N	2.33	0.44
1:A:245:ASP:O	1:A:248:LYS:HB2	2.17	0.44
1:E:423:MET:CE	1:E:423:MET:HA	2.48	0.44
2:F:481:GLU:HG2	2:F:482:GLN:HG2	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:297:GLU:HA	2:B:297:GLU:OE1	2.17	0.43
2:F:520:THR:HG23	2:F:521:PHE:N	2.33	0.43
1:E:453:LEU:HA	1:E:456:ILE:HG22	2.00	0.43
1:A:297:ARG:HD3	1:A:472:HIS:CD2	2.54	0.43
1:E:274:LEU:HB3	3:E:501:67S:C19	2.49	0.43
1:A:221:THR:HG22	1:A:223:ALA:H	1.83	0.42
2:F:297:GLU:HA	2:F:297:GLU:OE1	2.19	0.42
1:A:297:ARG:HD3	1:A:472:HIS:HD2	1.84	0.42
2:B:448:ASN:HA	2:B:464:ARG:NH1	2.35	0.42
2:F:524:GLU:CD	2:F:535:HIS:HA	2.40	0.42
1:E:408:ARG:NH2	1:E:415:GLN:O	2.49	0.42
1:E:356:PHE:CZ	1:E:360:MET:HE3	2.54	0.41
1:A:408:ARG:NH2	1:A:415:GLN:O	2.50	0.41
1:A:449:LEU:N	1:A:449:LEU:HD12	2.37	0.40
2:F:384:PHE:CD1	2:F:384:PHE:C	2.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/270 (96%)	254 (98%)	4 (2%)	0	100	100
1	E	239/270 (88%)	236 (99%)	3 (1%)	0	100	100
2	B	207/255 (81%)	202 (98%)	5 (2%)	0	100	100
2	F	210/255 (82%)	203 (97%)	7 (3%)	0	100	100
All	All	914/1050 (87%)	895 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/235 (97%)	225 (99%)	3 (1%)	76	91
1	E	219/235 (93%)	215 (98%)	4 (2%)	66	86
2	B	189/216 (88%)	184 (97%)	5 (3%)	54	79
2	F	189/216 (88%)	184 (97%)	5 (3%)	54	79
All	All	825/902 (92%)	808 (98%)	17 (2%)	61	84

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	322	HIS
1	A	408	ARG
1	A	476	GLN
2	B	345	LYS
2	B	363	LEU
2	B	384	PHE
2	B	521	PHE
2	B	542	LEU
1	E	346	GLN
1	E	361	ARG
1	E	408	ARG
1	E	453	LEU
2	F	345	LYS
2	F	384	PHE
2	F	399	THR
2	F	413	VAL
2	F	521	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	239	ASN
1	A	294	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	468	HIS
1	A	472	HIS
1	E	239	ASN
1	E	294	GLN
1	E	346	GLN
2	F	483	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	67S	A	501	-	34,35,35	2.17	4 (11%)	42,53,53	2.42	14 (33%)
3	67S	E	501	-	34,35,35	2.14	3 (8%)	42,53,53	2.52	14 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	67S	A	501	-	-	1/28/41/41	0/3/3/3
3	67S	E	501	-	-	1/28/41/41	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	67S	C18-S	-11.49	1.61	1.77
3	E	501	67S	C18-S	-11.19	1.62	1.77
3	E	501	67S	C19-S	-2.32	1.64	1.74
3	A	501	67S	C4-N1	-2.16	1.45	1.48
3	A	501	67S	C19-S	-2.13	1.65	1.74
3	E	501	67S	C13-N4	3.11	1.47	1.38
3	A	501	67S	C13-N4	3.35	1.48	1.38

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	501	67S	N3-C5-N2	-6.65	118.47	126.05
3	A	501	67S	N3-C5-N2	-5.96	119.26	126.05
3	E	501	67S	C20-C18-C16	-4.75	118.57	121.79
3	A	501	67S	C20-C18-C16	-4.72	118.59	121.79
3	E	501	67S	O4-S-O3	-4.32	108.29	117.73
3	A	501	67S	C8-C10-N3	-3.33	118.04	123.86
3	E	501	67S	C8-C10-N3	-2.93	118.74	123.86
3	A	501	67S	F1-C7-C6	-2.75	106.89	112.03
3	A	501	67S	O4-S-O3	-2.71	111.80	117.73
3	E	501	67S	C17-C16-C15	-2.70	114.69	120.15
3	E	501	67S	C3-C2-C1	-2.48	103.28	110.65
3	E	501	67S	F3-C7-C6	-2.39	107.56	112.03
3	A	501	67S	C17-C16-C15	-2.27	115.56	120.15
3	A	501	67S	O3-S-C18	-2.21	105.41	108.12
3	A	501	67S	C12-N4-C21	-2.06	108.97	113.33
3	A	501	67S	C11-N1-C4	2.23	120.36	115.50
3	A	501	67S	C11-C12-N4	2.28	114.85	110.59
3	E	501	67S	C7-C6-N2	2.29	118.88	113.00
3	E	501	67S	C5-N2-C6	2.31	119.34	116.22
3	A	501	67S	C15-C16-C18	2.39	119.80	116.04
3	A	501	67S	C17-C16-C18	2.50	124.64	122.46
3	E	501	67S	C15-C16-C18	2.68	120.25	116.04
3	E	501	67S	C11-N1-C4	2.75	121.49	115.50
3	E	501	67S	O3-S-C18	2.89	111.66	108.12
3	E	501	67S	C17-C16-C18	2.94	125.04	122.46
3	A	501	67S	O4-S-C18	2.96	111.75	108.12

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	501	67S	C10-N3-C5	8.57	124.00	115.67
3	A	501	67S	C10-N3-C5	8.69	124.12	115.67

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	501	67S	C16-C18-S-C19
3	A	501	67S	C16-C18-S-C19

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	501	67S	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	260/270 (96%)	-0.36	0 100 100	26, 39, 63, 78	0
1	E	245/270 (90%)	-0.18	2 (0%) 87 85	29, 41, 67, 80	0
2	B	215/255 (84%)	-0.12	6 (2%) 56 50	33, 52, 73, 93	0
2	F	218/255 (85%)	-0.23	1 (0%) 91 90	30, 48, 77, 90	0
All	All	938/1050 (89%)	-0.23	9 (0%) 84 81	26, 44, 70, 93	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	254	LEU	3.7
1	E	459	VAL	3.7
2	B	518	ILE	3.5
2	B	401	LEU	3.5
2	B	348	PHE	2.4
2	B	390	ASP	2.2
2	B	519	ASP	2.2
2	F	298	GLU	2.2
2	B	540	ARG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	67S	A	501	33/33	0.95	0.19	1.64	37,42,83,93	0
3	67S	E	501	33/33	0.90	0.23	1.60	41,51,109,126	0

6.5 Other polymers [i](#)

There are no such residues in this entry.