



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 07:58 PM GMT

PDB ID : 1I6L  
Title : 1.7 HIGH RESOLUTION EXPERIMENTAL PHASES FOR  
TRYPTOPHANYL-TRNA SYNTHETASE COMPLEXED WITH  
TRYPTOPHANYL-5'AMP  
Authors : Retailleau, P.; Carter, C.W.  
Deposited on : 2001-03-02  
Resolution : 1.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4425 (1.74-1.70)
Ramachandran outliers	100387	4360 (1.74-1.70)
Sidechain outliers	100360	4360 (1.74-1.70)
RSRZ outliers	91569	4010 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	<div> <div>5%</div> <div>82%</div> <div>16%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	360	-	-	-	X
2	SO4	A	364	-	-	X	-
2	SO4	A	365	-	-	-	X
5	GOL	A	370	-	X	-	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	371	-	X	-	X
5	GOL	A	372	-	X	-	-
5	GOL	A	373	-	X	-	-
5	GOL	A	374	-	X	-	-
5	GOL	A	375	-	X	-	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3064 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRYPTOPHANYL-TRNA SYNTHETASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	326	2604	1649	449	493	3	10	0	5	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP P00953
A	92	MSE	MET	MODIFIED RESIDUE	UNP P00953
A	105	MSE	MET	MODIFIED RESIDUE	UNP P00953
A	129	MSE	MET	MODIFIED RESIDUE	UNP P00953
A	184	MSE	MET	MODIFIED RESIDUE	UNP P00953
A	193	MSE	MET	MODIFIED RESIDUE	UNP P00953
A	291	MSE	MET	MODIFIED RESIDUE	UNP P00953
A	314	MSE	MET	MODIFIED RESIDUE	UNP P00953
A	318	MSE	MET	MODIFIED RESIDUE	UNP P00953
A	322	MSE	MET	MODIFIED RESIDUE	UNP P00953

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is AMMONIUM ION (three-letter code: NH4) (formula: H<sub>4</sub>N).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	N			0	0
			1	1				

- Molecule 4 is TRYPTOPHANYL-5'AMP (three-letter code: TYM) (formula: C<sub>21</sub>H<sub>24</sub>N<sub>7</sub>O<sub>8</sub>P).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	351	Total	O	0	0
			351	351		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TRYPTOPHANYL-TRNA SYNTHETASE





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.77Å 59.77Å 232.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.72 19.94 – 1.71	Depositor EDS
% Data completeness (in resolution range)	96.8 (20.00-1.72) 96.2 (19.94-1.71)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 1.71Å)	Xtriage
Refinement program	BUSTER-TNT	Depositor
R, $R_{free}$	0.209 , 0.241 0.194 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	19.0	Xtriage
Anisotropy	0.492	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 58.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 44725 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3064	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NH4, SO4, TYM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.02	3/2670 (0.1%)	0.83	3/3592 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	285	GLU	CG-CD	6.06	1.61	1.51
1	A	125	TYR	CD2-CE2	5.53	1.47	1.39
1	A	275	VAL	CB-CG1	5.41	1.64	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	117	ALA	N-CA-C	-7.13	91.75	111.00
1	A	291	MSE	CA-CB-CG	5.32	122.34	113.30
1	A	301	ASP	CB-CG-OD1	5.16	122.94	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2604	0	2621	43	0
2	A	35	0	0	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	1	0	0	0	0
4	A	37	0	23	1	0
5	A	36	0	24	2	0
6	A	351	0	0	16	0
All	All	3064	0	2668	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195[A]:LYS:HG2	5:A:374:GOL:H12	1.66	0.77
1:A:201:LYS:HG3	6:A:578:HOH:O	1.86	0.74
1:A:14:ILE:HD13	1:A:195[B]:LYS:HE2	1.73	0.69
1:A:107:GLN:HG2	6:A:467:HOH:O	1.95	0.66
1:A:228:ILE:HD12	1:A:260:TYR:HB3	1.77	0.65
1:A:285:GLU:HG2	6:A:655:HOH:O	1.98	0.64
1:A:107:GLN:HG3	1:A:150:HIS:CE1	2.33	0.63
1:A:86:HIS:HD2	1:A:132:ASP:OD2	1.82	0.62
1:A:214:GLU:O	1:A:218:LYS:HD3	2.00	0.62
1:A:112:SER:O	1:A:115:LYS:HB2	2.00	0.61
1:A:111:LYS:HD3	6:A:675:HOH:O	2.00	0.61
1:A:225:GLU:OE2	1:A:227:THR:HB	2.00	0.61
1:A:288:HIS:HE1	6:A:562:HOH:O	1.87	0.57
1:A:288:HIS:HD2	6:A:549:HOH:O	1.88	0.57
1:A:125:TYR:HB3	1:A:126:PRO:HD3	1.88	0.56
1:A:294:GLU:HG3	6:A:714:HOH:O	2.07	0.53
1:A:229:ARG:HB2	6:A:741:HOH:O	2.10	0.52
1:A:125:TYR:OH	4:A:350:TYM:NH3	2.44	0.51
1:A:86:HIS:HE1	1:A:136:TYR:OH	1.93	0.51
1:A:182:ARG:HD2	6:A:680:HOH:O	2.13	0.49
1:A:60:ARG:NH1	6:A:687:HOH:O	2.45	0.49
1:A:201:LYS:HZ1	1:A:219:SER:CB	2.26	0.48
1:A:60:ARG:NH2	1:A:291:MSE:HE2	2.28	0.48
1:A:124:THR:O	1:A:127:PRO:HD2	2.13	0.48
1:A:182:ARG:HG3	6:A:680:HOH:O	2.12	0.48
1:A:4:ILE:HD12	1:A:33:TYR:CG	2.50	0.47
1:A:60:ARG:NH2	2:A:366:SO4:S	2.89	0.45
1:A:211:LYS:NZ	6:A:588:HOH:O	2.50	0.45
1:A:259:GLN:O	1:A:263:LYS:HE3	2.17	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:373:GOL:H11	6:A:551:HOH:O	2.17	0.44
1:A:119[A]:SER:HB2	6:A:436:HOH:O	2.17	0.44
1:A:232:LYS:HE3	2:A:364:SO4:O1	2.18	0.44
1:A:252:SER:OG	1:A:255:GLU:HG3	2.18	0.44
1:A:218:LYS:HA	1:A:269:LYS:HD3	2.00	0.43
1:A:211:LYS:HG3	6:A:727:HOH:O	2.18	0.43
1:A:211:LYS:HE3	1:A:211:LYS:HB3	1.81	0.43
1:A:164:ARG:HD2	6:A:559:HOH:O	2.19	0.42
1:A:159:GLU:HG3	1:A:171:ILE:HD12	2.02	0.42
1:A:11:SER:OG	1:A:195[B]:LYS:HD3	2.20	0.42
1:A:6:SER:HB3	1:A:26:PHE:CZ	2.55	0.41
1:A:285:GLU:OE2	1:A:285:GLU:HA	2.20	0.41
1:A:232:LYS:NZ	2:A:364:SO4:O4	2.53	0.41
1:A:274:GLN:HG3	1:A:278:GLU:OE2	2.21	0.40
1:A:122:LEU:HA	1:A:122:LEU:HD23	1.82	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	329/328 (100%)	318 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	283/270 (105%)	271 (96%)	12 (4%)	36	14

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MSE
1	A	32	GLU
1	A	115	LYS
1	A	116	GLU
1	A	201	LYS
1	A	205	THR
1	A	218	LYS
1	A	225	GLU
1	A	245	TYR
1	A	254	GLU
1	A	285	GLU
1	A	295	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	GLN
1	A	86	HIS
1	A	274	GLN
1	A	288	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 1 is modelled with single atom - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	TYM	A	350	-	33,41,41	1.73	4 (12%)	31,61,61	1.53	3 (9%)
2	SO4	A	360	-	4,4,4	2.66	2 (50%)	6,6,6	0.54	0
2	SO4	A	361	-	4,4,4	2.69	2 (50%)	6,6,6	0.65	0
2	SO4	A	362	-	4,4,4	2.72	2 (50%)	6,6,6	0.68	0
2	SO4	A	363	-	4,4,4	2.66	2 (50%)	6,6,6	0.69	0
2	SO4	A	364	-	4,4,4	2.69	2 (50%)	6,6,6	0.67	0
2	SO4	A	365	-	4,4,4	2.58	2 (50%)	6,6,6	0.71	0
2	SO4	A	366	-	4,4,4	2.80	2 (50%)	6,6,6	0.67	0
5	GOL	A	370	-	5,5,5	4.63	5 (100%)	5,5,5	5.41	3 (60%)
5	GOL	A	371	-	5,5,5	4.82	4 (80%)	5,5,5	5.37	3 (60%)
5	GOL	A	372	-	5,5,5	4.78	5 (100%)	5,5,5	5.69	3 (60%)
5	GOL	A	373	-	5,5,5	4.79	5 (100%)	5,5,5	5.72	3 (60%)
5	GOL	A	374	-	5,5,5	5.00	5 (100%)	5,5,5	5.72	3 (60%)
5	GOL	A	375	-	5,5,5	4.72	5 (100%)	5,5,5	5.72	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TYM	A	350	-	-	0/16/39/39	0/5/5/5
2	SO4	A	360	-	-	0/0/0/0	0/0/0/0
2	SO4	A	361	-	-	0/0/0/0	0/0/0/0
2	SO4	A	362	-	-	0/0/0/0	0/0/0/0
2	SO4	A	363	-	-	0/0/0/0	0/0/0/0
2	SO4	A	364	-	-	0/0/0/0	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	365	-	-	0/0/0/0	0/0/0/0
2	SO4	A	366	-	-	0/0/0/0	0/0/0/0
5	GOL	A	370	-	-	0/4/4/4	0/0/0/0
5	GOL	A	371	-	-	0/4/4/4	0/0/0/0
5	GOL	A	372	-	-	0/4/4/4	0/0/0/0
5	GOL	A	373	-	-	0/4/4/4	0/0/0/0
5	GOL	A	374	-	-	0/4/4/4	0/0/0/0
5	GOL	A	375	-	-	0/4/4/4	0/0/0/0

All (47) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	371	GOL	C3-C2	-8.83	1.18	1.52
5	A	374	GOL	C3-C2	-8.43	1.20	1.52
5	A	372	GOL	C3-C2	-8.06	1.21	1.52
5	A	373	GOL	C3-C2	-8.06	1.21	1.52
5	A	370	GOL	C3-C2	-7.96	1.21	1.52
5	A	375	GOL	C3-C2	-7.88	1.22	1.52
2	A	366	SO4	O3-S	-3.97	1.33	1.47
2	A	360	SO4	O3-S	-3.89	1.33	1.47
5	A	374	GOL	C1-C2	-3.89	1.37	1.52
2	A	362	SO4	O3-S	-3.88	1.33	1.47
2	A	361	SO4	O3-S	-3.87	1.33	1.47
2	A	364	SO4	O3-S	-3.86	1.33	1.47
2	A	363	SO4	O3-S	-3.71	1.34	1.47
2	A	365	SO4	O3-S	-3.49	1.34	1.47
5	A	373	GOL	C1-C2	-3.42	1.39	1.52
5	A	371	GOL	C1-C2	-3.39	1.39	1.52
5	A	370	GOL	C1-C2	-3.32	1.39	1.52
5	A	372	GOL	C1-C2	-3.32	1.39	1.52
5	A	375	GOL	O2-C2	-2.96	1.34	1.43
5	A	373	GOL	O2-C2	-2.96	1.34	1.43
5	A	370	GOL	O2-C2	-2.90	1.34	1.43
5	A	375	GOL	C1-C2	-2.86	1.41	1.52
5	A	372	GOL	O2-C2	-2.85	1.35	1.43
5	A	374	GOL	O2-C2	-2.51	1.36	1.43
5	A	371	GOL	O2-C2	-2.47	1.36	1.43
4	A	350	TYM	C8-N7	-2.43	1.30	1.34
4	A	350	TYM	C3'-C4'	2.21	1.59	1.53
5	A	370	GOL	O3-C3	2.68	1.53	1.42
4	A	350	TYM	P-O5'	2.90	1.72	1.59
5	A	374	GOL	O3-C3	3.21	1.56	1.42

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	372	GOL	O3-C3	3.25	1.56	1.42
5	A	373	GOL	O3-C3	3.33	1.56	1.42
2	A	360	SO4	O1-S	3.47	1.59	1.47
5	A	375	GOL	O3-C3	3.48	1.57	1.42
2	A	361	SO4	O1-S	3.67	1.59	1.47
2	A	364	SO4	O1-S	3.69	1.59	1.47
2	A	363	SO4	O1-S	3.69	1.59	1.47
2	A	362	SO4	O1-S	3.72	1.59	1.47
2	A	365	SO4	O1-S	3.73	1.59	1.47
2	A	366	SO4	O1-S	3.86	1.60	1.47
5	A	370	GOL	O1-C1	4.11	1.60	1.42
5	A	371	GOL	O1-C1	4.22	1.60	1.42
5	A	373	GOL	O1-C1	4.27	1.60	1.42
5	A	372	GOL	O1-C1	4.42	1.61	1.42
5	A	375	GOL	O1-C1	4.50	1.61	1.42
5	A	374	GOL	O1-C1	4.73	1.62	1.42
4	A	350	TYM	P-OPP	7.54	1.75	1.60

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	350	TYM	CB-CG-CD1	-5.48	121.20	127.97
4	A	350	TYM	OPP-P-O5'	-3.94	92.59	102.86
4	A	350	TYM	C4'-O4'-C1'	2.23	112.17	109.72
5	A	371	GOL	O1-C1-C2	2.98	124.64	110.18
5	A	372	GOL	O1-C1-C2	3.08	125.13	110.18
5	A	370	GOL	O1-C1-C2	3.09	125.19	110.18
5	A	374	GOL	O1-C1-C2	3.14	125.42	110.18
5	A	373	GOL	O1-C1-C2	3.21	125.73	110.18
5	A	375	GOL	O1-C1-C2	3.48	127.07	110.18
5	A	371	GOL	O2-C2-C3	6.27	137.42	108.65
5	A	375	GOL	O2-C2-C3	6.51	138.52	108.65
5	A	373	GOL	O2-C2-C3	6.61	138.97	108.65
5	A	372	GOL	O2-C2-C3	6.69	139.31	108.65
5	A	374	GOL	O2-C2-C3	6.72	139.44	108.65
5	A	370	GOL	O2-C2-C3	7.06	141.03	108.65
5	A	370	GOL	O3-C3-C2	9.21	154.84	110.18
5	A	371	GOL	O3-C3-C2	9.76	157.52	110.18
5	A	372	GOL	O3-C3-C2	10.37	160.46	110.18
5	A	374	GOL	O3-C3-C2	10.41	160.67	110.18
5	A	375	GOL	O3-C3-C2	10.42	160.74	110.18
5	A	373	GOL	O3-C3-C2	10.46	160.93	110.18



There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	350	TYM	1	0
2	A	364	SO4	2	0
2	A	366	SO4	1	0
5	A	373	GOL	1	0
5	A	374	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	316/328 (96%)	0.23	17 (5%)	29 32	10, 19, 51, 79	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	114	GLY	10.5
1	A	113	ALA	9.4
1	A	115	LYS	6.8
1	A	224	SER	5.9
1	A	117	ALA	3.9
1	A	116	GLU	3.4
1	A	226	GLY	3.1
1	A	233	GLU	3.0
1	A	262	GLY	2.8
1	A	31[A]	HIS	2.5
1	A	12	GLY	2.5
1	A	106	THR	2.5
1	A	241	LEU	2.4
1	A	110	GLU	2.2
1	A	234	ALA	2.1
1	A	118	VAL	2.0
1	A	285	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	365	5/5	0.79	0.30	4.30	40,45,45,45	5
2	SO4	A	360	5/5	0.94	0.16	3.75	29,34,34,34	5
5	GOL	A	371	6/6	0.62	0.21	2.43	23,27,29,32	0
2	SO4	A	361	5/5	0.91	0.13	0.83	40,44,45,45	5
5	GOL	A	370	6/6	0.87	0.11	0.66	18,23,26,29	0
5	GOL	A	374	6/6	0.81	0.19	0.48	33,39,44,49	0
4	TYM	A	350	37/37	0.96	0.07	-0.63	9,14,20,23	0
3	NH4	A	400	1/1	0.96	0.11	-0.68	20,20,20,20	0
5	GOL	A	372	6/6	0.57	0.38	-	32,37,41,46	6
2	SO4	A	364	5/5	0.69	0.32	-	41,46,46,46	5
2	SO4	A	362	5/5	0.87	0.18	-	34,38,39,39	5
5	GOL	A	373	6/6	0.66	0.28	-	46,51,56,61	0
5	GOL	A	375	6/6	0.63	0.55	-	47,52,57,62	0
2	SO4	A	366	5/5	0.72	0.24	-	44,49,50,50	5
2	SO4	A	363	5/5	0.68	0.24	-	47,51,52,52	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.