



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:23 AM GMT

PDB ID : 3IAB
Title : Crystal structure of RNase P /RNase MRP proteins Pop6, Pop7 in a complex with the P3 domain of RNase MRP RNA
Authors : Perederina, A.; Esakova, O.A.; Quan, C.; Khanova, E.; Krasilnikov, A.S.
Deposited on : 2009-07-13
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)
RNA backbone	2183	1069 (3.10-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	158	<div> <div>13%</div> <div> <div></div> <div>79%</div> <div>14%</div> <div>• 6%</div> </div> </div>
2	B	140	<div> <div>9%</div> <div> <div></div> <div>61%</div> <div>15%</div> <div>• 24%</div> </div> </div>
3	R	46	<div> <div>9%</div> <div> <div></div> <div>33%</div> <div>48%</div> <div>15%</div> <div>•</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3087 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonucleases P/MRP protein subunit POP6.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	148	Total	C	N	O	S	Se	0	0	0
			1196	759	205	227	2	3			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	141	MSE	LEU	CONFLICT	UNP P53218

- Molecule 2 is a protein called Ribonucleases P/MRP protein subunit POP7.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	107	Total	C	N	O	S	Se	0	0	0
			853	548	151	152	1	1			

- Molecule 3 is a RNA chain called P3 domain of the RNA component of RNase MRP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	R	46	Total	C	N	O	P	0	0	0
			985	441	181	317	46			

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	3	Total	Zn	0	0
			3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	16	Total	O	0	0
			16	16		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	17	Total	O	0	0
			17	17		
5	R	17	Total	O	0	0
			17	17		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ribonucleases P/MRP protein subunit POP6



- Molecule 2: Ribonucleases P/MRP protein subunit POP7



- Molecule 3: P3 domain of the RNA component of RNase MRP



4 Data and refinement statistics

Property	Value	Source
Space group	P 42 2 2	Depositor
Cell constants a, b, c, α , β , γ	126.51Å 126.51Å 76.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.70 29.24 – 2.70	Depositor EDS
% Data completeness (in resolution range)	92.6 (30.00-2.70) 93.1 (29.24-2.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.72Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.250 , 0.265 0.252 , 0.261	Depositor DCC
R_{free} test set	822 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	74.1	Xtriage
Anisotropy	0.299	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 61.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 16350 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3087	wwPDB-VP
Average B, all atoms (Å ²)	90.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/1210	0.46	0/1619
2	B	0.29	0/864	0.47	0/1157
3	R	0.46	0/1103	0.96	5/1717 (0.3%)
All	All	0.37	0/3177	0.70	5/4493 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	R	94	A	C2'-C3'-O3'	7.18	125.30	109.50
3	R	72	U	N1-C1'-C2'	5.85	121.61	114.00
3	R	94	A	C4'-C3'-C2'	5.65	108.25	102.60
3	R	94	A	C4'-C3'-O3'	5.24	123.48	113.00
3	R	65	A	N9-C1'-C2'	5.05	120.56	114.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1196	0	1227	20	0
2	B	853	0	914	18	0
3	R	985	0	496	25	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	3	0	0	0	0
5	A	16	0	0	6	0
5	B	17	0	0	7	0
5	R	17	0	0	3	0
All	All	3087	0	2637	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88:ILE:HG13	5:A:174:HOH:O	1.59	1.02
1:A:88:ILE:N	5:A:174:HOH:O	2.05	0.87
2:B:59:GLY:HA2	5:B:149:HOH:O	1.75	0.86
1:A:52:ASN:HD22	3:R:64:C:H41	1.23	0.82
1:A:40:ASN:HB2	1:A:149:ASP:HB3	1.62	0.80
1:A:52:ASN:ND2	3:R:64:C:H41	1.80	0.79
1:A:148:ILE:HG22	1:A:150:LEU:H	1.53	0.74
3:R:59:G:N7	5:R:41:HOH:O	2.22	0.72
2:B:37:SER:O	3:R:68:A:H5'	1.90	0.71
1:A:53:ASP:O	1:A:90:LYS:HD3	1.91	0.70
1:A:51:LYS:H	1:A:87:HIS:HD2	1.40	0.70
3:R:78:G:OP2	5:R:15:HOH:O	2.11	0.69
2:B:68:MSE:HG2	3:R:91:A:H2'	1.75	0.68
3:R:72:U:O2'	3:R:73:G:OP2	2.10	0.67
1:A:84:TYR:C	5:A:174:HOH:O	2.34	0.65
3:R:93:A:O2'	3:R:95:A:OP2	2.15	0.64
3:R:66:G:H5'	3:R:67:U:OP2	1.99	0.62
1:A:108:LYS:NZ	5:A:168:HOH:O	2.32	0.62
1:A:35:ALA:HB1	1:A:150:LEU:HD22	1.84	0.59
2:B:68:MSE:HE3	2:B:131:VAL:HB	1.87	0.56
3:R:87:G:O2'	3:R:88:C:H5'	2.08	0.54
2:B:99:VAL:HG11	3:R:92:C:H42	1.72	0.54
3:R:86:A:H2'	3:R:87:G:O4'	2.09	0.52
5:B:160:HOH:O	3:R:94:A:H8	1.95	0.50
1:A:52:ASN:HD22	3:R:64:C:N4	2.03	0.50
1:A:26:LEU:O	1:A:31:ILE:HG12	2.11	0.49
1:A:84:TYR:O	5:A:174:HOH:O	2.18	0.49
2:B:31:THR:HG22	2:B:55:VAL:HG23	1.96	0.48
1:A:102:TYR:CD2	1:A:109:ILE:HG12	2.49	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:R:74:C:H2'	3:R:75:U:C6	2.48	0.48
3:R:82:A:H2'	3:R:83:C:O4'	2.14	0.48
1:A:51:LYS:H	1:A:87:HIS:CD2	2.26	0.47
2:B:14:ARG:CZ	5:B:160:HOH:O	2.62	0.47
2:B:34:PHE:H	3:R:89:U:H6	1.62	0.46
3:R:81:A:H2'	3:R:82:A:H8	1.80	0.45
3:R:59:G:C8	5:R:41:HOH:O	2.68	0.45
2:B:103:VAL:HG22	2:B:103:VAL:O	2.17	0.45
2:B:90:ILE:HG22	2:B:138:ILE:HA	1.99	0.43
2:B:58:GLN:HG2	5:B:154:HOH:O	2.16	0.43
2:B:16:THR:HG22	2:B:102:GLU:CG	2.48	0.43
1:A:143:SER:HB2	1:A:148:ILE:HD11	2.01	0.43
1:A:69:ALA:HB3	1:A:77:GLN:HG2	2.01	0.43
2:B:27:LYS:HG2	5:B:155:HOH:O	2.18	0.42
3:R:99:G:O2'	3:R:100:A:H5'	2.19	0.42
2:B:86:LYS:HE3	5:B:157:HOH:O	2.18	0.42
2:B:23:THR:N	5:B:145:HOH:O	2.52	0.42
2:B:99:VAL:HG11	3:R:92:C:N4	2.33	0.42
1:A:152:LEU:HD12	5:A:171:HOH:O	2.20	0.41
3:R:70:U:C2	3:R:74:C:H4'	2.55	0.41
3:R:81:A:H2'	3:R:82:A:C8	2.55	0.41
2:B:76:LEU:HD21	2:B:134:VAL:HG11	2.03	0.41
1:A:30:VAL:O	1:A:33:SER:HB3	2.22	0.40
3:R:80:A:H2'	3:R:81:A:O4'	2.21	0.40
2:B:46:LYS:O	2:B:50:LYS:HG2	2.21	0.40
3:R:73:G:H8	3:R:73:G:OP2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	144/158 (91%)	138 (96%)	6 (4%)	0	100	100
2	B	103/140 (74%)	100 (97%)	3 (3%)	0	100	100
All	All	247/298 (83%)	238 (96%)	9 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/145 (97%)	139 (99%)	1 (1%)	88	96
2	B	97/124 (78%)	96 (99%)	1 (1%)	82	94
All	All	237/269 (88%)	235 (99%)	2 (1%)	86	96

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	152	LEU
2	B	55	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	GLN
1	A	52	ASN
1	A	87	HIS
2	B	56	HIS
2	B	85	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	R	45/46 (97%)	12 (26%)	3 (6%)

All (12) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	R	66	G
3	R	67	U
3	R	69	A
3	R	70	U
3	R	72	U
3	R	73	G
3	R	90	U
3	R	94	A
3	R	95	A
3	R	97	U
3	R	98	G
3	R	99	G

All (3) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
3	R	65	A
3	R	72	U
3	R	94	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	145/158 (91%)	0.91	20 (13%) 4 3	54, 78, 123, 140	0
2	B	106/140 (75%)	0.94	13 (12%) 5 4	59, 88, 117, 131	0
3	R	46/46 (100%)	0.36	4 (8%) 13 10	65, 86, 146, 176	0
All	All	297/344 (86%)	0.83	37 (12%) 5 4	54, 84, 131, 176	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	14	ARG	6.0
1	A	70	ASP	5.3
1	A	129	GLN	5.0
1	A	72	SER	4.3
2	B	57	LYS	4.2
2	B	103	VAL	4.2
1	A	133	LEU	3.8
1	A	131	GLU	3.8
2	B	125	GLN	3.6
1	A	153	HIS	3.5
1	A	105	ASN	3.4
1	A	71	ARG	3.1
2	B	59	GLY	3.1
1	A	6	TYR	3.1
2	B	56	HIS	3.1
2	B	15	VAL	3.0
2	B	23	THR	2.8
1	A	106	ASN	2.8
3	R	79	G	2.7
1	A	104	LYS	2.7
3	R	82	A	2.6
1	A	151	ASN	2.5
2	B	126	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	9	GLU	2.4
1	A	130	GLU	2.4
1	A	88	ILE	2.3
2	B	134	VAL	2.3
1	A	90	LYS	2.3
1	A	120	ILE	2.2
1	A	39	ASN	2.2
3	R	59	G	2.1
3	R	78	G	2.1
1	A	73	LEU	2.1
2	B	133	GLY	2.1
1	A	40	ASN	2.0
2	B	136	LEU	2.0
2	B	77	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	ZN	B	142	1/1	0.97	0.15	-	107,107,107,107	1
4	ZN	B	143	1/1	0.98	0.09	-	82,82,82,82	1
4	ZN	B	141	1/1	0.98	0.03	-	55,55,55,55	1

6.5 Other polymers [i](#)

There are no such residues in this entry.