



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:23 AM GMT

PDB ID : 3IAG  
Title : CSL (RBP-Jk) bound to HES-1 nonconsensus site  
Authors : Friedmann, D.R.; Kovall, R.A.  
Deposited on : 2009-07-13  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	15	
2	B	15	
3	C	422	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	C	1	-	-	X	-
4	EDO	C	3	-	-	X	-
5	XYL	A	17	-	-	-	X

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4195 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(\*AP\*AP\*TP\*CP\*TP\*TP\*TP\*CP\*AP\*CP\*AP\*CP\*GP\*AP\*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	15	Total	C	N	O	P	0	0	0
			300	146	52	88	14			

- Molecule 2 is a DNA chain called 5'-D(\*TP\*TP\*AP\*TP\*CP\*GP\*TP\*GP\*TP\*GP\*AP\*AP\*AP\*GP\*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	15	Total	C	N	O	P	0	0	0
			309	149	58	88	14			

- Molecule 3 is a protein called Recombining binding protein suppressor of hairless.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	422	Total	C	N	O	S	0	8	0
			3394	2149	583	635	27			

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is SUGAR (D-XYLITOL) (three-letter code: XYL) (formula: C<sub>5</sub>H<sub>12</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			10	5	5		


- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	O	0	0
			2	2		
6	B	3	Total	O	0	0
			3	3		
6	C	153	Total	O	0	0
			153	153		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(\*AP\*AP\*TP\*CP\*TP\*TP\*TP\*CP\*AP\*CP\*AP\*CP\*GP\*AP\*T)-3'

Chain A: 





- Molecule 2: 5'-D(\*TP\*TP\*AP\*TP\*CP\*GP\*TP\*GP\*TP\*GP\*AP\*AP\*AP\*GP\*A)-3'

Chain B: 



- Molecule 3: Recombining binding protein suppressor of hairless

Chain C: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.45Å 93.14Å 112.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.80 – 2.00 24.80 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.2 (24.80-2.00) 98.2 (24.80-2.00)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.50 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.197 , 0.243 0.216 , 0.253	Depositor DCC
$R_{free}$ test set	2258 reflections (5.04%)	DCC
Wilson B-factor (Å <sup>2</sup> )	37.8	Xtriage
Anisotropy	0.187	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 48.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 44820 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4195	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XYL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.79	5/335 (1.5%)	2.69	39/514 (7.6%)
2	B	1.69	6/347 (1.7%)	2.74	41/535 (7.7%)
3	C	0.85	1/3492 (0.0%)	0.87	6/4717 (0.1%)
All	All	1.05	12/4174 (0.3%)	1.40	86/5766 (1.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	DA	C5-C4	12.47	1.47	1.38
1	A	1	DA	N9-C8	11.34	1.46	1.37
2	B	19	DT	C4-C5	10.06	1.54	1.45
2	B	19	DT	N1-C2	9.89	1.46	1.38
1	A	1	DA	N3-C4	9.70	1.40	1.34
1	A	1	DA	N1-C2	9.02	1.42	1.34
2	B	19	DT	C5-C7	8.25	1.55	1.50
1	A	1	DA	C6-N1	7.63	1.40	1.35
2	B	30	DA	O4'-C1'	6.09	1.49	1.42
2	B	19	DT	N1-C6	5.68	1.42	1.38
2	B	19	DT	C5-C6	5.48	1.38	1.34
3	C	89	GLU	CB-CG	-5.33	1.42	1.52

All (86) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	25	DG	O3'-P-O5'	-14.58	76.29	104.00
2	B	24	DT	O3'-P-O5'	-13.13	79.05	104.00
1	A	1	DA	OP1-P-O3'	-13.02	76.55	105.20
1	A	13	DG	O3'-P-O5'	-12.58	80.09	104.00
2	B	23	DG	O3'-P-O5'	-12.11	81.00	104.00
2	B	22	DT	OP1-P-O3'	-11.46	80.00	105.20
1	A	1	DA	P-O3'-C3'	10.85	132.72	119.70
2	B	17	DT	OP1-P-O3'	-10.82	81.39	105.20
1	A	2	DA	OP1-P-O3'	-10.61	81.86	105.20
1	A	6	DT	OP1-P-O3'	-10.41	82.30	105.20
1	A	3	DT	O3'-P-O5'	-10.26	84.50	104.00
2	B	26	DA	OP1-P-O3'	-10.26	82.62	105.20
3	C	218	ARG	NE-CZ-NH1	-10.21	115.19	120.30
2	B	17	DT	O3'-P-O5'	-9.59	85.78	104.00
2	B	25	DG	OP1-P-O3'	-9.56	84.16	105.20
1	A	2	DA	O3'-P-O5'	-9.42	86.11	104.00
2	B	26	DA	O3'-P-O5'	-9.36	86.22	104.00
2	B	16	DT	N3-C4-O4	9.31	125.48	119.90
2	B	20	DC	O4'-C1'-N1	-9.23	101.53	108.00
2	B	19	DT	C6-C5-C7	-9.03	117.48	122.90
1	A	3	DT	OP1-P-O3'	-9.00	85.41	105.20
2	B	16	DT	C5-C4-O4	-8.96	118.63	124.90
1	A	6	DT	P-O3'-C3'	8.89	130.37	119.70
2	B	25	DG	O4'-C1'-N9	8.17	113.72	108.00
1	A	6	DT	O3'-P-O5'	-8.07	88.67	104.00
2	B	23	DG	OP1-P-OP2	7.97	131.56	119.60
1	A	11	DA	O4'-C1'-N9	-7.80	102.54	108.00
2	B	24	DT	OP1-P-O3'	-7.76	88.14	105.20
2	B	24	DT	OP2-P-O3'	-7.62	88.44	105.20
2	B	27	DA	O4'-C1'-N9	-7.42	102.80	108.00
1	A	2	DA	O5'-P-OP1	7.36	119.53	110.70
1	A	1	DA	O3'-P-O5'	-7.32	90.09	104.00
1	A	7	DT	P-O3'-C3'	7.00	128.10	119.70
1	A	13	DG	P-O3'-C3'	6.98	128.07	119.70
1	A	14	DA	O5'-P-OP2	6.76	118.81	110.70
1	A	13	DG	OP1-P-O3'	-6.75	90.36	105.20
1	A	6	DT	C4-C5-C7	6.71	123.03	119.00
1	A	8	DC	O4'-C1'-N1	-6.66	103.34	108.00
1	A	4	DC	O4'-C1'-N1	6.61	112.63	108.00
2	B	23	DG	OP2-P-O3'	-6.53	90.83	105.20
1	A	7	DT	OP1-P-OP2	6.46	129.29	119.60
1	A	7	DT	O5'-P-OP1	6.39	118.37	110.70
1	A	3	DT	OP2-P-O3'	-6.38	91.15	105.20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	6	DT	O4'-C1'-N1	-6.36	103.55	108.00
2	B	26	DA	O5'-P-OP1	6.30	118.26	110.70
1	A	2	DA	OP1-P-OP2	6.24	128.97	119.60
2	B	19	DT	C5-C6-N1	-6.20	119.98	123.70
1	A	2	DA	OP2-P-O3'	-6.09	91.81	105.20
3	C	259	GLU	C-N-CA	6.05	136.83	121.70
3	C	326	LEU	CA-CB-CG	6.04	129.19	115.30
2	B	24	DT	O5'-P-OP1	5.99	117.88	110.70
2	B	22	DT	P-O3'-C3'	5.98	126.88	119.70
1	A	7	DT	C4-C5-C7	5.97	122.58	119.00
2	B	24	DT	O5'-P-OP2	5.93	117.82	110.70
1	A	4	DC	O5'-P-OP1	5.89	117.77	110.70
1	A	2	DA	C2-N3-C4	-5.88	107.66	110.60
2	B	18	DA	O5'-P-OP1	5.87	117.74	110.70
2	B	22	DT	O3'-P-O5'	-5.82	92.94	104.00
2	B	19	DT	C4-C5-C7	5.82	122.49	119.00
2	B	22	DT	OP2-P-O3'	-5.79	92.47	105.20
2	B	24	DT	O4'-C1'-N1	5.75	112.03	108.00
2	B	26	DA	OP2-P-O3'	-5.75	92.56	105.20
2	B	22	DT	C1'-O4'-C4'	-5.71	104.39	110.10
1	A	15	DT	C4-C5-C7	5.71	122.42	119.00
2	B	27	DA	O5'-P-OP1	5.70	117.54	110.70
2	B	25	DG	P-O3'-C3'	5.68	126.52	119.70
1	A	13	DG	OP2-P-O3'	-5.54	93.02	105.20
2	B	26	DA	P-O3'-C3'	5.50	126.31	119.70
3	C	218	ARG	NE-CZ-NH2	5.47	123.03	120.30
3	C	326	LEU	CB-CG-CD1	-5.44	101.75	111.00
1	A	15	DT	C6-C5-C7	-5.42	119.65	122.90
1	A	12	DC	O4'-C1'-N1	-5.41	104.21	108.00
3	C	89	GLU	CB-CA-C	-5.41	99.59	110.40
1	A	6	DT	OP2-P-O3'	-5.40	93.33	105.20
2	B	23	DG	OP1-P-O3'	-5.38	93.36	105.20
1	A	14	DA	OP1-P-OP2	5.37	127.66	119.60
1	A	3	DT	O5'-P-OP1	5.30	117.06	110.70
1	A	13	DG	C6-C5-N7	5.27	133.56	130.40
1	A	13	DG	O4'-C1'-C2'	-5.23	101.72	105.90
2	B	18	DA	OP1-P-OP2	5.16	127.33	119.60
2	B	17	DT	OP2-P-O3'	-5.10	93.97	105.20
2	B	27	DA	OP1-P-OP2	5.08	127.23	119.60
1	A	11	DA	N1-C2-N3	-5.08	126.76	129.30
2	B	25	DG	OP2-P-O3'	-5.08	94.03	105.20
2	B	24	DT	C5-C4-O4	-5.06	121.36	124.90

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	25	DG	O5'-P-OP2	5.03	116.73	110.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	258	GLY	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	300	0	172	5	0
2	B	309	0	172	2	0
3	C	3394	0	3391	57	0
4	A	4	0	6	0	0
4	C	20	0	30	11	0
5	A	10	0	10	0	0
6	A	2	0	0	0	0
6	B	3	0	0	0	0
6	C	153	0	0	1	0
All	All	4195	0	3781	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:134:SER:HB3	3:C:136:GLN:H	0.91	1.07
3:C:134:SER:HA	3:C:135:ASP:HB2	1.36	1.02
3:C:134:SER:HB3	3:C:136:GLN:N	1.76	1.00
3:C:136:GLN:HG3	4:C:1:EDO:O1	1.75	0.86
3:C:365:THR:H	4:C:3:EDO:H12	1.41	0.86
3:C:134:SER:HA	3:C:135:ASP:CB	2.09	0.81

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:259:GLU:N	3:C:260:GLU:CB	2.44	0.81
3:C:218:ARG:HD3	3:C:226:THR:OG1	1.82	0.80
3:C:230:HIS:HD2	3:C:232:GLU:OE2	1.66	0.78
3:C:134:SER:CB	3:C:136:GLN:H	1.85	0.78
3:C:131:ILE:CD1	4:C:1:EDO:H11	2.14	0.77
3:C:258:GLY:O	3:C:260:GLU:HB3	1.87	0.74
3:C:94[A]:CYS:HG	3:C:228:TYR:HH	1.29	0.74
3:C:131:ILE:HD11	4:C:1:EDO:H11	1.68	0.74
3:C:259:GLU:H	3:C:260:GLU:CB	2.02	0.72
1:A:14:DA:H2''	1:A:15:DT:H5'	1.74	0.70
3:C:259:GLU:H	3:C:260:GLU:HB2	1.56	0.70
3:C:365:THR:H	4:C:3:EDO:C1	2.06	0.68
3:C:136:GLN:CG	4:C:1:EDO:O1	2.45	0.65
3:C:259:GLU:N	3:C:260:GLU:HB3	2.10	0.65
3:C:422:ARG:HH11	3:C:422:ARG:CG	2.10	0.64
3:C:258:GLY:C	3:C:260:GLU:HB3	2.20	0.62
3:C:131:ILE:HD11	4:C:1:EDO:C1	2.29	0.61
2:B:25:DG:H2''	2:B:26:DA:C8	2.38	0.57
3:C:131:ILE:HD13	4:C:1:EDO:H11	1.86	0.57
3:C:259:GLU:N	3:C:260:GLU:HB2	2.16	0.55
3:C:150:THR:OG1	3:C:152:LYS:HE3	2.07	0.54
3:C:134:SER:CA	3:C:135:ASP:HB2	2.24	0.54
3:C:275[A]:LYS:HE3	3:C:284:ALA:HB1	1.91	0.53
3:C:295:LYS:HB2	6:C:531:HOH:O	2.08	0.53
1:A:14:DA:C2'	1:A:15:DT:H5'	2.40	0.52
1:A:1:DA:H2''	1:A:2:DA:C8	2.45	0.51
3:C:79:ALA:HB2	3:C:95:PRO:HD2	1.93	0.50
1:A:4:DC:OP2	3:C:196:LYS:HE2	2.12	0.49
3:C:382:VAL:HG22	4:C:3:EDO:O1	2.12	0.49
3:C:216:PHE:HB3	3:C:228:TYR:CD2	2.48	0.49
3:C:78:HIS:HD2	3:C:79:ALA:O	1.97	0.48
3:C:60:GLU:HB2	4:C:4:EDO:H21	1.96	0.48
3:C:198:GLN:HG3	3:C:307:SER:HB3	1.95	0.47
3:C:261:PHE:HB3	3:C:262:THR:H	1.52	0.47
3:C:132:GLY:O	3:C:133:ASN:OD1	2.32	0.47
3:C:56:ARG:NH2	3:C:457:ASN:OD1	2.48	0.46
3:C:422:ARG:NH1	3:C:428:LEU:HD12	2.31	0.46
3:C:422:ARG:NH1	3:C:422:ARG:CG	2.75	0.46
3:C:196:LYS:HA	3:C:196:LYS:HD3	1.90	0.45
3:C:219:LEU:O	3:C:219:LEU:HD12	2.17	0.45
3:C:422:ARG:NH1	3:C:422:ARG:HG2	2.34	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:258:GLY:C	3:C:260:GLU:CB	2.82	0.43
3:C:198:GLN:HG3	3:C:307:SER:CB	2.47	0.43
3:C:365:THR:N	4:C:3:EDO:H12	2.21	0.43
3:C:78:HIS:CE1	3:C:184:SER:HB2	2.54	0.43
3:C:422:ARG:HH11	3:C:422:ARG:HG2	1.80	0.42
1:A:4:DC:P	3:C:196:LYS:HE2	2.59	0.42
3:C:260:GLU:HG2	3:C:261:PHE:N	2.35	0.42
2:B:29:DG:C6	2:B:30:DA:C6	3.07	0.42
3:C:89:GLU:OE1	3:C:91:ARG:NH2	2.52	0.42
3:C:389:ASN:HB2	3:C:390:GLY:C	2.40	0.41
3:C:150:THR:OG1	3:C:152:LYS:CE	2.68	0.41
3:C:396:MET:SD	3:C:430:VAL:HG22	2.60	0.41
3:C:390:GLY:N	3:C:391:GLY:HA2	2.36	0.41
3:C:75:LEU:HB2	3:C:100:TYR:HB2	2.03	0.41
3:C:219:LEU:HD11	3:C:220:ARG:NH1	2.37	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	C	428/422 (101%)	412 (96%)	10 (2%)	6 (1%)	<div>14</div> <div>6</div>

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	260	GLU
3	C	261	PHE
3	C	262	THR
3	C	135	ASP
3	C	134	SER
3	C	391	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	380/372 (102%)	377 (99%)	3 (1%)	86	89

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	221	SER
3	C	420	MET
3	C	422	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
3	C	64	ASN
3	C	72	GLN
3	C	78	HIS
3	C	83	GLN
3	C	230	HIS
3	C	335	GLN
3	C	407	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	EDO	A	16	-	3,3,3	0.92	0	2,2,2	0.86	0
5	XYL	A	17	-	9,9,9	0.65	0	10,11,11	1.46	1 (10%)
4	EDO	C	1	-	3,3,3	0.35	0	2,2,2	0.79	0
4	EDO	C	2	-	3,3,3	0.39	0	2,2,2	0.96	0
4	EDO	C	3	-	3,3,3	0.34	0	2,2,2	0.83	0
4	EDO	C	4	-	3,3,3	0.43	0	2,2,2	0.65	0
4	EDO	C	6	-	3,3,3	0.60	0	2,2,2	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	16	-	-	0/1/1/1	0/0/0/0
5	XYL	A	17	-	-	0/12/12/12	0/0/0/0
4	EDO	C	1	-	-	0/1/1/1	0/0/0/0
4	EDO	C	2	-	-	0/1/1/1	0/0/0/0
4	EDO	C	3	-	-	0/1/1/1	0/0/0/0
4	EDO	C	4	-	-	0/1/1/1	0/0/0/0
4	EDO	C	6	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	17	XYL	C1-C2-C3	-2.80	105.90	112.48

There are no chirality outliers.

There are no torsion outliers.



There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	1	EDO	6	0
4	C	3	EDO	4	0
4	C	4	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	15/15 (100%)	0.11	0	100 100	39, 46, 55, 67	0
2	B	15/15 (100%)	0.69	0	100 100	33, 46, 56, 66	0
3	C	422/422 (100%)	0.94	80 (18%)	2 2	19, 31, 53, 69	0
All	All	452/452 (100%)	0.91	80 (17%)	2 2	19, 32, 55, 69	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	391	GLY	7.5
3	C	135	ASP	7.2
3	C	393	ASP	6.2
3	C	263	VAL	5.8
3	C	474	PRO	5.6
3	C	343	GLN	4.9
3	C	340	PRO	4.8
3	C	454	LEU	4.5
3	C	200	LEU	4.4
3	C	366	PHE	4.4
3	C	410	VAL	4.4
3	C	254	ASP	4.3
3	C	341	LYS	4.3
3	C	314	PHE	4.1
3	C	412	PHE	4.1
3	C	394	VAL	4.1
3	C	252	ASP	4.1
3	C	261	PHE	3.9
3	C	253	ASP	3.9
3	C	390	GLY	3.8
3	C	453	THR	3.8
3	C	380	VAL	3.7
3	C	201	LYS	3.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	C	74	VAL	3.7
3	C	204	ASP	3.6
3	C	342	GLU	3.5
3	C	219	LEU	3.5
3	C	262	THR	3.5
3	C	455	VAL	3.3
3	C	260	GLU	3.3
3	C	354	TRP	3.3
3	C	339	CYS	3.3
3	C	356	ILE	3.2
3	C	462	TYR	3.2
3	C	294	ASP	3.2
3	C	411	TRP	3.2
3	C	328	GLN	3.1
3	C	461	ILE	3.1
3	C	452	VAL	3.1
3	C	199	SER	3.1
3	C	203	ALA	3.0
3	C	169	VAL	3.0
3	C	137	GLU	3.0
3	C	258	GLY	2.9
3	C	312[A]	CYS	2.9
3	C	473	GLU	2.9
3	C	53	PRO	2.9
3	C	215	LEU	2.9
3	C	195	LYS	2.8
3	C	276	LEU	2.8
3	C	290	ILE	2.8
3	C	392	GLY	2.7
3	C	164[A]	HIS	2.7
3	C	182	PHE	2.6
3	C	289	ILE	2.6
3	C	259	GLU	2.6
3	C	229	LEU	2.5
3	C	202	ASN	2.4
3	C	136	GLN	2.4
3	C	295	LYS	2.4
3	C	134	SER	2.4
3	C	99	VAL	2.3
3	C	460	VAL	2.3
3	C	377	VAL	2.3
3	C	301	ASP	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	C	413	GLY	2.3
3	C	255	GLU	2.2
3	C	365	THR	2.2
3	C	214	ALA	2.2
3	C	313	ALA	2.2
3	C	265	ASP	2.2
3	C	196	LYS	2.2
3	C	415	VAL	2.1
3	C	285	LEU	2.1
3	C	389	ASN	2.1
3	C	88	ASN	2.1
3	C	288	LEU	2.1
3	C	68	GLU	2.0
3	C	324	LEU	2.0
3	C	216	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	XYL	A	17	10/10	0.84	0.28	4.02	74,77,78,78	0
4	EDO	C	2	4/4	0.86	0.26	1.39	44,50,53,55	0
4	EDO	C	3	4/4	0.93	0.26	0.78	43,46,50,53	0
4	EDO	C	6	4/4	0.87	0.22	-0.07	63,64,64,65	0
4	EDO	C	1	4/4	0.94	0.15	-0.53	48,50,51,51	0
4	EDO	A	16	4/4	0.79	0.18	-	84,84,85,85	0
4	EDO	C	4	4/4	0.79	0.41	-	63,68,69,70	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.