



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:26 PM GMT

PDB ID : 4IB5
Title : Structure of human protein kinase CK2 catalytic subunit in complex with a CK2beta-competitive cyclic peptide
Authors : Raaf, J.; Guerra, B.; Neundorff, I.; Bopp, B.; Issinger, O.-G.; Jose, J.; Pietsch, M.; Niefind, K.
Deposited on : 2012-12-08
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	<div> <div>2%</div> <div>87%</div> <div>11%</div> <div>..</div> </div>
1	B	335	<div> <div>8%</div> <div>91%</div> <div>7%</div> <div>..</div> </div>
1	C	335	<div> <div>7%</div> <div>89%</div> <div>10%</div> <div>.</div> </div>
2	D	13	<div> <div>100%</div> </div>
2	E	13	<div> <div>54%</div> <div>77%</div> <div>23%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	13	
2	G	13	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	404	-	-	-	X
3	GOL	B	401[A]	-	-	-	X
3	GOL	B	401[B]	-	-	-	X
3	GOL	C	403	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9387 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Casein kinase II subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	331	Total	C	N	O	S	0	11	0
			2882	1844	510	517	11			
1	B	329	Total	C	N	O	S	0	0	0
			2782	1783	489	499	11			
1	C	332	Total	C	N	O	S	0	0	0
			2804	1795	495	503	11			

- Molecule 2 is a protein called CK2beta-derived cyclic peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	13	Total	C	N	O	S	0	0	0
			97	62	19	14	2			
2	E	13	Total	C	N	O	S	0	0	0
			97	62	19	14	2			
2	F	13	Total	C	N	O	S	0	0	0
			97	62	19	14	2			
2	G	13	Total	C	N	O	S	0	0	0
			97	62	19	14	2			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	1
			12	6	6		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	1
			12	6	6		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Cl	0	0
			2	2		
4	C	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	190	Total 190	O 190	0	0
5	B	106	Total 106	O 106	0	0
5	C	154	Total 154	O 154	0	0
5	D	7	Total 7	O 7	0	0
5	E	4	Total 4	O 4	0	0
5	G	1	Total 1	O 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Casein kinase II subunit alpha



- Molecule 1: Casein kinase II subunit alpha



- Molecule 1: Casein kinase II subunit alpha




- Molecule 2: CK2beta-derived cyclic peptide




There are no outlier residues recorded for this chain.


- Molecule 2: CK2beta-derived cyclic peptide



- Molecule 2: CK2beta-derived cyclic peptide



- Molecule 2: CK2beta-derived cyclic peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	77.38Å 105.42Å 152.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.69 – 2.20 45.69 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.6 (45.69-2.20) 99.6 (45.69-2.20)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.177 , 0.218 0.177 , 0.219	Depositor DCC
R_{free} test set	3183 reflections (4.99%)	DCC
Wilson B-factor (Å ²)	39.1	Xtriage
Anisotropy	0.417	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 63978 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9387	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/2961	0.51	0/4000
1	B	0.23	0/2857	0.47	1/3864 (0.0%)
1	C	0.24	0/2879	0.51	1/3893 (0.0%)
2	D	0.24	0/99	0.37	0/129
2	E	0.25	0/99	0.36	0/129
2	F	0.22	0/99	0.36	0/129
2	G	0.22	0/99	0.31	0/129
All	All	0.24	0/9093	0.49	2/12273 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	48	GLY	N-CA-C	5.73	127.43	113.10
1	C	71	LYS	C-N-CD	5.67	140.31	128.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2882	0	2829	25	1
1	B	2782	0	2725	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2804	0	2748	20	0
2	D	97	0	92	0	0
2	E	97	0	92	2	0
2	F	97	0	92	2	0
2	G	97	0	92	1	0
3	A	36	0	48	6	0
3	B	12	0	16	0	0
3	C	18	0	24	1	0
4	A	2	0	0	1	0
4	C	1	0	0	0	0
5	A	190	0	0	3	0
5	B	106	0	0	0	0
5	C	154	0	0	3	0
5	D	7	0	0	0	0
5	E	4	0	0	0	0
5	G	1	0	0	0	0
All	All	9387	0	8758	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:174:ILE:HB	3:C:402:GOL:H32	1.69	0.73
1:A:278:ARG:NH2	1:A:302:ASP:OD1	2.24	0.70
1:A:47[B]:ARG:NH1	5:A:574:HOH:O	2.24	0.69
1:B:132:ASP:OD1	1:B:169:ARG:NH2	2.26	0.68
1:A:43[A]:ARG:NH1	1:A:55:GLU:OE1	2.27	0.67
1:C:129:THR:HG23	1:C:132:ASP:H	1.60	0.67
1:A:186[B]:GLN:NE2	5:A:578:HOH:O	2.32	0.62
1:C:47:ARG:HG3	1:C:48:GLY:H	1.68	0.59
1:C:278:ARG:NH2	5:C:528:HOH:O	2.37	0.57
1:A:174:ILE:HD12	3:A:401[B]:GOL:H32	1.87	0.56
1:A:174:ILE:HD12	3:A:401[A]:GOL:H32	1.89	0.55
1:A:84:ILE:HG23	1:A:152:ILE:HD13	1.88	0.55
1:C:116:VAL:HG21	1:C:172:ARG:HG3	1.90	0.54
1:C:259:ASP:OD1	5:C:649:HOH:O	2.19	0.54
2:F:186:ARG:NH1	2:F:189:GLY:O	2.40	0.53
1:A:70:LEU:HD13	1:A:78:ILE:HG12	1.91	0.53
1:A:286:HIS:CE1	1:A:288:GLU:HB2	2.44	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:64:LYS:O	1:A:115:HIS:HB2	2.09	0.52
1:B:134:ARG:HG2	1:B:323:TYR:CZ	2.46	0.51
1:B:48:GLY:O	1:B:49:LYS:HB2	2.10	0.51
1:C:134:ARG:HG2	1:C:323:TYR:CZ	2.46	0.51
1:A:52[A]:GLU:HG3	1:A:54:PHE:HE2	1.76	0.50
1:B:71:LYS:O	1:B:71:LYS:HG3	2.11	0.50
1:B:159:PRO:HD3	1:B:221:MET:HG2	1.94	0.49
1:A:134:ARG:HG2	1:A:323:TYR:CZ	2.48	0.48
1:B:47:ARG:HG2	1:B:53:VAL:HG21	1.94	0.48
1:A:172:ARG:HH21	3:A:404:GOL:H32	1.78	0.48
1:C:66:VAL:HG23	1:C:115:HIS:HA	1.95	0.47
1:C:44:LYS:HZ1	1:C:47:ARG:NE	2.11	0.47
1:C:165:ASP:HB3	1:C:170:LYS:HG2	1.95	0.47
1:B:290:GLN:HA	1:B:293:VAL:HG12	1.97	0.47
1:A:159:PRO:HD3	1:A:221:MET:HG2	1.97	0.47
1:A:51[A]:SER:HB3	1:A:70:LEU:HD23	1.97	0.47
1:B:40:GLN:HG2	2:E:191:LYS:HB2	1.97	0.46
1:A:48[A]:GLY:O	1:A:49[A]:LYS:HB2	2.16	0.45
1:A:96:THR:H	3:A:404:GOL:H12	1.81	0.45
1:A:45[B]:LEU:HD21	1:A:55:GLU:HB2	1.99	0.44
1:A:148:HIS:ND1	4:A:406:CL:CL	2.80	0.44
1:C:84:ILE:HG23	1:C:152:ILE:HD13	1.99	0.44
1:B:281:TRP:HB3	1:B:298:LEU:HD22	1.99	0.44
1:A:132:ASP:OD2	5:A:673:HOH:O	2.21	0.44
1:C:130:ASP:O	1:C:134:ARG:HG3	2.18	0.43
1:C:49:LYS:HA	1:C:49:LYS:HD2	1.78	0.43
2:G:191:LYS:HB3	2:G:191:LYS:HE3	1.73	0.43
1:A:73:VAL:HG13	1:A:78:ILE:CD1	2.49	0.43
1:C:108:THR:HG21	2:F:188:TYR:CD1	2.54	0.43
1:C:134:ARG:HD3	1:C:323:TYR:O	2.19	0.43
1:A:174:ILE:HB	3:A:401[A]:GOL:H11	2.00	0.43
1:C:230:GLU:HA	1:C:231:PRO:HA	1.91	0.42
1:B:153:MET:HE3	1:B:182:TYR:HB2	2.00	0.42
1:C:303:LYS:HB3	1:C:313:LEU:HG	2.00	0.42
2:E:185:CYS:SG	2:E:196:GLY:N	2.92	0.42
1:B:130:ASP:O	1:B:134:ARG:HG3	2.20	0.41
1:B:84:ILE:HG12	1:B:152:ILE:HD12	2.02	0.41
1:C:73:VAL:HG13	1:C:74:LYS:HG2	2.02	0.41
1:A:44:LYS:O	3:A:402:GOL:H12	2.21	0.41
1:C:103:ASP:OD1	1:C:106:SER:HB3	2.21	0.41
1:A:47[A]:ARG:O	1:A:51[A]:SER:O	2.39	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:153:MET:CE	1:B:182:TYR:HB2	2.51	0.40
1:B:145:ASP:OD2	1:B:316:ARG:NH2	2.53	0.40
1:A:132:ASP:OD1	1:A:169:ARG:NH1	2.54	0.40
1:C:186:GLN:NE2	5:C:627:HOH:O	2.54	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:ARG:NH2	1:A:120:ASP:OD2[4_545]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/335 (102%)	320 (94%)	18 (5%)	2 (1%)	30	29
1	B	327/335 (98%)	311 (95%)	15 (5%)	1 (0%)	46	50
1	C	330/335 (98%)	316 (96%)	13 (4%)	1 (0%)	46	50
2	D	11/13 (85%)	11 (100%)	0	0	100	100
2	E	11/13 (85%)	11 (100%)	0	0	100	100
2	F	11/13 (85%)	10 (91%)	1 (9%)	0	100	100
2	G	11/13 (85%)	11 (100%)	0	0	100	100
All	All	1041/1057 (98%)	990 (95%)	47 (4%)	4 (0%)	46	42

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	49	LYS
1	A	50[A]	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	50[B]	TYR
1	C	49	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/307 (102%)	307 (98%)	6 (2%)	65	77
1	B	303/307 (99%)	300 (99%)	3 (1%)	82	91
1	C	305/307 (99%)	302 (99%)	3 (1%)	82	91
2	D	9/9 (100%)	9 (100%)	0	100	100
2	E	9/9 (100%)	9 (100%)	0	100	100
2	F	9/9 (100%)	9 (100%)	0	100	100
2	G	9/9 (100%)	8 (89%)	1 (11%)	8	6
All	All	957/957 (100%)	944 (99%)	13 (1%)	76	85

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	ARG
1	A	43[A]	ARG
1	A	43[B]	ARG
1	A	121	PHE
1	A	153	MET
1	A	228	ARG
1	B	71	LYS
1	B	120	ASP
1	B	283	ARG
1	C	49	LYS
1	C	57	ILE
1	C	120	ASP
2	G	193	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such

sidechains are listed below:

Mol	Chain	Res	Type
1	B	310	GLN
1	C	166	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 3 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	401[A]	-	5,5,5	0.35	0	5,5,5	0.29	0
3	GOL	A	401[B]	-	5,5,5	0.33	0	5,5,5	0.28	0
3	GOL	A	402	-	5,5,5	0.37	0	5,5,5	0.32	0
3	GOL	A	403	-	5,5,5	0.33	0	5,5,5	0.23	0
3	GOL	A	404	-	5,5,5	0.36	0	5,5,5	0.42	0
3	GOL	A	405	-	5,5,5	0.35	0	5,5,5	0.32	0
3	GOL	B	401[A]	-	5,5,5	0.36	0	5,5,5	0.21	0
3	GOL	B	401[B]	-	5,5,5	0.34	0	5,5,5	0.24	0
3	GOL	C	401	-	5,5,5	0.34	0	5,5,5	0.25	0
3	GOL	C	402	-	5,5,5	0.35	0	5,5,5	0.27	0
3	GOL	C	403	-	5,5,5	0.35	0	5,5,5	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	401[A]	-	-	0/4/4/4	0/0/0/0
3	GOL	A	401[B]	-	-	0/4/4/4	0/0/0/0
3	GOL	A	402	-	-	0/4/4/4	0/0/0/0
3	GOL	A	403	-	-	0/4/4/4	0/0/0/0
3	GOL	A	404	-	-	0/4/4/4	0/0/0/0
3	GOL	A	405	-	-	0/4/4/4	0/0/0/0
3	GOL	B	401[A]	-	-	0/4/4/4	0/0/0/0
3	GOL	B	401[B]	-	-	0/4/4/4	0/0/0/0
3	GOL	C	401	-	-	0/4/4/4	0/0/0/0
3	GOL	C	402	-	-	0/4/4/4	0/0/0/0
3	GOL	C	403	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401[A]	GOL	2	0
3	A	401[B]	GOL	1	0
3	A	402	GOL	1	0
3	A	404	GOL	2	0
3	C	402	GOL	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	331/335 (98%)	-0.23	6 (1%) 71 70	27, 50, 84, 106	0
1	B	329/335 (98%)	0.16	26 (7%) 15 15	41, 74, 112, 143	0
1	C	332/335 (99%)	0.16	25 (7%) 17 17	32, 58, 128, 180	0
2	D	13/13 (100%)	-0.02	0 100 100	50, 74, 109, 113	0
2	E	13/13 (100%)	2.39	7 (53%) 0 0	70, 93, 123, 125	0
2	F	13/13 (100%)	4.99	12 (92%) 0 0	102, 127, 157, 168	0
2	G	13/13 (100%)	0.42	2 (15%) 3 2	64, 82, 94, 99	0
All	All	1044/1057 (98%)	0.13	78 (7%) 17 17	27, 62, 120, 180	0

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	48[A]	GLY	9.0
2	F	185	CYS	8.8
1	C	48	GLY	8.5
2	F	196	GLY	8.1
2	F	195	CYS	7.6
1	C	50	TYR	7.3
1	C	45	LEU	6.2
1	B	18	HIS	6.0
2	F	187	LEU	5.9
1	C	43	ARG	5.9
1	C	42	VAL	5.8
1	C	49	LYS	5.7
2	E	184	GLY	5.6
1	C	41	LEU	5.5
1	A	49[A]	LYS	5.4
2	E	196	GLY	5.2
1	A	50[A]	TYR	5.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	E	185	CYS	5.2
1	A	47[A]	ARG	5.1
1	B	329	LYS	5.1
2	E	194	GLY	5.1
2	F	190	PHE	5.0
1	B	331	GLN	5.0
2	F	188	TYR	4.8
2	F	194	GLY	4.7
2	F	192	ILE	4.5
2	F	184	GLY	4.4
1	C	54	PHE	4.3
1	C	119	THR	4.3
1	B	15	VAL	4.2
1	B	123	GLN	4.0
2	F	193	HIS	3.8
1	C	62	ASN	3.7
1	B	48	GLY	3.7
2	F	191	LYS	3.6
1	C	70	LEU	3.6
2	F	186	ARG	3.5
2	E	195	CYS	3.5
1	C	332	ALA	3.4
2	G	196	GLY	3.4
1	C	73	VAL	3.3
1	B	169	ARG	3.2
1	C	333	ARG	3.2
1	C	59	ILE	3.2
1	B	327	VAL	3.1
1	B	316	ARG	3.1
1	B	49	LYS	3.0
1	C	72	PRO	3.0
1	B	170	LYS	2.9
2	E	193	HIS	2.9
1	C	125	TYR	2.9
1	C	55	GLU	2.8
1	B	17	THR	2.8
1	C	53	VAL	2.7
1	C	103	ASP	2.7
1	B	16	ASN	2.7
1	C	47	ARG	2.7
1	B	166	HIS	2.7
1	A	117	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	119	THR	2.6
1	B	19	ARG	2.5
1	B	330	ASP	2.5
1	B	135	PHE	2.5
1	C	71	LYS	2.4
1	A	46[A]	GLY	2.4
1	B	121	PHE	2.3
1	B	50	TYR	2.3
1	C	51	SER	2.2
1	B	11	VAL	2.2
1	B	14	ASP	2.1
2	E	192	ILE	2.1
1	C	104	PRO	2.1
1	B	145	ASP	2.1
2	G	195	CYS	2.1
1	C	52	GLU	2.1
1	B	128	LEU	2.1
1	B	167	GLU	2.0
1	B	131	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GOL	A	404	6/6	0.84	0.23	8.82	69,88,91,91	0
3	GOL	C	403	6/6	0.92	0.25	3.89	86,91,91,94	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	B	401[B]	6/6	0.91	0.26	3.69	40,42,46,49	6
3	GOL	B	401[A]	6/6	0.91	0.26	3.23	39,42,44,48	6
3	GOL	A	403	6/6	0.88	0.17	1.91	85,86,86,87	0
4	CL	A	406	1/1	0.98	0.16	1.10	59,59,59,59	0
4	CL	C	404	1/1	0.99	0.15	0.89	58,58,58,58	0
3	GOL	C	401	6/6	0.59	0.28	0.67	92,98,101,106	0
3	GOL	A	401[B]	6/6	0.85	0.20	0.43	23,43,46,47	6
3	GOL	A	401[A]	6/6	0.85	0.20	0.38	28,43,44,47	6
3	GOL	A	402	6/6	0.87	0.19	0.35	61,68,72,72	0
3	GOL	C	402	6/6	0.77	0.14	0.01	59,64,67,71	0
3	GOL	A	405	6/6	0.84	0.21	-0.06	65,73,77,80	0
4	CL	A	407	1/1	0.99	0.07	-	55,55,55,55	0

6.5 Other polymers

There are no such residues in this entry.