



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 08:01 PM GMT

PDB ID : 1IE7
Title : PHOSPHATE INHIBITED BACILLUS PASTEURII UREASE CRYSTAL
STRUCTURE
Authors : Benini, S.; Rypniewski, W.R.; Wilson, K.S.; Cieurli, S.; Mangani, S.
Deposited on : 2001-04-09
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	100	 78% 19% ..
2	B	126	 85% 11% ..
3	C	570	 82% 15% .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6964 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UREASE GAMMA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	100	Total	C	N	O	S	5	0	0
			781	493	133	149	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	CXM	MET	MODIFIED RESIDUE	UNP P41022

- Molecule 2 is a protein called UREASE BETA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	122	Total	C	N	O	S	21	0	0
			951	589	171	190	1			

- Molecule 3 is a protein called UREASE ALPHA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	570	Total	C	N	O	S	57	2	0
			4329	2719	743	843	24			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	19	GLU	ARG	CONFLICT	UNP P41020
C	28	TRP	-	INSERTION	UNP P41020
C	29	ILE	GLY	CONFLICT	UNP P41020
C	36	THR	TYR	CONFLICT	UNP P41020
C	37	THR	TYR	CONFLICT	UNP P41020
C	38	TYR	LEU	CONFLICT	UNP P41020
C	220	KCX	LYS	MODIFIED RESIDUE	UNP P41020
C	263	LEU	VAL	CONFLICT	UNP P41020
C	327	ASN	GLN	CONFLICT	UNP P41020

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	420	ILE	MET	CONFLICT	UNP P41020

- Molecule 4 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	2	Total Ni 2 2	0	0

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	1	Total O P 5 4 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	130	Total O 130 130	0	0
6	B	186	Total O 186 186	0	0
6	C	580	Total O 580 580	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: UREASE GAMMA SUBUNIT

Chain A:  78% 19% ..





• Molecule 2: UREASE BETA SUBUNIT

Chain B:  85% 11% ..



• Molecule 3: UREASE ALPHA SUBUNIT

Chain C:  82% 15% .



4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	131.49 Å 131.49 Å 189.49 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	11.99 – 1.85 11.99 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.3 (11.99-1.85) 97.6 (11.99-1.85)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 1.85 Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.170 , 0.210 0.172 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	23.6	Xtriage
Anisotropy	0.350	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 78.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 80351 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6964	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, PO4, CXM, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.50	2/781 (0.3%)	2.40	17/1053 (1.6%)
2	B	2.39	4/963 (0.4%)	1.51	13/1296 (1.0%)
3	C	0.73	7/4407 (0.2%)	1.45	55/5975 (0.9%)
All	All	1.44	13/6151 (0.2%)	1.61	85/8324 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
3	C	1	6
All	All	1	9

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	22	ARG	CZ-NH2	57.60	2.08	1.33
2	B	126	GLU	CA-C	53.72	2.92	1.52
2	B	126	GLU	CA-CB	45.71	2.54	1.53
1	A	22	ARG	CZ-NH1	-35.59	0.86	1.33
3	C	403	LEU	CB-CG	17.39	2.02	1.52
2	B	5	ASN	CB-CG	12.92	1.80	1.51
3	C	326	LYS	C-N	-6.52	1.19	1.34
3	C	329	ILE	CA-CB	-6.21	1.40	1.54
2	B	115	GLN	CB-CG	-6.02	1.36	1.52
3	C	386	LYS	CE-NZ	-5.52	1.35	1.49
3	C	324	HIS	CB-CG	-5.14	1.40	1.50
3	C	102	ASN	C-N	5.03	1.43	1.34
3	C	337	ASP	CG-OD1	-5.01	1.13	1.25

All (85) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	22	ARG	NE-CZ-NH2	-53.34	93.63	120.30
2	B	126	GLU	CB-CA-C	-28.13	54.13	110.40
1	A	22	ARG	NH1-CZ-NH2	-26.11	90.68	119.40
3	C	5	ARG	NE-CZ-NH2	-21.61	109.50	120.30
1	A	22	ARG	NE-CZ-NH1	16.33	128.47	120.30
3	C	64	GLU	OE1-CD-OE2	13.51	139.51	123.30
3	C	64	GLU	CG-CD-OE2	-13.21	91.89	118.30
3	C	513	ARG	NE-CZ-NH2	-12.64	113.98	120.30
3	C	388	ARG	NE-CZ-NH1	12.53	126.57	120.30
3	C	403	LEU	CA-CB-CG	-12.48	86.60	115.30
3	C	337	ASP	CB-CG-OD2	-11.18	108.24	118.30
1	A	37	ALA	O-C-N	10.76	139.92	122.70
3	C	5	ARG	NE-CZ-NH1	10.39	125.50	120.30
2	B	5	ASN	CA-CB-CG	-10.09	91.20	113.40
3	C	388	ARG	NE-CZ-NH2	-9.85	115.37	120.30
3	C	42	VAL	CA-CB-CG1	9.65	125.37	110.90
3	C	326	LYS	C-N-CA	9.12	144.51	121.70
3	C	566	ARG	NE-CZ-NH2	-9.06	115.77	120.30
3	C	403	LEU	CB-CG-CD2	8.88	126.09	111.00
3	C	327	ASN	N-CA-CB	8.84	126.52	110.60
1	A	37	ALA	CA-C-O	-8.72	101.78	120.10
2	B	54	GLU	OE1-CD-OE2	8.27	133.22	123.30
3	C	337	ASP	CB-CG-OD1	7.97	125.47	118.30
3	C	326	LYS	O-C-N	-7.68	110.42	122.70
1	A	14	PHE	CA-C-N	7.62	133.97	117.20
3	C	326	LYS	CA-C-N	7.48	133.65	117.20
1	A	79	ASP	CB-CG-OD2	-7.39	111.65	118.30
1	A	26	ARG	NE-CZ-NH2	-7.28	116.66	120.30
3	C	264	ARG	NE-CZ-NH2	-7.25	116.68	120.30
3	C	234	ARG	NE-CZ-NH2	-7.06	116.77	120.30
3	C	35	TYR	CB-CG-CD1	-7.03	116.78	121.00
2	B	115	GLN	CA-CB-CG	7.00	128.80	113.40
3	C	35	TYR	CB-CG-CD2	6.90	125.14	121.00
3	C	324	HIS	CA-CB-CG	6.82	125.20	113.60
3	C	386	LYS	CD-CE-NZ	6.73	127.19	111.70
2	B	5	ASN	CB-CG-OD1	6.65	134.91	121.60
3	C	107	ASP	CB-CG-OD1	6.46	124.12	118.30
3	C	141	ILE	CB-CG1-CD1	-6.46	95.81	113.90
1	A	44	MET	CA-CB-CG	-6.43	102.36	113.30
3	C	512	ARG	NE-CZ-NH1	6.43	123.51	120.30
1	A	48	ARG	NE-CZ-NH2	-6.28	117.16	120.30
2	B	66	ARG	NE-CZ-NH1	6.16	123.38	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	14	PHE	O-C-N	-6.07	112.99	122.70
3	C	402	ARG	NE-CZ-NH1	5.92	123.26	120.30
3	C	269	ARG	NE-CZ-NH2	-5.89	117.36	120.30
3	C	520	CYS	CA-CB-SG	5.88	124.59	114.00
3	C	102	ASN	CA-C-O	5.86	132.41	120.10
3	C	513	ARG	CG-CD-NE	-5.80	99.62	111.80
3	C	141	ILE	CA-CB-CG2	5.74	122.38	110.90
1	A	15	LEU	CB-CA-C	5.74	121.11	110.20
3	C	62	ARG	NE-CZ-NH2	-5.72	117.44	120.30
3	C	5	ARG	CD-NE-CZ	5.70	131.58	123.60
1	A	66	ARG	NE-CZ-NH2	-5.70	117.45	120.30
2	B	126	GLU	N-CA-C	-5.69	95.65	111.00
1	A	66	ARG	CA-CB-CG	5.68	125.89	113.40
3	C	286	ASP	CB-CG-OD1	5.63	123.37	118.30
1	A	59	GLU	OE1-CD-OE2	-5.63	116.55	123.30
3	C	305	ARG	NE-CZ-NH1	5.59	123.10	120.30
2	B	38	ARG	NE-CZ-NH1	-5.56	117.52	120.30
3	C	376	ARG	NE-CZ-NH2	-5.50	117.55	120.30
3	C	52	GLU	OE1-CD-OE2	-5.46	116.75	123.30
3	C	369	ARG	NE-CZ-NH2	-5.46	117.57	120.30
3	C	234	ARG	CB-CG-CD	5.44	125.75	111.60
2	B	126	GLU	CA-C-O	5.43	131.50	120.10
3	C	234	ARG	CG-CD-NE	-5.43	100.41	111.80
1	A	14	PHE	CA-C-O	-5.39	108.77	120.10
3	C	64	GLU	CA-CB-CG	-5.35	101.64	113.40
3	C	102	ASN	C-N-CD	5.33	139.59	128.40
3	C	566	ARG	NE-CZ-NH1	5.33	122.96	120.30
3	C	143	PRO	N-CA-CB	5.32	109.68	103.30
2	B	33	SER	N-CA-CB	-5.32	102.53	110.50
3	C	478	ARG	NE-CZ-NH1	5.32	122.96	120.30
1	A	37	ALA	N-CA-CB	5.25	117.46	110.10
3	C	329	ILE	CA-CB-CG2	5.25	121.40	110.90
3	C	61	THR	CA-CB-CG2	-5.25	105.06	112.40
3	C	388	ARG	CD-NE-CZ	5.23	130.93	123.60
3	C	120	GLU	OE1-CD-OE2	-5.22	117.04	123.30
3	C	243	ASP	CB-CG-OD1	5.17	122.96	118.30
2	B	5	ASN	CB-CG-ND2	-5.15	104.33	116.70
3	C	341	ARG	NE-CZ-NH2	-5.14	117.73	120.30
3	C	64	GLU	CG-CD-OE1	5.14	128.59	118.30
2	B	13	ARG	CA-CB-CG	5.11	124.64	113.40
3	C	327	ASN	C-N-CA	5.09	134.41	121.70
3	C	5	ARG	NH1-CZ-NH2	5.08	124.99	119.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	37	ASP	CB-CG-OD1	5.05	122.84	118.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	C	327	ASN	CA

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1	CXM	Mainchain
1	A	14	PHE	Mainchain
1	A	22	ARG	Sidechain
3	C	329	ILE	Mainchain
3	C	340	ILE	Mainchain
3	C	366	ALA	Mainchain
3	C	427	ILE	Mainchain
3	C	46	GLY	Mainchain
3	C	522	ASN	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	781	0	801	11	0
2	B	951	0	937	6	0
3	C	4329	0	4297	54	2
4	C	2	0	0	0	0
5	C	5	0	0	0	0
6	A	130	0	0	1	0
6	B	186	0	0	1	0
6	C	580	0	0	10	4
All	All	6964	0	6035	68	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:201:HIS:HE1	3:C:222:HIS:H	1.24	0.85
1:A:14:PHE:O	1:A:15:LEU:C	2.18	0.80
1:A:13:ILE:O	1:A:14:PHE:O	2.05	0.74
3:C:201:HIS:CE1	3:C:222:HIS:H	2.06	0.73
3:C:43:ASN:HD22	3:C:48:LYS:HB3	1.58	0.69
3:C:383:LYS:NZ	3:C:387:GLN:HE22	1.92	0.67
1:A:14:PHE:O	1:A:16:ALA:N	2.31	0.63
3:C:26:ASP:HB3	6:C:1366:HOH:O	1.99	0.60
2:B:25:ARG:NH2	6:B:233:HOH:O	2.37	0.57
3:C:352:HIS:HD2	3:C:409:LYS:NZ	2.03	0.56
2:B:22:ASN:HD21	2:B:66:ARG:HH22	1.52	0.56
3:C:418:GLN:NE2	6:C:1239:HOH:O	2.38	0.56
3:C:321:VAL:HA	6:C:1389:HOH:O	2.06	0.56
3:C:470:THR:N	3:C:471:PRO:CD	2.71	0.54
3:C:383:LYS:HZ3	3:C:387:GLN:NE2	2.05	0.54
3:C:81:GLY:HA2	3:C:404:LYS:HE2	1.89	0.54
1:A:12:GLN:HG3	6:A:198:HOH:O	2.08	0.53
3:C:6:GLN:NE2	6:C:1302:HOH:O	2.41	0.53
3:C:383:LYS:HZ3	3:C:387:GLN:HE22	1.58	0.52
3:C:300:SER:OG	3:C:352:HIS:HE1	1.93	0.51
2:B:95:GLU:O	3:C:104:ASP:HB3	2.11	0.51
3:C:249:HIS:CE1	3:C:281:GLY:HA3	2.45	0.51
3:C:274:PHE:O	3:C:275:HIS:C	2.49	0.51
3:C:383:LYS:HZ1	3:C:387:GLN:HE22	1.59	0.50
1:A:4:ASN:HB2	1:A:5:PRO:HD2	1.92	0.50
3:C:220:KCX:HE2	3:C:274:PHE:CD2	2.47	0.50
3:C:253:LEU:HD23	6:C:1274:HOH:O	2.11	0.50
1:A:13:ILE:C	1:A:14:PHE:O	2.51	0.49
3:C:141:ILE:HD12	3:C:141:ILE:O	2.12	0.49
3:C:352:HIS:HD2	3:C:409:LYS:HZ2	1.61	0.48
3:C:302:ASN:N	3:C:303:PRO:CD	2.76	0.48
3:C:386:LYS:HG3	6:C:1236:HOH:O	2.14	0.48
3:C:26:ASP:HB2	6:C:1236:HOH:O	2.13	0.48
3:C:468:ILE:HB	3:C:469:PRO:HD2	1.95	0.47
3:C:481:GLY:HA3	6:C:1333:HOH:O	2.14	0.47
2:B:10:GLY:HA2	3:C:21:ARG:O	2.16	0.46
3:C:362:THR:O	3:C:368:GLY:HA3	2.16	0.45
2:B:22:ASN:ND2	2:B:66:ARG:HH22	2.15	0.45
3:C:220:KCX:HE2	3:C:274:PHE:HD2	1.82	0.45
3:C:383:LYS:NZ	3:C:387:GLN:NE2	2.60	0.45
3:C:201:HIS:CE1	3:C:225:TRP:HB2	2.52	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:61:THR:OG1	3:C:64:GLU:HG3	2.17	0.44
1:A:23:ARG:HB3	1:A:28:LEU:HD12	1.99	0.44
3:C:70:LEU:HD11	3:C:86:ASP:HB3	1.99	0.44
3:C:470:THR:N	3:C:471:PRO:HD3	2.33	0.44
3:C:110:THR:HA	3:C:111:PRO:HD3	1.85	0.43
3:C:177:PRO:O	3:C:181:GLU:HG3	2.19	0.43
3:C:305:ARG:HA	3:C:306:PRO:HA	1.75	0.43
3:C:284:ALA:HA	3:C:285:PRO:HA	1.78	0.43
3:C:303:PRO:HG2	3:C:367:MET:O	2.19	0.43
1:A:73:VAL:N	1:A:74:PRO:CD	2.81	0.43
3:C:452:LYS:HG3	6:C:1045:HOH:O	2.19	0.43
3:C:220:KCX:CX	3:C:222:HIS:HD2	2.32	0.43
3:C:51:ARG:HD2	3:C:51:ARG:HH11	1.63	0.42
3:C:43:ASN:ND2	3:C:48:LYS:HB3	2.30	0.42
3:C:301:THR:CG2	3:C:363:ASP:HB2	2.50	0.42
3:C:386:LYS:HE3	3:C:386:LYS:HB2	1.84	0.42
3:C:526:LYS:HD3	6:C:1293:HOH:O	2.19	0.42
1:A:75:GLU:HG2	2:B:8:VAL:HG22	2.01	0.41
3:C:468:ILE:HB	3:C:469:PRO:CD	2.50	0.41
3:C:472:GLN:HB3	3:C:473:PRO:HA	2.02	0.41
3:C:513:ARG:HB3	3:C:513:ARG:HE	1.71	0.41
1:A:15:LEU:HD12	1:A:15:LEU:O	2.20	0.41
3:C:141:ILE:HG13	3:C:365:LEU:HD13	2.02	0.41
3:C:132:GLY:HA3	3:C:155:THR:OG1	2.21	0.41
3:C:137:HIS:CE1	3:C:274:PHE:CD2	3.09	0.40
1:A:4:ASN:HB2	1:A:5:PRO:CD	2.51	0.40
3:C:282:GLY:O	3:C:283:HIS:C	2.60	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:64:GLU:OE2	6:C:1432:HOH:O[11_555]	1.60	0.60
6:C:1032:HOH:O	6:C:1473:HOH:O[12_565]	1.91	0.29
6:C:1032:HOH:O	6:C:1480:HOH:O[12_565]	2.01	0.19
3:C:63:THR:OG1	6:C:1470:HOH:O[11_555]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	98/100 (98%)	93 (95%)	4 (4%)	1 (1%)	19	6
2	B	120/126 (95%)	116 (97%)	3 (2%)	1 (1%)	24	9
3	C	569/570 (100%)	544 (96%)	22 (4%)	3 (0%)	34	17
All	All	787/796 (99%)	753 (96%)	29 (4%)	5 (1%)	30	13

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	327	ASN
1	A	14	PHE
2	B	99	ILE
3	C	275	HIS
3	C	367	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	84/84 (100%)	83 (99%)	1 (1%)	78	69
2	B	101/105 (96%)	101 (100%)	0	100	100
3	C	462/460 (100%)	446 (96%)	16 (4%)	43	23
All	All	647/649 (100%)	630 (97%)	17 (3%)	54	36

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	67	ASP
3	C	1	MET
3	C	43	ASN
3	C	141	ILE
3	C	158	PHE
3	C	249	HIS
3	C	253	LEU
3	C	285	PRO
3	C	324	HIS
3	C	326	LYS
3	C	327	ASN
3	C	328	ASN
3	C	395	LYS
3	C	396	ASN
3	C	520	CYS
3	C	526	LYS
3	C	536	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	HIS
2	B	22	ASN
3	C	7	GLN
3	C	43	ASN
3	C	201	HIS
3	C	267	ASN
3	C	352	HIS
3	C	387	GLN
3	C	472	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CXM	A	1	1	6,10,11	1.86	2 (33%)	4,11,13	3.32	2 (50%)
3	KCX	C	220	3,4	7,11,12	0.82	0	7,12,14	3.53	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CXM	A	1	1	-	0/6/10/12	0/0/0/0
3	KCX	C	220	3,4	-	0/6/10/12	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	CXM	CB-CA	-3.45	1.47	1.53
1	A	1	CXM	CA-N	-2.01	1.43	1.46

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	220	KCX	CE-NZ-CX	-8.79	113.54	123.49
1	A	1	CXM	O-C-CA	-4.23	114.27	125.44
3	C	220	KCX	CD-CG-CB	-2.08	106.26	113.66
1	A	1	CXM	CB-CG-SD	4.97	134.12	113.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	220	KCX	3	0

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	PO4	C	901	4	4,4,4	0.69	0	6,6,6	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PO4	C	901	4	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	99/100 (99%)	-0.76	0 100 100	21, 26, 36, 43	2 (2%)
2	B	122/126 (96%)	-0.62	1 (0%) 87 87	22, 28, 40, 50	5 (4%)
3	C	567/570 (99%)	-0.63	10 (1%) 71 71	19, 25, 46, 64	12 (2%)
All	All	788/796 (98%)	-0.65	11 (1%) 78 78	19, 26, 44, 64	19 (2%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	126	GLU	4.8
3	C	322	CYS	3.6
3	C	326	LYS	3.1
3	C	550	GLY	2.8
3	C	333	VAL	2.4
3	C	316	LEU	2.3
3	C	321	VAL	2.3
3	C	327	ASN	2.2
3	C	320	MET	2.2
3	C	542	GLU	2.1
3	C	549	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	KCX	C	220	12/13	0.96	0.09	-	19,22,35,40	0
1	CXM	A	1	11/12	0.97	0.07	-	21,27,34,34	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	PO4	C	901	5/5	0.97	0.11	0.12	33,35,39,41	5
4	NI	C	601	1/1	0.99	0.06	-0.95	34,34,34,34	0
4	NI	C	600	1/1	1.00	0.06	-0.95	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.