



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:29 PM GMT

PDB ID : 4III
Title : Lambda-[Ru(TAP)2(11-Cl-dppz)] with a DNA decamer at atomic resolution
Authors : Hall, J.P.; Cardin, C.J.
Deposited on : 2012-12-20
Resolution : 1.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1152 (1.10-0.94)
Clashscore	102246	1237 (1.10-0.94)
RSRZ outliers	91569	1154 (1.10-0.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	10	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 609 atoms, of which 178 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a DNA chain called 5'-D(*TP*CP*GP*GP*CP*GP*CP*CP*GP*A)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	10	Total	C	H	N	O	P	1	2	0
			381	116	136	49	69	11			

- Molecule 2 is BARIUM ION (three-letter code: BA) (formula: Ba).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ba	0	0
			1	1		

- Molecule 3 is RU(TAP)2(CL-DPPZ) COMPLEX (three-letter code: RCZ) (formula: C₃₈H₂₁ClN₁₂Ru).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Cl	H	N	Ru	0	1
			146	76	2	42	24	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	81	Total	O	0	0
			81	81		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(*TP*CP*GP*GP*CP*GP*CP*CP*GP*A)-3'

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	42.34Å 42.34Å 39.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.93 – 1.02 23.89 – 1.02	Depositor EDS
% Data completeness (in resolution range)	97.9 (28.93-1.02) 97.9 (23.89-1.02)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.81 (at 1.02Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.101 , 0.114 0.104 , 0.118	Depositor DCC
R_{free} test set	946 reflections (5.40%)	DCC
Wilson B-factor (Å ²)	8.0	Xtriage
Anisotropy	1.193	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 52.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 18474 reflections	Xtriage
F_o, F_c correlation	0.99	EDS
Total number of atoms	609	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BA, RCZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.18	1/274 (0.4%)	1.50	4/419 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	5	DC	O3'-P	-5.00	1.55	1.61

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	10[A]	DA	O5'-P-OP1	-8.21	98.31	105.70
1	A	10[B]	DA	O5'-P-OP1	-8.21	98.31	105.70
1	A	5	DC	O4'-C4'-C3'	-7.79	101.33	106.00
1	A	7	DC	O4'-C4'-C3'	-5.40	102.34	104.50

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	5	DC	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	245	136	136	3	0
2	A	1	0	0	0	0
3	A	104	42	42	1	0
4	A	81	0	0	2	0
All	All	431	178	178	3	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (3) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10[A]:DA:OP2	4:A:277:HOH:O	1.58	1.19
1:A:10[A]:DA:P	4:A:277:HOH:O	2.40	0.62
1:A:9[B]:DG:H21	3:A:102[B]:RCZ:C20	2.34	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	RCZ	A	102[A]	-	65,65,65	1.63	13 (20%)	76,111,111	1.40	12 (15%)
3	RCZ	A	102[B]	-	65,65,65	2.07	15 (23%)	76,111,111	2.11	25 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RCZ	A	102[A]	-	-	0/0/60/60	0/14/14/14
3	RCZ	A	102[B]	-	-	0/0/60/60	0/14/14/14

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	102[B]	RCZ	RU-N5	-8.41	1.94	2.07
3	A	102[B]	RCZ	RU-N12	-4.88	1.99	2.07
3	A	102[B]	RCZ	RU-N9	-4.27	2.00	2.07
3	A	102[B]	RCZ	RU-N2	-4.14	2.00	2.07
3	A	102[B]	RCZ	C22-N6	-3.58	1.30	1.37
3	A	102[A]	RCZ	C38-N12	-2.77	1.28	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	102[A]	RCZ	C9-C8	-2.46	1.36	1.41
3	A	102[B]	RCZ	RU-N8	-2.39	2.03	2.07
3	A	102[B]	RCZ	C8-C7	-2.32	1.40	1.45
3	A	102[A]	RCZ	C28-N8	-2.10	1.29	1.33
3	A	102[B]	RCZ	RU-N1	-2.08	2.03	2.07
3	A	102[A]	RCZ	C29-C36	-2.05	1.39	1.43
3	A	102[B]	RCZ	C29-N9	2.12	1.40	1.37
3	A	102[B]	RCZ	C29-C36	2.21	1.48	1.43
3	A	102[A]	RCZ	C16-C17	2.31	1.41	1.36
3	A	102[B]	RCZ	C7-N4	2.33	1.37	1.33
3	A	102[A]	RCZ	C15-N3	2.34	1.39	1.35
3	A	102[B]	RCZ	C5-C1	2.48	1.45	1.40
3	A	102[B]	RCZ	C16-C17	2.85	1.42	1.36
3	A	102[A]	RCZ	C36-N12	2.89	1.41	1.37
3	A	102[A]	RCZ	C1-N2	2.89	1.41	1.37
3	A	102[A]	RCZ	C29-C32	3.02	1.47	1.40
3	A	102[A]	RCZ	C20-C21	3.04	1.45	1.38
3	A	102[B]	RCZ	C19-C22	3.40	1.48	1.40
3	A	102[A]	RCZ	C29-N9	3.44	1.42	1.37
3	A	102[A]	RCZ	C7-N4	4.49	1.40	1.33
3	A	102[B]	RCZ	C26-N8	4.99	1.45	1.37
3	A	102[A]	RCZ	C26-N8	5.57	1.45	1.37

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	102[B]	RCZ	C38-N12-C36	-5.74	110.18	117.53
3	A	102[B]	RCZ	C29-C36-C35	-4.31	110.84	119.63
3	A	102[B]	RCZ	C36-C35-N11	-3.92	115.81	120.87
3	A	102[B]	RCZ	C31-N10-C32	-3.79	110.45	116.87
3	A	102[B]	RCZ	C20-N5-C19	-3.70	112.78	117.53
3	A	102[B]	RCZ	C19-C22-N6	-3.67	116.14	120.87
3	A	102[B]	RCZ	C7-C6-N3	-3.10	118.56	121.87
3	A	102[B]	RCZ	C5-C1-C10	-3.09	117.21	120.60
3	A	102[A]	RCZ	C17-C16-C15	-2.71	117.94	120.88
3	A	102[A]	RCZ	C18-C14-C13	-2.68	116.96	119.51
3	A	102[B]	RCZ	C26-C25-N7	-2.66	117.44	120.87
3	A	102[A]	RCZ	C28-N8-C26	-2.65	114.13	117.53
3	A	102[B]	RCZ	C8-C10-N1	-2.60	120.19	122.84
3	A	102[A]	RCZ	C14-C13-N4	-2.51	115.73	118.96
3	A	102[B]	RCZ	C1-C5-C6	-2.11	116.70	119.22
3	A	102[A]	RCZ	C9-C8-C10	-2.09	115.35	117.79

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	102[A]	RCZ	C12-N1-C10	-2.07	114.87	117.53
3	A	102[B]	RCZ	C24-C23-C22	2.01	124.93	120.64
3	A	102[B]	RCZ	C14-C13-C15	2.15	121.73	119.47
3	A	102[B]	RCZ	C26-C19-N5	2.24	119.27	116.58
3	A	102[A]	RCZ	C9-C8-C7	2.29	125.89	122.61
3	A	102[B]	RCZ	C30-C31-N10	2.30	127.05	123.41
3	A	102[B]	RCZ	C4-C5-C6	2.32	125.95	122.61
3	A	102[B]	RCZ	C6-N3-C15	2.46	119.60	116.76
3	A	102[A]	RCZ	C16-C17-C18	2.50	122.39	119.28
3	A	102[A]	RCZ	C23-C22-N6	2.55	122.75	118.52
3	A	102[A]	RCZ	C2-C3-C4	2.65	123.16	119.39
3	A	102[B]	RCZ	C21-N6-C22	2.84	121.68	116.87
3	A	102[B]	RCZ	C9-C8-C10	2.87	121.13	117.79
3	A	102[B]	RCZ	C29-C36-N12	3.13	120.33	116.58
3	A	102[B]	RCZ	C5-C6-C7	3.39	123.31	120.05
3	A	102[A]	RCZ	C14-C13-C15	3.41	123.05	119.47
3	A	102[B]	RCZ	C23-C22-N6	3.49	124.32	118.52
3	A	102[A]	RCZ	C33-C32-N10	3.61	124.53	118.52
3	A	102[B]	RCZ	C8-C10-C1	3.68	124.64	120.60
3	A	102[B]	RCZ	C36-C29-C32	3.90	127.57	119.63
3	A	102[B]	RCZ	C29-C32-N10	4.14	126.20	120.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	102[B]	RCZ	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	10/10 (100%)	-0.37	0 100 100	12, 15, 21, 26	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	RCZ	A	102[A]	52/52	1.00	0.07	0.53	8,11,19,27	73
3	RCZ	A	102[B]	52/52	1.00	0.07	0.53	12,16,20,27	73
2	BA	A	101	1/1	1.00	0.05	-	11,11,11,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.