



## wwPDB EM Map/Model Validation Report ⓘ

Apr 10, 2016 – 01:49 PM BST

PDB ID : 3IKU  
EMDB ID: : EMD-5128  
Title : Structural model of ParM filament in closed state from cryo-EM  
Authors : Galkin, V.E.; Orlova, A.; Rivera, C.; Mullins, R.D.; Egelman, E.H.  
Deposited on : 2009-08-06  
Resolution : 18.00 Å(reported)

This is a wwPDB EM Map/Model Validation Report for a publicly released PDB/EMDB entry.  
For rigid body fitted models, validation errors reported here could stem from errors in the original structure(s) used in the fitting.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/EMValidationReportHelp>

---

MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk27241


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 18.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	114402	924
Ramachandran outliers	111179	726
Sidechain outliers	111093	686

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	320	56% 39% ..
1	B	320	56% 40% ..
1	C	320	53% 43% ..
1	D	320	53% 43% ..
1	E	320	53% 43% ..
1	F	320	53% 43% ..
1	G	320	53% 43% ..
1	H	320	53% 43% ..
1	I	320	52% 44% ..

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	J	320	<div><div></div><div>53%</div><div>43%</div><div></div><div>• •</div></div>
1	K	320	<div><div></div><div>57%</div><div>39%</div><div></div><div>• •</div></div>
1	L	320	<div><div></div><div>58%</div><div>38%</div><div></div><div>• •</div></div>

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 29880 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Plasmid segregation protein parM.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	316	Total	C	N	O	S	0	0
			2490	1569	421	492	8		
1	B	316	Total	C	N	O	S	0	0
			2490	1569	421	492	8		
1	C	316	Total	C	N	O	S	0	0
			2490	1569	421	492	8		
1	D	316	Total	C	N	O	S	0	0
			2490	1569	421	492	8		
1	E	316	Total	C	N	O	S	0	0
			2490	1569	421	492	8		
1	F	316	Total	C	N	O	S	0	0
			2490	1569	421	492	8		
1	G	316	Total	C	N	O	S	0	0
			2490	1569	421	492	8		
1	H	316	Total	C	N	O	S	0	0
			2490	1569	421	492	8		
1	I	316	Total	C	N	O	S	0	0
			2490	1569	421	492	8		
1	J	316	Total	C	N	O	S	0	0
			2490	1569	421	492	8		
1	K	316	Total	C	N	O	S	0	0
			2490	1569	421	492	8		
1	L	316	Total	C	N	O	S	0	0
			2490	1569	421	492	8		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Plasmid segregation protein parM




- Molecule 1: Plasmid segregation protein parM




- Molecule 1: Plasmid segregation protein parM






• Molecule 1: Plasmid segregation protein parM




• Molecule 1: Plasmid segregation protein parM




• Molecule 1: Plasmid segregation protein parM






• Molecule 1: Plasmid segregation protein parM

Chain G: 53% 43% ..




• Molecule 1: Plasmid segregation protein parM

Chain H: 53% 43% ..




• Molecule 1: Plasmid segregation protein parM

Chain I: 52% 44% ..



• Molecule 1: Plasmid segregation protein parM





## 4 Experimental information

Property	Value	Source
Reconstruction method	HELICAL	Depositor
Imposed symmetry	POINT, Not provided	Depositor
Number of images	Not provided	Depositor
Resolution determination method	Not provided	Depositor
CTF correction method	The micrographs were multiplied by the CTF to correct phases	Depositor
Microscope	Not provided	Depositor
Voltage (kV)	Not provided	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	Not provided	Depositor
Minimum defocus (nm)	Not provided	Depositor
Maximum defocus (nm)	Not provided	Depositor
Magnification	Not provided	Depositor
Image detector	Not provided	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >2	RMSZ	# Z  >2
1	A	0.40	0/2527	0.67	0/3413
1	B	0.40	0/2527	0.67	0/3413
1	C	0.40	0/2527	0.67	0/3413
1	D	0.40	0/2527	0.67	0/3413
1	E	0.40	0/2527	0.67	0/3413
1	F	0.40	0/2527	0.67	0/3413
1	G	0.40	0/2527	0.67	0/3413
1	H	0.40	0/2527	0.67	0/3413
1	I	0.40	0/2527	0.67	0/3413
1	J	0.40	0/2527	0.67	0/3413
1	K	0.40	0/2527	0.67	0/3413
1	L	0.40	0/2527	0.67	0/3413
All	All	0.40	0/30324	0.67	0/40956

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2490	0	2467	304	0
1	B	2490	0	2466	315	0
1	C	2490	0	2464	411	0
1	D	2490	0	2464	412	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2490	0	2464	407	0
1	F	2490	0	2464	408	0
1	G	2490	0	2464	412	0
1	H	2490	0	2464	416	0
1	I	2490	0	2464	413	0
1	J	2490	0	2464	412	0
1	K	2490	0	2468	314	0
1	L	2490	0	2468	299	0
All	All	29880	0	29581	3422	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 58.

The worst 5 of 3422 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:112:ASN:HB3	1:I:43:LYS:CE	1.24	1.66
1:A:112:ASN:HB3	1:C:43:LYS:CE	1.24	1.65
1:E:112:ASN:HB3	1:G:43:LYS:CE	1.24	1.65
1:F:112:ASN:HB3	1:H:43:LYS:CE	1.24	1.61
1:H:112:ASN:HB3	1:J:43:LYS:CE	1.24	1.59

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	306/320 (96%)	290 (95%)	14 (5%)	2 (1%)	26	71
1	B	306/320 (96%)	290 (95%)	14 (5%)	2 (1%)	26	71
1	C	306/320 (96%)	290 (95%)	14 (5%)	2 (1%)	26	71

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	306/320 (96%)	290 (95%)	14 (5%)	2 (1%)	26	71
1	E	306/320 (96%)	290 (95%)	14 (5%)	2 (1%)	26	71
1	F	306/320 (96%)	290 (95%)	14 (5%)	2 (1%)	26	71
1	G	306/320 (96%)	290 (95%)	14 (5%)	2 (1%)	26	71
1	H	306/320 (96%)	290 (95%)	14 (5%)	2 (1%)	26	71
1	I	306/320 (96%)	290 (95%)	14 (5%)	2 (1%)	26	71
1	J	306/320 (96%)	290 (95%)	14 (5%)	2 (1%)	26	71
1	K	306/320 (96%)	290 (95%)	14 (5%)	2 (1%)	26	71
1	L	306/320 (96%)	290 (95%)	14 (5%)	2 (1%)	26	71
All	All	3672/3840 (96%)	3480 (95%)	168 (5%)	24 (1%)	31	71

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	211	SER
1	B	211	SER
1	C	211	SER
1	D	211	SER
1	E	211	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	281/284 (99%)	263 (94%)	18 (6%)	22	58
1	B	281/284 (99%)	263 (94%)	18 (6%)	22	58
1	C	281/284 (99%)	263 (94%)	18 (6%)	22	58
1	D	281/284 (99%)	263 (94%)	18 (6%)	22	58
1	E	281/284 (99%)	263 (94%)	18 (6%)	22	58
1	F	281/284 (99%)	263 (94%)	18 (6%)	22	58
1	G	281/284 (99%)	263 (94%)	18 (6%)	22	58

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	281/284 (99%)	263 (94%)	18 (6%)	22	58
1	I	281/284 (99%)	263 (94%)	18 (6%)	22	58
1	J	281/284 (99%)	263 (94%)	18 (6%)	22	58
1	K	281/284 (99%)	263 (94%)	18 (6%)	22	58
1	L	281/284 (99%)	263 (94%)	18 (6%)	22	58
All	All	3372/3408 (99%)	3156 (94%)	216 (6%)	26	58

5 of 216 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	127	ARG
1	G	221	LEU
1	L	13	LYS
1	F	185	LEU
1	G	13	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 176 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	112	ASN
1	G	240	ASN
1	K	308	GLN
1	F	228	HIS
1	G	47	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.