



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 20, 2016 – 05:22 AM EDT

PDB ID : 5IL0
Title : Crystal structural of the METTL3-METTL14 complex for N6-adenosine methylation
Authors : Wang, X.; Guan, Z.; Zou, T.; Yin, P.
Deposited on : 2016-03-04
Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027790
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027790


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6965 (1.90-1.86)
Clashscore	102246	7778 (1.90-1.86)
Ramachandran outliers	100387	7691 (1.90-1.86)
Sidechain outliers	100360	7692 (1.90-1.86)
RSRZ outliers	91569	6979 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	212	<div> <div>9%</div> <div>84%</div> <div>14%</div> <div>.</div> </div>
2	B	300	<div> <div>6%</div> <div>83%</div> <div>9%</div> <div>6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	B	501	-	-	X	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4347 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called METTL3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	211	Total	C	N	O	S	0	0	0
			1706	1090	304	302	10			

- Molecule 2 is a protein called METTL14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	282	Total	C	N	O	P S	0	0	0
			2301	1463	398	427	1 12			

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	Br 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	102	Total 102	O 102	0	0
5	B	229	Total 229	O 229	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: METTL3



• Molecule 2: METTL14



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	101.69Å 101.69Å 116.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.27 – 1.88 36.27 – 1.88	Depositor EDS
% Data completeness (in resolution range)	99.9 (36.27-1.88) 95.0 (36.27-1.88)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 1.88Å)	Xtriage
Refinement program	PHENIX (1.10.1 _2155: ???)	Depositor
R, R_{free}	0.175 , 0.205 0.171 , 0.202	Depositor DCC
R_{free} test set	2393 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	27.8	Xtriage
Anisotropy	0.009	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4347	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, BR, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/1752	0.55	0/2381
2	B	0.39	0/2348	0.58	0/3180
All	All	0.37	0/4100	0.56	0/5561

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1706	0	1685	20	0
2	B	2301	0	2246	22	0
3	A	4	0	6	2	0
3	B	4	0	6	6	0
4	B	1	0	0	0	0
5	A	102	0	0	0	0
5	B	229	0	0	1	0
All	All	4347	0	3943	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:276:ASP:HB2	2:B:277:PRO:HD3	1.57	0.86
2:B:311:ILE:HG22	3:B:501:EDO:H11	1.58	0.84
1:A:537:PRO:HB3	3:A:601:EDO:H22	1.77	0.65
2:B:168:MET:HE3	2:B:369:TYR:HA	1.80	0.64
1:A:373:TRP:HB2	1:A:551:LEU:HD13	1.81	0.62
1:A:383:VAL:HG22	1:A:422:LEU:HD21	1.83	0.60
2:B:200:ARG:NH1	5:B:604:HOH:O	2.34	0.60
1:A:456:ILE:HD11	2:B:262:ILE:HD11	1.84	0.59
2:B:276:ASP:HB2	2:B:277:PRO:CD	2.31	0.59
1:A:411:ASP:CG	1:A:435:ARG:HH22	2.05	0.58
2:B:191:LEU:HD22	2:B:193:PRO:HD3	1.85	0.58
1:A:411:ASP:OD1	1:A:435:ARG:NH2	2.39	0.56
2:B:311:ILE:H	3:B:501:EDO:C1	2.20	0.55
1:A:533:LEU:HD12	1:A:547:LEU:HD12	1.88	0.55
2:B:297:LYS:HB2	2:B:300:THR:HG22	1.90	0.54
2:B:325:GLU:CD	2:B:349:ARG:HH11	2.12	0.53
2:B:311:ILE:H	3:B:501:EDO:H12	1.72	0.53
1:A:377:ASP:OD2	1:A:379:ARG:HD3	2.08	0.53
2:B:200:ARG:NH2	2:B:276:ASP:HB3	2.24	0.52
1:A:480:LYS:HD2	2:B:313:LEU:HD21	1.91	0.52
1:A:434:GLY:O	1:A:437:MET:HG2	2.11	0.50
2:B:310:ASP:HA	3:B:501:EDO:H12	1.93	0.49
1:A:468:ARG:HD3	3:B:501:EDO:O1	2.13	0.48
2:B:310:ASP:OD1	3:B:501:EDO:H21	2.13	0.48
2:B:196:GLU:HG2	2:B:210:CYS:HB3	1.93	0.48
1:A:396:PRO:HG2	1:A:398:TRP:CE2	2.52	0.45
1:A:381:LEU:HD23	1:A:574:ILE:HG21	2.00	0.44
1:A:456:ILE:HD11	2:B:262:ILE:CD1	2.47	0.44
2:B:277:PRO:HB2	2:B:278:LYS:H	1.50	0.44
1:A:459:LYS:HE2	1:A:481:GLU:HG2	1.99	0.44
2:B:142:ARG:NH2	2:B:329:GLU:OE2	2.39	0.42
1:A:436:ALA:HA	1:A:439:LEU:HB3	2.02	0.42
1:A:552:ASP:N	3:A:601:EDO:O2	2.40	0.42
1:A:378:ILE:HG13	1:A:534:PHE:CE1	2.55	0.41
2:B:390:GLU:OE1	2:B:394:ARG:HD3	2.19	0.41
1:A:383:VAL:CG2	1:A:422:LEU:HD21	2.50	0.41
2:B:261:TRP:CE2	2:B:327:PRO:HG3	2.56	0.40
2:B:269:PRO:HA	2:B:270:GLY:HA2	1.67	0.40
1:A:388:LYS:O	1:A:529:ARG:HD3	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/212 (99%)	203 (97%)	5 (2%)	1 (0%)	34	20
2	B	275/300 (92%)	267 (97%)	5 (2%)	3 (1%)	17	6
All	All	484/512 (94%)	470 (97%)	10 (2%)	4 (1%)	24	10

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	276	ASP
2	B	277	PRO
1	A	578	LYS
2	B	270	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/188 (99%)	180 (97%)	6 (3%)	46	33
2	B	250/268 (93%)	242 (97%)	8 (3%)	46	33
All	All	436/456 (96%)	422 (97%)	14 (3%)	46	33

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	369	PHE
1	A	474	HIS
1	A	480	LYS
1	A	529	ARG
1	A	533	LEU
1	A	556	LEU
2	B	191	LEU
2	B	207	ASN
2	B	243	LEU
2	B	271	LYS
2	B	274	THR
2	B	278	LYS
2	B	298	ARG
2	B	320	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SEP	B	399	2	7,9,10	1.75	3 (42%)	8,12,14	1.76	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SEP	B	399	2	-	0/5/8/10	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	399	SEP	P-O2P	2.03	1.61	1.54
2	B	399	SEP	P-O3P	2.10	1.62	1.54
2	B	399	SEP	P-O1P	3.29	1.61	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	399	SEP	OG-CB-CA	4.24	111.95	108.26

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	601	-	3,3,3	0.42	0	2,2,2	0.52	0
3	EDO	B	501	-	3,3,3	0.43	0	2,2,2	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	601	-	-	0/1/1/1	0/0/0/0
3	EDO	B	501	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	601	EDO	2	0
3	B	501	EDO	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	211/212 (99%)	0.34	19 (9%) 12 12	20, 43, 80, 102	0
2	B	281/300 (93%)	0.10	18 (6%) 23 24	19, 31, 70, 109	0
All	All	492/512 (96%)	0.20	37 (7%) 17 18	19, 37, 77, 109	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	272	THR	6.4
2	B	206	ALA	5.6
2	B	321	ILE	5.2
1	A	402	MET	4.6
2	B	270	GLY	4.3
1	A	369	PHE	4.0
2	B	274	THR	3.7
2	B	199	TYR	3.7
1	A	579	ASN	3.5
2	B	207	ASN	3.4
2	B	205	THR	3.4
2	B	401	PRO	3.3
1	A	403	GLU	3.2
1	A	492	GLN	3.1
1	A	573	ILE	3.1
1	A	571	ASP	3.0
2	B	322	GLY	3.0
1	A	578	LYS	2.9
2	B	116	HIS	2.8
1	A	566	LYS	2.8
2	B	208	GLU	2.7
1	A	490	ASN	2.7
1	A	400(A)	HIS	2.7
2	B	269	PRO	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	577	PRO	2.6
2	B	271	LYS	2.6
1	A	572	GLY	2.5
1	A	575	SER	2.5
2	B	268	ASN	2.5
2	B	200	ARG	2.3
1	A	370	PRO	2.3
1	A	565	PHE	2.3
1	A	554	ILE	2.3
2	B	260	CYS	2.2
2	B	277	PRO	2.2
1	A	561	VAL	2.2
1	A	559	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SEP	B	399	10/11	0.81	0.15	-	32,55,86,87	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	EDO	B	501	4/4	0.87	0.28	3.56	41,45,45,52	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	EDO	A	601	4/4	0.92	0.14	0.80	54,63,64,65	0
4	BR	B	502	1/1	1.00	0.04	-3.63	38,38,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.