



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:32 PM GMT

PDB ID : 4IN5
Title : (M)L214G mutant of the Rhodobacter sphaeroides Reaction Center
Authors : Saer, R.G.; Hardjasa, A.; Murphy, M.E.; Beatty, J.T.
Deposited on : 2013-01-04
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	282	<div> <div>%</div> <div> <div></div> <div>93%</div> <div>6%</div> <div>.</div> </div> </div>
2	M	307	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>10%</div> <div>..</div> </div> </div>
3	H	266	<div> <div>%</div> <div> <div></div> <div>80%</div> <div>9%</div> <div>.</div> <div>10%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	GOL	H	301	-	-	X	X
10	GOL	H	305	-	-	-	X
10	GOL	H	307	-	-	-	X
10	GOL	L	309	-	-	-	X
10	GOL	L	311	-	-	-	X
10	GOL	L	312	-	-	-	X
10	GOL	L	314	-	-	X	X
13	PC1	M	408	-	-	-	X
4	LDA	H	309	-	-	-	X
4	LDA	L	301	-	-	-	X
4	LDA	L	302	-	-	-	X
4	LDA	M	404	-	-	-	X
4	LDA	M	412	-	-	-	X
6	U10	L	304[A]	-	-	-	X
6	U10	L	304[B]	-	-	-	X
7	BCL	M	401[A]	-	-	-	X
7	BCL	M	401[B]	-	-	-	X
8	PO4	M	411	-	-	X	-
9	HTO	L	307	-	-	-	X

2 Entry composition

There are 15 unique types of molecules in this entry. The entry contains 7489 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reaction center protein L chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	281	Total	C	N	O	S	0	2	0
			2242	1515	356	363	8			

- Molecule 2 is a protein called Reaction center protein M chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	M	302	Total	C	N	O	S	0	1	0
			2407	1604	395	398	10			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	214	GLY	LEU	ENGINEERED MUTATION	UNP P0C0Y9


- Molecule 3 is a protein called Reaction center protein H chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	240	Total	C	N	O	S	0	5	0
			1850	1184	320	337	9			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
H	-5	HIS	-	EXPRESSION TAG	UNP P0C0Y7
H	-4	HIS	-	EXPRESSION TAG	UNP P0C0Y7
H	-3	HIS	-	EXPRESSION TAG	UNP P0C0Y7
H	-2	HIS	-	EXPRESSION TAG	UNP P0C0Y7
H	-1	HIS	-	EXPRESSION TAG	UNP P0C0Y7
H	0	HIS	-	EXPRESSION TAG	UNP P0C0Y7

- Molecule 4 is LAURYL DIMETHYLAMINE-N-OXIDE (three-letter code: LDA) (formula: C₁₄H₃₁NO).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	L	1	Total	C	N	O	0	0
			16	14	1	1		
4	L	1	Total	C	N	O	0	0
			16	14	1	1		
4	M	1	Total	C	N	O	0	0
			16	14	1	1		
4	M	1	Total	C	N	O	0	0
			16	14	1	1		
4	H	1	Total	C	N	O	0	0
			16	14	1	1		
4	H	1	Total	C	N	O	0	0
			16	14	1	1		

- Molecule 5 is BACTERIOPHEOPHYTIN A (three-letter code: BPH) (formula: $C_{55}H_{76}N_4O_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	L	1	Total	C	N	O	0	0
			65	55	4	6		
5	L	1	Total	C	N	O	0	0
			65	55	4	6		

- Molecule 6 is UBIQUINONE-10 (three-letter code: U10) (formula: $C_{59}H_{90}O_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	L	1	Total	C	O	0	1
			46	38	8		
6	M	1	Total	C	O	0	0
			48	44	4		

- Molecule 7 is BACTERIOCHLOROPHYLL A (three-letter code: BCL) (formula: C₅₅H₇₄MgN₄O₆).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	L	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
7	M	1	Total	C	Mg	N	O	0	1
			86	75	1	4	6		
7	M	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
7	M	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		

- Molecule 8 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	L	1	Total	O	P	0	0
			5	4	1		
8	M	1	Total	O	P	0	0
			5	4	1		
8	M	1	Total	O	P	0	0
			5	4	1		

- Molecule 9 is HEPTANE-1,2,3-TRIOL (three-letter code: HTO) (formula: $C_7H_{16}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	L	1	Total	C	O	0	0
			10	7	3		
9	H	1	Total	C	O	0	0
			10	7	3		

- Molecule 10 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	L	1	Total	C	O	0	0
			6	3	3		
10	L	1	Total	C	O	0	0
			6	3	3		
10	L	1	Total	C	O	0	0
			6	3	3		
10	L	1	Total	C	O	0	0
			6	3	3		
10	L	1	Total	C	O	0	0
			6	3	3		
10	L	1	Total	C	O	0	0
			6	3	3		
10	M	1	Total	C	O	0	0
			6	3	3		
10	H	1	Total	C	O	0	0
			6	3	3		
10	H	1	Total	C	O	0	0
			6	3	3		
10	H	1	Total	C	O	0	0
			6	3	3		
10	H	1	Total	C	O	0	0
			6	3	3		
10	H	1	Total	C	O	0	0
			6	3	3		

- Molecule 11 is FE (III) ION (three-letter code: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	M	1	Total	Fe	0	0
			1	1		

- Molecule 12 is SPHEROIDENE (three-letter code: SPO) (formula: $C_{41}H_{60}O$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	M	1	Total	C	O	0	0
			42	41	1		

- Molecule 13 is 1,2-DIACYL-SN-GLYCERO-3-PHOSPHOCHOLINE (three-letter code: PC1) (formula: $C_{44}H_{88}NO_8P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
13	M	1	Total	C	N	O	P	0	0
			43	33	1	8	1		

- Molecule 14 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	H	1	Total	K	0	0
			1	1		


- Molecule 15 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	L	56	Total	O	0	0
			56	56		
15	M	53	Total	O	0	0
			53	53		
15	H	83	Total	O	0	0
			83	83		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Reaction center protein L chain



- Molecule 2: Reaction center protein M chain



- Molecule 3: Reaction center protein H chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	139.11Å 139.11Å 184.69Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	55.62 – 2.20 55.56 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (55.62-2.20) 99.9 (55.56-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.181 , 0.206 0.188 , 0.211	Depositor DCC
R_{free} test set	5239 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	39.1	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.0	EDS
Estimated twinning fraction	0.017 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 104887 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7489	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BCL, GOL, LDA, HTO, BPH, K, PC1, FE, SPO, U10, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.89	1/2342 (0.0%)	0.76	1/3206 (0.0%)
2	M	0.82	1/2504 (0.0%)	0.82	8/3418 (0.2%)
3	H	0.85	0/1923	0.87	2/2611 (0.1%)
All	All	0.85	2/6769 (0.0%)	0.81	11/9235 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	H	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	67	TYR	CB-CG	5.65	1.60	1.51
2	M	41	TRP	CB-CG	-5.04	1.41	1.50

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	83	ARG	NE-CZ-NH2	-8.13	116.23	120.30
3	H	83	ARG	NE-CZ-NH1	7.88	124.24	120.30
2	M	228	ARG	NE-CZ-NH2	-7.79	116.41	120.30
2	M	29	ARG	NE-CZ-NH1	-7.01	116.79	120.30
2	M	253	ARG	NE-CZ-NH2	6.99	123.80	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	H	248	ARG	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	2242	0	2194	15	0
2	M	2407	0	2315	19	0
3	H	1850	0	1875	21	0
4	H	32	0	62	3	0
4	L	32	0	62	1	0
4	M	32	0	62	3	0
5	L	130	0	152	9	0
6	L	46	0	46	11	0
6	M	48	0	63	2	0
7	L	66	0	74	2	0
7	M	218	0	226	19	0
8	L	5	0	0	1	0
8	M	10	0	0	3	0
9	H	10	0	16	1	0
9	L	10	0	16	0	0
10	H	30	0	40	5	0
10	L	36	0	48	7	0
10	M	6	0	8	0	0
11	M	1	0	0	0	0
12	M	42	0	60	1	0
13	M	43	0	60	0	0
14	H	1	0	0	0	0
15	H	83	0	0	3	0
15	L	56	0	0	3	0
15	M	53	0	0	2	0
All	All	7489	0	7379	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 94 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:L:304[B]:U10:C8	6:L:304[B]:U10:H1M1	1.67	1.23
6:L:304[B]:U10:H8	6:L:304[B]:U10:H1M1	1.54	0.88
5:L:313:BPH:HHC	5:L:313:BPH:HBB3	1.60	0.84
5:L:303:BPH:HHC	5:L:303:BPH:HBB3	1.60	0.83
10:L:314:GOL:O1	15:L:454:HOH:O	2.00	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	281/282 (100%)	276 (98%)	5 (2%)	0	100	100
2	M	301/307 (98%)	290 (96%)	10 (3%)	1 (0%)	46	50
3	H	243/266 (91%)	240 (99%)	1 (0%)	2 (1%)	24	22
All	All	825/855 (96%)	806 (98%)	16 (2%)	3 (0%)	52	42

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	H	249[A]	LYS
3	H	249[B]	LYS
2	M	301	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	222/221 (100%)	215 (97%)	7 (3%)	46	57
2	M	236/239 (99%)	232 (98%)	4 (2%)	68	81
3	H	200/214 (94%)	191 (96%)	9 (4%)	34	41
All	All	658/674 (98%)	638 (97%)	20 (3%)	54	60

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	M	72	ILE
2	M	216	PHE
3	H	220[B]	LYS
2	M	52	LEU
2	M	55	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
2	M	77	GLN
3	H	201	ASN
2	M	193	HIS
1	L	183	ASN
2	M	187	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 37 ligands modelled in this entry, 2 are monoatomic - leaving 35 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
10	GOL	H	301	-	5,5,5	1.01	0	5,5,5	1.76	2 (40%)
10	GOL	H	302	-	5,5,5	0.53	0	5,5,5	0.74	0
10	GOL	H	303	-	5,5,5	0.43	0	5,5,5	0.49	0
4	LDA	H	304	-	15,15,15	3.91	3 (20%)	16,17,17	1.53	2 (12%)
10	GOL	H	305	-	5,5,5	0.52	0	5,5,5	0.77	0
10	GOL	H	307	-	5,5,5	0.26	0	5,5,5	0.56	0
9	HTO	H	308	-	9,9,9	0.59	0	8,10,10	0.65	0
4	LDA	H	309	-	15,15,15	3.82	1 (6%)	16,17,17	0.42	0
4	LDA	L	301	-	15,15,15	3.68	1 (6%)	16,17,17	0.36	0
4	LDA	L	302	-	15,15,15	3.98	1 (6%)	16,17,17	1.12	2 (12%)
5	BPH	L	303	-	64,70,70	1.88	16 (25%)	73,101,101	1.94	15 (20%)
6	U10	L	304[A]	-	23,23,63	1.70	2 (8%)	28,31,79	1.88	9 (32%)
6	U10	L	304[B]	-	23,23,63	1.64	2 (8%)	28,31,79	1.10	1 (3%)
7	BCL	L	305	-	53,74,74	1.27	7 (13%)	57,115,115	1.61	9 (15%)
8	PO4	L	306	-	4,4,4	0.37	0	6,6,6	0.31	0
9	HTO	L	307	-	9,9,9	0.88	0	8,10,10	0.66	0
10	GOL	L	308	-	5,5,5	1.16	0	5,5,5	1.26	0
10	GOL	L	309	-	5,5,5	0.36	0	5,5,5	0.32	0
10	GOL	L	310	-	5,5,5	1.21	0	5,5,5	1.43	1 (20%)
10	GOL	L	311	-	5,5,5	0.48	0	5,5,5	0.84	0
10	GOL	L	312	-	5,5,5	0.40	0	5,5,5	0.64	0
5	BPH	L	313	-	64,70,70	1.80	15 (23%)	73,101,101	1.80	16 (21%)
10	GOL	L	314	-	5,5,5	0.72	0	5,5,5	0.61	0
7	BCL	M	401[A]	-	53,74,74	1.43	7 (13%)	57,115,115	2.11	15 (26%)
7	BCL	M	401[B]	-	53,74,74	1.42	7 (13%)	57,115,115	1.92	13 (22%)
7	BCL	M	402	-	53,74,74	1.35	5 (9%)	57,115,115	1.61	13 (22%)
7	BCL	M	403	-	53,74,74	1.61	8 (15%)	57,115,115	1.61	12 (21%)
4	LDA	M	404	-	15,15,15	3.68	1 (6%)	16,17,17	1.01	1 (6%)
6	U10	M	406	-	48,48,63	1.20	4 (8%)	58,61,79	1.61	15 (25%)
12	SPO	M	407	-	40,41,41	0.90	1 (2%)	45,50,50	1.72	7 (15%)
13	PC1	M	408	-	42,42,53	1.13	2 (4%)	46,50,61	1.50	7 (15%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	PO4	M	409	-	4,4,4	0.44	0	6,6,6	0.30	0
10	GOL	M	410	-	5,5,5	0.75	0	5,5,5	0.79	0
8	PO4	M	411	-	4,4,4	1.33	0	6,6,6	0.32	0
4	LDA	M	412	-	15,15,15	3.63	1 (6%)	16,17,17	1.80	3 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	GOL	H	301	-	-	0/4/4/4	0/0/0/0
10	GOL	H	302	-	-	0/4/4/4	0/0/0/0
10	GOL	H	303	-	-	0/4/4/4	0/0/0/0
4	LDA	H	304	-	-	0/13/13/13	0/0/0/0
10	GOL	H	305	-	-	0/4/4/4	0/0/0/0
10	GOL	H	307	-	-	0/4/4/4	0/0/0/0
9	HTO	H	308	-	-	0/10/10/10	0/0/0/0
4	LDA	H	309	-	-	0/13/13/13	0/0/0/0
4	LDA	L	301	-	-	0/13/13/13	0/0/0/0
4	LDA	L	302	-	-	0/13/13/13	0/0/0/0
5	BPH	L	303	-	-	0/54/105/105	0/1/6/6
6	U10	L	304[A]	-	-	0/15/39/87	0/1/1/1
6	U10	L	304[B]	-	-	0/15/39/87	0/1/1/1
7	BCL	L	305	-	-	0/37/137/137	0/0/9/9
8	PO4	L	306	-	-	0/0/0/0	0/0/0/0
9	HTO	L	307	-	-	0/10/10/10	0/0/0/0
10	GOL	L	308	-	-	0/4/4/4	0/0/0/0
10	GOL	L	309	-	-	0/4/4/4	0/0/0/0
10	GOL	L	310	-	-	0/4/4/4	0/0/0/0
10	GOL	L	311	-	-	0/4/4/4	0/0/0/0
10	GOL	L	312	-	-	0/4/4/4	0/0/0/0
5	BPH	L	313	-	-	0/54/105/105	0/1/6/6
10	GOL	L	314	-	-	0/4/4/4	0/0/0/0
7	BCL	M	401[A]	-	-	0/37/137/137	0/0/9/9
7	BCL	M	401[B]	-	-	0/37/137/137	0/0/9/9
7	BCL	M	402	-	-	1/37/137/137	0/0/9/9
7	BCL	M	403	-	-	0/37/137/137	0/0/9/9
4	LDA	M	404	-	-	0/13/13/13	0/0/0/0
6	U10	M	406	-	-	0/45/69/87	0/1/1/1
12	SPO	M	407	-	-	0/47/47/47	0/0/0/0
13	PC1	M	408	-	-	1/46/46/57	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	PO4	M	409	-	-	0/0/0/0	0/0/0/0
10	GOL	M	410	-	-	0/4/4/4	0/0/0/0
8	PO4	M	411	-	-	0/0/0/0	0/0/0/0
4	LDA	M	412	-	-	0/13/13/13	0/0/0/0

The worst 5 of 84 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	L	302	LDA	O1-N1	-15.25	1.25	1.39
4	H	309	LDA	O1-N1	-14.68	1.25	1.39
4	H	304	LDA	O1-N1	-14.53	1.25	1.39
4	M	404	LDA	O1-N1	-14.12	1.26	1.39
4	L	301	LDA	O1-N1	-14.05	1.26	1.39

The worst 5 of 143 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	M	407	SPO	C4-C5-C6	-6.74	115.08	124.67
5	L	303	BPH	CBB-CAB-C3B	-5.96	107.27	120.52
7	M	401[B]	BCL	C1D-CHD-C4C	-5.56	117.58	126.07
7	M	401[A]	BCL	C1D-CHD-C4C	-5.56	117.58	126.07
7	M	401[B]	BCL	C3C-C4C-CHD	-5.01	112.15	123.33

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
13	M	408	PC1	C2-O21-C21-C22
7	M	402	BCL	CGA-O2A-C1-C2

There are no ring outliers.

22 monomers are involved in 62 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	H	301	GOL	4	0
4	H	304	LDA	3	0
10	H	307	GOL	1	0
9	H	308	HTO	1	0
4	L	302	LDA	1	0
5	L	303	BPH	3	0
6	L	304[A]	U10	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	L	304[B]	U10	8	0
7	L	305	BCL	2	0
8	L	306	PO4	1	0
10	L	308	GOL	1	0
10	L	312	GOL	1	0
5	L	313	BPH	6	0
10	L	314	GOL	5	0
7	M	401[A]	BCL	1	0
7	M	401[B]	BCL	3	0
7	M	402	BCL	6	0
7	M	403	BCL	11	0
6	M	406	U10	2	0
12	M	407	SPO	1	0
8	M	411	PO4	3	0
4	M	412	LDA	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	281/282 (99%)	-0.42	2 (0%) 89 88	29, 38, 63, 83	0
2	M	302/307 (98%)	-0.44	6 (1%) 68 67	29, 42, 64, 104	6 (1%)
3	H	240/266 (90%)	-0.68	2 (0%) 87 87	30, 41, 56, 106	3 (1%)
All	All	823/855 (96%)	-0.50	10 (1%) 81 80	29, 40, 63, 106	9 (1%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	H	250	SER	6.0
1	L	277	GLY	3.2
2	M	105	PHE	3.0
2	M	1	ALA	2.9
2	M	109	LEU	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
9	HTO	L	307	10/10	0.67	0.40	22.07	75,90,100,103	0
4	LDA	L	302	16/16	0.57	0.46	11.36	91,97,115,118	0
13	PC1	M	408	43/54	0.35	0.45	10.03	78,108,166,179	0
10	GOL	L	311	6/6	0.84	0.21	8.35	54,67,77,82	0
10	GOL	H	301	6/6	0.85	0.20	8.35	42,59,64,66	0
10	GOL	H	307	6/6	0.95	0.25	7.75	56,73,81,85	0
4	LDA	H	309	16/16	0.56	0.26	7.42	78,89,134,144	0
4	LDA	M	404	16/16	0.76	0.24	6.82	66,78,98,98	0
10	GOL	L	309	6/6	0.87	0.26	6.41	82,82,88,89	0
4	LDA	L	301	16/16	0.62	0.32	5.12	84,96,122,123	0
6	U10	L	304[B]	23/63	0.83	0.30	4.70	47,64,69,74	23
10	GOL	H	305	6/6	0.94	0.19	4.70	41,49,58,68	0
6	U10	L	304[A]	23/63	0.83	0.30	4.24	32,59,75,78	23
7	BCL	M	401[B]	66/66	0.94	0.21	4.15	24,35,62,70	20
7	BCL	M	401[A]	66/66	0.94	0.21	4.15	24,35,53,65	20
10	GOL	L	312	6/6	0.95	0.20	3.66	53,55,60,61	0
4	LDA	M	412	16/16	0.77	0.22	3.43	66,76,87,101	0
10	GOL	L	314	6/6	0.91	0.14	2.26	58,69,75,77	0
4	LDA	H	304	16/16	0.90	0.14	1.91	49,66,81,82	0
10	GOL	L	310	6/6	0.77	0.15	1.71	49,64,77,81	0
8	PO4	M	409	5/5	0.93	0.13	1.57	63,69,76,81	0
6	U10	M	406	48/63	0.93	0.15	1.48	30,53,78,90	0
10	GOL	L	308	6/6	0.91	0.14	1.13	42,58,66,72	0
12	SPO	M	407	42/42	0.96	0.16	1.02	34,43,80,96	0
5	BPH	L	313	65/65	0.96	0.14	0.65	31,41,118,122	0
7	BCL	M	403	66/66	0.98	0.12	0.62	28,33,55,81	0
7	BCL	M	402	66/66	0.97	0.14	0.46	29,36,81,84	0
7	BCL	L	305	66/66	0.97	0.14	0.33	28,35,50,71	0
5	BPH	L	303	65/65	0.96	0.11	0.05	27,34,44,49	0
8	PO4	L	306	5/5	0.97	0.10	-0.13	56,56,61,64	0
11	FE	M	405	1/1	1.00	0.12	-0.93	30,30,30,30	0
14	K	H	306	1/1	0.97	0.05	-2.85	46,46,46,46	0
9	HTO	H	308	10/10	0.58	0.31	-	71,98,111,112	0
10	GOL	M	410	6/6	0.80	0.16	-	61,61,69,69	0
10	GOL	H	303	6/6	0.80	0.19	-	73,80,83,83	0
10	GOL	H	302	6/6	0.80	0.35	-	68,76,78,88	0
8	PO4	M	411	5/5	0.99	0.09	-	47,48,60,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.