



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:31 PM GMT

PDB ID : 4INQ
Title : Crystal structure of Osh3 ORD in complex with PI(4)P from *Saccharomyces cerevisiae*
Authors : Tong, J.; Im, Y.J.
Deposited on : 2013-01-05
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	397	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3199 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Oxysterol-binding protein homolog 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	372	3041	1947	521	568	5	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	600	GLY	-	EXPRESSION TAG	UNP P38713
A	601	ALA	-	EXPRESSION TAG	UNP P38713
A	602	MET	-	EXPRESSION TAG	UNP P38713
A	603	ASP	-	EXPRESSION TAG	UNP P38713
A	604	PRO	-	EXPRESSION TAG	UNP P38713

- Molecule 2 is (2R)-3-[[[(S)-HYDROXY{[(1R,2R,3R,4R,5S,6R)-2,3,5,6-TETRAHYDROXY-4-(PHOSPHONOOXY)CYCLOHEXYL]OXY}PHOSPHORYL]OXY}PROPANE-1,2-DIYL DIOCTANOATE (three-letter code: PIF) (formula: C₂₅H₄₈O₁₆P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			43	25	16	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	115	Total	O	0	0
			115	115		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Oxysterol-binding protein homolog 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	39.89Å 88.99Å 95.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.35 – 2.20 40.35 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (40.35-2.20) 100.0 (40.35-2.20)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.44 (at 2.20Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.232 , 0.280 0.234 , 0.226	Depositor DCC
R_{free} test set	915 reflections (5.09%)	DCC
Wilson B-factor (Å ²)	24.5	Xtriage
Anisotropy	0.497	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 17960 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3199	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PIF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/3120	0.63	0/4218

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3041	0	3034	83	0
2	A	43	0	45	1	0
3	A	115	0	0	3	0
All	All	3199	0	3079	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:860:LYS:HD3	1:A:861:LYS:H	1.17	1.09

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:627:VAL:HG11	1:A:910:ILE:HG12	1.55	0.88
1:A:860:LYS:HD3	1:A:861:LYS:N	1.88	0.87
1:A:702:ILE:HD11	1:A:832:ILE:HG12	1.65	0.77
1:A:739:ILE:HD13	1:A:740:SER:N	1.99	0.77
1:A:631:ASN:ND2	1:A:903:ASN:HD21	1.83	0.76
1:A:776:ILE:HG13	1:A:807:LEU:HD11	1.78	0.66
1:A:914:ASN:ND2	1:A:987:HIS:HE1	1.95	0.65
1:A:643:LEU:HD11	1:A:776:ILE:HD11	1.78	0.64
1:A:655:LEU:HD22	2:A:1001:PIF:H19	1.79	0.64
1:A:650:ASN:ND2	1:A:658:ILE:HG21	2.13	0.63
1:A:847:PHE:CE1	1:A:872:LEU:HD13	2.35	0.61
1:A:738:LEU:C	1:A:738:LEU:HD23	2.21	0.61
1:A:716:ARG:HD2	1:A:895:TYR:CD1	2.36	0.60
1:A:847:PHE:HE1	1:A:872:LEU:HD13	1.66	0.60
1:A:631:ASN:HD22	1:A:903:ASN:HD21	1.51	0.57
1:A:648:ARG:HA	1:A:651:VAL:HG23	1.85	0.57
1:A:858:ASN:HD22	1:A:859:ARG:N	2.03	0.56
1:A:738:LEU:HD23	1:A:739:ILE:N	2.20	0.56
1:A:742:LYS:NZ	1:A:745:HIS:HD2	2.04	0.56
1:A:653:LYS:O	1:A:654:ASP:HB2	2.05	0.55
1:A:887:LEU:HA	1:A:898:THR:HA	1.87	0.55
1:A:674:ILE:HB	1:A:752:PHE:CZ	2.42	0.54
1:A:716:ARG:HD2	1:A:895:TYR:HD1	1.72	0.54
1:A:742:LYS:HE2	1:A:745:HIS:HA	1.90	0.54
1:A:655:LEU:O	1:A:656:SER:HB3	2.08	0.54
1:A:723:LEU:O	1:A:724:ALA:HB3	2.08	0.54
1:A:777:GLU:HG2	1:A:779:ASN:OD1	2.07	0.53
1:A:828:ASP:OD1	1:A:854:PRO:HD3	2.08	0.53
1:A:742:LYS:HZ3	1:A:745:HIS:HD2	1.58	0.52
1:A:690:ARG:O	1:A:693:PRO:HD3	2.10	0.52
1:A:692:ASP:O	1:A:695:THR:HG23	2.10	0.51
1:A:855:PRO:O	1:A:856:SER:HB2	2.10	0.51
1:A:686:LYS:HB2	1:A:686:LYS:NZ	2.26	0.51
1:A:845:GLU:HA	1:A:868:TRP:NE1	2.26	0.51
1:A:946:GLN:NE2	1:A:950:GLU:OE2	2.44	0.51
1:A:858:ASN:N	1:A:858:ASN:HD22	2.09	0.50
1:A:858:ASN:ND2	1:A:858:ASN:N	2.57	0.50
1:A:821:GLU:OE2	1:A:831:HIS:HD2	1.94	0.49
1:A:779:ASN:ND2	1:A:801:THR:OG1	2.45	0.49
1:A:853:PRO:HG3	1:A:860:LYS:CA	2.43	0.49
1:A:665:ASN:HA	1:A:717:LYS:O	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:744:SER:HB3	1:A:749:VAL:HB	1.93	0.49
1:A:837:ALA:HB2	3:A:1179:HOH:O	2.13	0.49
1:A:853:PRO:HG3	1:A:860:LYS:C	2.33	0.48
1:A:648:ARG:HA	1:A:651:VAL:CG2	2.44	0.48
1:A:654:ASP:O	1:A:655:LEU:O	2.32	0.48
1:A:845:GLU:O	1:A:867:LYS:HA	2.15	0.47
1:A:905:ASN:HD21	1:A:921:ARG:HA	1.80	0.47
1:A:627:VAL:HG11	1:A:910:ILE:CG1	2.34	0.47
1:A:757:LEU:HB2	3:A:1120:HOH:O	2.13	0.47
1:A:905:ASN:ND2	1:A:921:ARG:HA	2.30	0.47
1:A:827:GLY:C	1:A:854:PRO:HG3	2.35	0.47
1:A:994:GLN:HG2	1:A:996:TRP:O	2.16	0.46
1:A:914:ASN:HD22	1:A:987:HIS:HE1	1.61	0.46
1:A:832:ILE:HD12	1:A:832:ILE:N	2.31	0.45
1:A:643:LEU:HD13	1:A:647:LEU:HD13	1.97	0.45
1:A:783:ILE:HG21	1:A:785:ARG:NH1	2.31	0.45
1:A:655:LEU:HB2	1:A:656:SER:H	1.49	0.45
1:A:647:LEU:HD23	1:A:771:PHE:CZ	2.52	0.45
1:A:627:VAL:HG12	1:A:909:GLU:CD	2.37	0.45
1:A:855:PRO:O	1:A:856:SER:CB	2.65	0.45
1:A:858:ASN:HD22	1:A:859:ARG:H	1.63	0.45
1:A:783:ILE:HD12	1:A:783:ILE:N	2.33	0.44
1:A:681:ALA:N	1:A:682:PRO:CD	2.81	0.43
1:A:650:ASN:HD21	1:A:658:ILE:HG21	1.82	0.43
1:A:994:GLN:CG	1:A:996:TRP:O	2.67	0.43
1:A:821:GLU:OE2	1:A:831:HIS:CD2	2.72	0.43
1:A:838:GLY:O	1:A:839:MET:HB2	2.18	0.43
1:A:865:ALA:O	1:A:872:LEU:HD12	2.19	0.42
1:A:625:LYS:NZ	3:A:1177:HOH:O	2.52	0.42
1:A:858:ASN:ND2	1:A:858:ASN:H	2.16	0.42
1:A:849:VAL:HB	1:A:864:LEU:HB2	2.02	0.41
1:A:640:PRO:HG3	1:A:716:ARG:CZ	2.50	0.41
1:A:651:VAL:O	1:A:652:GLY:C	2.57	0.41
1:A:627:VAL:HG12	1:A:909:GLU:CG	2.50	0.41
1:A:643:LEU:HD12	1:A:807:LEU:HD22	2.02	0.41
1:A:716:ARG:HD2	1:A:895:TYR:CE1	2.56	0.41
1:A:686:LYS:O	1:A:690:ARG:HG2	2.21	0.40
1:A:706:SER:O	1:A:709:ARG:HG2	2.21	0.40
1:A:739:ILE:HG13	1:A:980:TYR:CG	2.57	0.40
1:A:648:ARG:O	1:A:651:VAL:HG23	2.22	0.40
1:A:673:LEU:HD22	1:A:718:PRO:HG2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	370/397 (93%)	344 (93%)	15 (4%)	11 (3%)	5 2

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	655	LEU
1	A	808	ILE
1	A	653	LYS
1	A	656	SER
1	A	839	MET
1	A	840	PHE
1	A	856	SER
1	A	651	VAL
1	A	654	ASP
1	A	841	SER
1	A	855	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	337/358 (94%)	324 (96%)	13 (4%)	39 48

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	654	ASP
1	A	695	THR
1	A	720	ASN
1	A	738	LEU
1	A	739	ILE
1	A	757	LEU
1	A	769	GLN
1	A	819	GLU
1	A	833	LEU
1	A	858	ASN
1	A	870	GLN
1	A	875	GLU
1	A	879	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	631	ASN
1	A	650	ASN
1	A	665	ASN
1	A	745	HIS
1	A	769	GLN
1	A	780	ASN
1	A	799	GLN
1	A	806	ASN
1	A	818	ASN
1	A	858	ASN
1	A	905	ASN
1	A	914	ASN
1	A	946	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PIF	A	1001	-	43,43,43	1.22	2 (4%)	52,58,58	1.13	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PIF	A	1001	-	-	0/39/63/63	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	PIF	C7-C8	2.33	1.57	1.50
2	A	1001	PIF	P4-O43	3.75	1.63	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	PIF	C23-C22-C21	2.04	125.06	114.53
2	A	1001	PIF	O15-C10-C11	2.38	119.17	111.90
2	A	1001	PIF	O14-C18-C19	4.19	120.64	111.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	PIF	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	372/397 (93%)	0.22	29 (7%) 16 15	7, 27, 58, 81	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	652	GLY	13.4
1	A	838	GLY	8.1
1	A	837	ALA	7.1
1	A	839	MET	7.0
1	A	655	LEU	5.2
1	A	653	LYS	5.0
1	A	810	GLY	4.7
1	A	654	ASP	4.6
1	A	651	VAL	4.5
1	A	808	ILE	4.4
1	A	840	PHE	4.0
1	A	626	SER	3.8
1	A	858	ASN	3.6
1	A	855	PRO	3.4
1	A	627	VAL	3.3
1	A	650	ASN	3.0
1	A	649	LYS	2.8
1	A	772	TRP	2.8
1	A	656	SER	2.6
1	A	643	LEU	2.5
1	A	841	SER	2.4
1	A	966	VAL	2.3
1	A	860	LYS	2.2
1	A	690	ARG	2.2
1	A	625	LYS	2.1
1	A	977	PRO	2.1
1	A	648	ARG	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	862	GLU	2.0
1	A	640	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	PIF	A	1001	43/43	0.88	0.21	0.77	37,49,57,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.