



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 9, 2016 – 10:37 AM EST

PDB ID : 5IOI
Title : X-RAY STRUCTURE OF THE N-TERMINAL DOMAIN OF HUMAN DOUBLECORTIN
Authors : Ruf, A.; Benz, J.; Burger, D.; D'Arcy, B.; Debulpaep, M.; Di Lello, P.; Fry, D.; Huber, W.; Kremer, T.; Laeremans, T.; Matile, H.; Ross, A.; Rudolph, M.G.; Rufer, A.C.; Sharma, A.; Steinmetz, M.O.; Steyaert, J.; Schoch, G.; Stihle, M.; Thoma, R.
Deposited on : 2016-03-08
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


| | | |
|--------------------------------|---|--|
| MolProbity | : | 4.02b-467 |
| Mogul | : | unknown |
| Xtriage (Phenix) | : | 1.9-1692 |
| EDS | : | rb-20028442 |
| Percentile statistics | : | 20151230.v01 (using entries in the PDB archive December 30th 2015) |
| Refmac | : | 5.8.0135 |
| CCP4 | : | 6.5.0 |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | rb-20028442 |

i

X-RAY DIFFRACTION

A

| Metric | Percentile Rank | Value |
|--------|-----------------|-------|
|--------|-----------------|-------|

Ramachandran outliers

electron density. The numeric value is given above the bar.

| | |
|--|--|
| | |
| | |

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5085 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neuronal migration protein doublecortin.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 1 | A | 107 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 866 | 544 | 151 | 168 | 3 | | | |
| 1 | B | 88 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 716 | 450 | 121 | 143 | 2 | | | |
| 1 | C | 96 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 783 | 492 | 134 | 155 | 2 | | | |
| 1 | D | 89 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 724 | 455 | 122 | 144 | 3 | | | |
| 1 | E | 107 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 866 | 544 | 151 | 168 | 3 | | | |
| 1 | F | 95 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 777 | 489 | 133 | 153 | 2 | | | |

There are 60 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|---------------------|------------|
| A | 125 | LEU | - | expression tag | UNP O43602 |
| A | 126 | VAL | - | expression tag | UNP O43602 |
| A | 127 | PRO | - | expression tag | UNP O43602 |
| A | 128 | ARG | - | expression tag | UNP O43602 |
| A | 129 | GLY | - | expression tag | UNP O43602 |
| A | 130 | SER | - | expression tag | UNP O43602 |
| A | 131 | HIS | - | expression tag | UNP O43602 |
| A | 132 | MET | - | expression tag | UNP O43602 |
| A | 215 | ASP | LYS | engineered mutation | UNP O43602 |
| A | 216 | ASP | LYS | engineered mutation | UNP O43602 |
| B | 125 | LEU | - | expression tag | UNP O43602 |
| B | 126 | VAL | - | expression tag | UNP O43602 |
| B | 127 | PRO | - | expression tag | UNP O43602 |
| B | 128 | ARG | - | expression tag | UNP O43602 |
| B | 129 | GLY | - | expression tag | UNP O43602 |
| B | 130 | SER | - | expression tag | UNP O43602 |
| B | 131 | HIS | - | expression tag | UNP O43602 |

Continued on next page...

Continued from previous page...

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|---------------------|------------|
| B | 132 | MET | - | expression tag | UNP O43602 |
| B | 215 | ASP | LYS | engineered mutation | UNP O43602 |
| B | 216 | ASP | LYS | engineered mutation | UNP O43602 |
| C | 125 | LEU | - | expression tag | UNP O43602 |
| C | 126 | VAL | - | expression tag | UNP O43602 |
| C | 127 | PRO | - | expression tag | UNP O43602 |
| C | 128 | ARG | - | expression tag | UNP O43602 |
| C | 129 | GLY | - | expression tag | UNP O43602 |
| C | 130 | SER | - | expression tag | UNP O43602 |
| C | 131 | HIS | - | expression tag | UNP O43602 |
| C | 132 | MET | - | expression tag | UNP O43602 |
| C | 215 | ASP | LYS | engineered mutation | UNP O43602 |
| C | 216 | ASP | LYS | engineered mutation | UNP O43602 |
| D | 125 | LEU | - | expression tag | UNP O43602 |
| D | 126 | VAL | - | expression tag | UNP O43602 |
| D | 127 | PRO | - | expression tag | UNP O43602 |
| D | 128 | ARG | - | expression tag | UNP O43602 |
| D | 129 | GLY | - | expression tag | UNP O43602 |
| D | 130 | SER | - | expression tag | UNP O43602 |
| D | 131 | HIS | - | expression tag | UNP O43602 |
| D | 132 | MET | - | expression tag | UNP O43602 |
| D | 215 | ASP | LYS | engineered mutation | UNP O43602 |
| D | 216 | ASP | LYS | engineered mutation | UNP O43602 |
| E | 125 | LEU | - | expression tag | UNP O43602 |
| E | 126 | VAL | - | expression tag | UNP O43602 |
| E | 127 | PRO | - | expression tag | UNP O43602 |
| E | 128 | ARG | - | expression tag | UNP O43602 |
| E | 129 | GLY | - | expression tag | UNP O43602 |
| E | 130 | SER | - | expression tag | UNP O43602 |
| E | 131 | HIS | - | expression tag | UNP O43602 |
| E | 132 | MET | - | expression tag | UNP O43602 |
| E | 215 | ASP | LYS | engineered mutation | UNP O43602 |
| E | 216 | ASP | LYS | engineered mutation | UNP O43602 |
| F | 125 | LEU | - | expression tag | UNP O43602 |
| F | 126 | VAL | - | expression tag | UNP O43602 |
| F | 127 | PRO | - | expression tag | UNP O43602 |
| F | 128 | ARG | - | expression tag | UNP O43602 |
| F | 129 | GLY | - | expression tag | UNP O43602 |
| F | 130 | SER | - | expression tag | UNP O43602 |
| F | 131 | HIS | - | expression tag | UNP O43602 |
| F | 132 | MET | - | expression tag | UNP O43602 |
| F | 215 | ASP | LYS | engineered mutation | UNP O43602 |

Continued on next page...

Continued from previous page...

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|---------------------|------------|
| F | 216 | ASP | LYS | engineered mutation | UNP O43602 |


- Molecule 2 is water.

| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|------------------|---------|---------|
| 2 | A | 95 | Total O 95 95 | 0 | 0 |
| 2 | B | 50 | Total O 50 50 | 0 | 0 |
| 2 | C | 58 | Total O 58 58 | 0 | 0 |
| 2 | D | 26 | Total O 26 26 | 0 | 0 |
| 2 | E | 69 | Total O 69 69 | 0 | 0 |
| 2 | F | 55 | Total O 55 55 | 0 | 0 |

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Neuronal migration protein doublecortin

Chain A: 




- Molecule 1: Neuronal migration protein doublecortin

Chain B: 





- Molecule 1: Neuronal migration protein doublecortin

Chain C: 




- Molecule 1: Neuronal migration protein doublecortin

Chain D: 



- Molecule 1: Neuronal migration protein doublecortin

Chain E: 



- Molecule 1: Neuronal migration protein doublecortin

Chain F: 



4 Data and refinement statistics

| Property | Value | Source |
|---|---|------------------|
| Space group | P 61 2 2 | Depositor |
| Cell constants a, b, c, α , β , γ | 97.72Å 97.72Å 377.09Å 90.00° 90.00° 120.00° | Depositor |
| Resolution (Å) | 35.14 – 2.40 35.10 – 2.40 | Depositor EDS |
| % Data completeness (in resolution range) | 99.9 (35.14-2.40) 100.0 (35.10-2.40) | Depositor EDS |
| R_{merge} | (Not available) | Depositor |
| R_{sym} | 0.12 | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 1.10 (at 2.39Å) | Xtriage |
| Refinement program | BUSTER 2.9.2 | Depositor |
| R, R_{free} | 0.196 , 0.234 0.211 , 0.251 | Depositor DCC |
| R_{free} test set | 2200 reflections (5.41%) | DCC |
| Wilson B-factor (Å ²) | 42.0 | Xtriage |
| Anisotropy | 0.060 | Xtriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.36 , 56.1 | EDS |
| L-test for twinning ² | $\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$ | Xtriage |
| Estimated twinning fraction | No twinning to report. | Xtriage |
| F_o, F_c correlation | 0.94 | EDS |
| Total number of atoms | 5085 | wwPDB-VP |
| Average B, all atoms (Å ²) | 49.0 | wwPDB-VP |

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------------|-------------|---------------|
| | | RMSZ | $\# Z > 5$ | RMSZ | $\# Z > 5$ |
| 1 | A | 0.55 | 0/884 | 0.79 | 0/1193 |
| 1 | B | 0.49 | 0/729 | 0.72 | 0/980 |
| 1 | C | 0.48 | 0/799 | 0.72 | 0/1077 |
| 1 | D | 0.97 | 1/737 (0.1%) | 1.63 | 3/990 (0.3%) |
| 1 | E | 0.53 | 0/884 | 0.78 | 0/1193 |
| 1 | F | 0.52 | 0/793 | 0.77 | 0/1069 |
| All | All | 0.61 | 1/4826 (0.0%) | 0.95 | 3/6502 (0.0%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1 | D | 0 | 1 |

All (1) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|--------|-------------|----------|
| 1 | D | 132 | MET | C-N | -22.87 | 0.81 | 1.34 |

All (3) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|--------|--------|-------------|----------|
| 1 | D | 132 | MET | O-C-N | -35.17 | 66.43 | 122.70 |
| 1 | D | 132 | MET | CA-C-N | 24.65 | 171.43 | 117.20 |
| 1 | D | 132 | MET | C-N-CA | 17.91 | 166.47 | 121.70 |

There are no chirality outliers.

All (1) planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|-----------|
| 1 | D | 132 | MET | Mainchain |

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 866 | 0 | 829 | 14 | 0 |
| 1 | B | 716 | 0 | 679 | 7 | 0 |
| 1 | C | 783 | 0 | 741 | 4 | 0 |
| 1 | D | 724 | 0 | 687 | 4 | 0 |
| 1 | E | 866 | 0 | 829 | 8 | 0 |
| 1 | F | 777 | 0 | 736 | 10 | 0 |
| 2 | A | 95 | 0 | 0 | 0 | 0 |
| 2 | B | 50 | 0 | 0 | 1 | 0 |
| 2 | C | 58 | 0 | 0 | 2 | 0 |
| 2 | D | 26 | 0 | 0 | 0 | 0 |
| 2 | E | 69 | 0 | 0 | 0 | 0 |
| 2 | F | 55 | 0 | 0 | 1 | 0 |
| All | All | 5085 | 0 | 4501 | 34 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 34 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|------------------|--------------------------|-------------------|
| 1:A:230:ASN:HD22 | 1:E:150:VAL:H | 1.19 | 0.90 |
| 1:A:224:ASN:HD22 | 1:A:224:ASN:H | 1.27 | 0.81 |
| 1:B:147:LYS:HE3 | 1:B:147:LYS:H | 1.49 | 0.77 |
| 1:A:128:ARG:NH2 | 1:B:171:SER:OG | 2.26 | 0.68 |
| 1:A:150:VAL:H | 1:E:230:ASN:HD22 | 1.46 | 0.60 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|---------------|-----------|---------|----------|-------------|-----|
| 1 | A | 105/107 (98%) | 104 (99%) | 1 (1%) | 0 | 100 | 100 |
| 1 | B | 86/107 (80%) | 84 (98%) | 2 (2%) | 0 | 100 | 100 |
| 1 | C | 94/107 (88%) | 90 (96%) | 3 (3%) | 1 (1%) | 17 | 25 |
| 1 | D | 87/107 (81%) | 85 (98%) | 2 (2%) | 0 | 100 | 100 |
| 1 | E | 105/107 (98%) | 98 (93%) | 7 (7%) | 0 | 100 | 100 |
| 1 | F | 93/107 (87%) | 91 (98%) | 1 (1%) | 1 (1%) | 17 | 25 |
| All | All | 570/642 (89%) | 552 (97%) | 16 (3%) | 2 (0%) | 39 | 56 |

All (2) Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | C | 227 | TRP |
| 1 | F | 173 | SER |

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|--------------|-----------|----------|-------------|----|
| 1 | A | 96/96 (100%) | 89 (93%) | 7 (7%) | 17 | 27 |
| 1 | B | 78/96 (81%) | 72 (92%) | 6 (8%) | 16 | 24 |
| 1 | C | 86/96 (90%) | 77 (90%) | 9 (10%) | 8 | 12 |
| 1 | D | 79/96 (82%) | 72 (91%) | 7 (9%) | 12 | 18 |
| 1 | E | 96/96 (100%) | 90 (94%) | 6 (6%) | 22 | 35 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|---------------|-----------|----------|-------------|----|
| 1 | F | 85/96 (88%) | 78 (92%) | 7 (8%) | 14 | 21 |
| All | All | 520/576 (90%) | 478 (92%) | 42 (8%) | 15 | 22 |

5 of 42 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | C | 180 | GLN |
| 1 | D | 135 | LYS |
| 1 | F | 211 | ASP |
| 1 | C | 211 | ASP |
| 1 | C | 228 | SER |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | B | 180 | GLN |
| 1 | F | 226 | ASN |
| 1 | E | 230 | ASN |
| 1 | A | 230 | ASN |
| 1 | F | 180 | GLN |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

| Mol | Chain | Number of breaks |
|-----|-------|------------------|
| 1 | D | 1 |

All chain breaks are listed below:

| Model | Chain | Residue-1 | Atom-1 | Residue-2 | Atom-2 | Distance (Å) |
|-------|-------|-----------|--------|-----------|--------|--------------|
| 1 | D | 132:MET | C | 133:ALA | N | 0.81 |

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 |
|-----|-------|----------------|--------|---|-----------------------|-------|
| 1 | A | 107/107 (100%) | -0.46 | 0 100 100 | 24, 34, 53, 72 | 0 |
| 1 | B | 88/107 (82%) | 0.12 | 5 (5%) 27 27 | 30, 49, 86, 89 | 0 |
| 1 | C | 96/107 (89%) | -0.19 | 2 (2%) 67 66 | 30, 47, 78, 82 | 0 |
| 1 | D | 89/107 (83%) | 0.37 | 8 (8%) 12 11 | 33, 63, 94, 105 | 0 |
| 1 | E | 107/107 (100%) | -0.26 | 1 (0%) 85 85 | 27, 41, 67, 79 | 0 |
| 1 | F | 95/107 (88%) | -0.26 | 2 (2%) 67 66 | 31, 45, 74, 83 | 0 |
| All | All | 582/642 (90%) | -0.13 | 18 (3%) 52 52 | 24, 45, 81, 105 | 0 |

The worst 5 of 18 RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 1 | D | 213 | PHE | 4.8 |
| 1 | B | 213 | PHE | 4.3 |
| 1 | E | 213 | PHE | 3.9 |
| 1 | B | 176 | ILE | 3.8 |
| 1 | F | 212 | ASN | 3.5 |

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.