



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:30 AM GMT

PDB ID : 3IQJ  
Title : Crystal Structure of human 14-3-3 sigma in Complex with Raf1 peptide (10mer)  
Authors : Ottmann, C.; Weyand, M.  
Deposited on : 2009-08-20  
Resolution : 1.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1495 (1.20-1.08)
Clashscore	102246	1593 (1.20-1.08)
Ramachandran outliers	100387	1521 (1.20-1.08)
Sidechain outliers	100360	1518 (1.20-1.08)
RSRZ outliers	91569	1498 (1.20-1.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	
2	P	10	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2517 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 14-3-3 protein sigma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	236	Total	C	N	O	S	0	37	0
			2021	1290	321	394	16			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	EXPRESSION TAG	UNP P31947
A	-3	ALA	-	EXPRESSION TAG	UNP P31947
A	-2	MET	-	EXPRESSION TAG	UNP P31947
A	-1	GLY	-	EXPRESSION TAG	UNP P31947
A	0	SER	-	EXPRESSION TAG	UNP P31947

- Molecule 2 is a protein called 10-mer peptide from RAF proto-oncogene serine/threonine-protein kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	8	Total	C	N	O	P	0	1	0
			63	34	10	18	1			

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Mg	0	0
			3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	412	Total 412	O 412	0	0
5	P	17	Total 17	O 17	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 14-3-3 protein sigma



- Molecule 2: 10-mer peptide from RAF proto-oncogene serine/threonine-protein kinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.61Å 112.56Å 62.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.49 – 1.15 34.49 – 1.15	Depositor EDS
% Data completeness (in resolution range)	99.4 (34.49-1.15) 99.4 (34.49-1.15)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.69 (at 1.15Å)	Xtriage
Refinement program	REFMAC 5.5.0070	Depositor
R, $R_{free}$	0.127 , 0.153 0.127 , 0.150	Depositor DCC
$R_{free}$ test set	5105 reflections (5.22%)	DCC
Wilson B-factor (Å <sup>2</sup> )	11.0	Xtriage
Anisotropy	0.207	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 41.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 102954 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2517	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSO, CL, MG, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.27	19/2145 (0.9%)	1.09	10/2882 (0.3%)
2	P	1.29	0/55	1.24	0/75
All	All	1.27	19/2200 (0.9%)	1.09	10/2957 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	84	TYR	CD1-CE1	8.83	1.52	1.39
1	A	115[A]	GLU	CD-OE1	6.78	1.33	1.25
1	A	115[B]	GLU	CD-OE1	6.78	1.33	1.25
1	A	84	TYR	CG-CD1	-6.63	1.30	1.39
1	A	84	TYR	CD2-CE2	-6.63	1.29	1.39
1	A	256	GLU	CD-OE2	-6.51	1.18	1.25
1	A	0[A]	SER	CB-OG	6.23	1.50	1.42
1	A	0[B]	SER	CB-OG	6.23	1.50	1.42
1	A	115[A]	GLU	CG-CD	5.63	1.60	1.51
1	A	115[B]	GLU	CG-CD	5.63	1.60	1.51
1	A	115[A]	GLU	CD-OE2	5.45	1.31	1.25
1	A	115[B]	GLU	CD-OE2	5.45	1.31	1.25
1	A	145	ASP	CB-CG	5.42	1.63	1.51
1	A	2	GLU	CG-CD	-5.22	1.44	1.51
1	A	19	TYR	CE1-CZ	5.16	1.45	1.38
1	A	153	GLU	CD-OE2	5.12	1.31	1.25

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	213[A]	TYR	CD1-CE1	-5.07	1.31	1.39
1	A	213[B]	TYR	CD1-CE1	-5.07	1.31	1.39
1	A	86	GLU	CG-CD	5.04	1.59	1.51

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	82	ARG	NE-CZ-NH1	9.75	125.17	120.30
1	A	84	TYR	CD1-CE1-CZ	-7.87	112.71	119.80
1	A	129	ARG	NE-CZ-NH1	7.43	124.02	120.30
1	A	136	THR	C-N-CA	-7.24	107.09	122.30
1	A	84	TYR	CB-CG-CD1	-6.69	116.99	121.00
1	A	86	GLU	OE1-CD-OE2	-6.23	115.82	123.30
1	A	3	ARG	NE-CZ-NH1	5.87	123.24	120.30
1	A	49	LYS	CD-CE-NZ	5.47	124.28	111.70
1	A	256	GLU	OE1-CD-OE2	-5.16	117.11	123.30
1	A	136	THR	O-C-N	-5.08	114.56	123.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	137	GLY	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2021	0	2084	58	3
2	P	63	0	51	22	0
3	A	1	0	0	0	0
4	A	3	0	0	0	0
5	A	412	0	0	25	10
5	P	17	0	0	11	0
All	All	2517	0	2135	68	10

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 16.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:ASP:HB2	5:A:400:HOH:O	1.41	1.20
2:P:258:THR:HB	5:P:344:HOH:O	1.42	1.19
1:A:200[A]:GLU:HG3	5:A:352:HOH:O	1.43	1.15
1:A:77:LYS:O	5:A:420:HOH:O	1.67	1.11
1:A:52[B]:VAL:HG13	1:A:92[B]:LEU:CD1	1.85	1.06
2:P:260[A]:THR:HG23	5:P:404:HOH:O	0.89	1.05
1:A:229[B]:LEU:HD21	2:P:255:GLN:OE1	1.57	1.03
1:A:86:GLU:HG2	5:A:367:HOH:O	1.64	0.98
1:A:42:ASN:HB3	5:A:398:HOH:O	1.65	0.96
1:A:177[A]:SER:OG	5:A:414:HOH:O	1.84	0.93
1:A:195[B]:LYS:HE2	5:A:358:HOH:O	1.67	0.92
1:A:52[B]:VAL:HG13	1:A:92[B]:LEU:HD11	1.49	0.92
1:A:175:ASN:OD1	2:P:260[B]:THR:HG23	1.71	0.89
2:P:260[B]:THR:HB	5:P:404:HOH:O	1.76	0.85
1:A:19:TYR:CE1	1:A:50[B]:ASN:ND2	2.48	0.81
1:A:119[B]:PHE:CE2	1:A:123[B]:MET:CE	2.67	0.78
1:A:52[B]:VAL:CG1	1:A:92[B]:LEU:HD11	2.13	0.77
2:P:260[B]:THR:CB	5:P:404:HOH:O	2.30	0.76
1:A:119[B]:PHE:CE2	1:A:123[B]:MET:HE1	2.22	0.75
1:A:42:ASN:OD1	5:A:397:HOH:O	2.05	0.74
1:A:52[B]:VAL:HG13	1:A:92[B]:LEU:HD12	1.70	0.73
1:A:42:ASN:CG	5:A:397:HOH:O	2.28	0.72
1:A:229[B]:LEU:HD23	5:P:204:HOH:O	1.87	0.72
1:A:229[B]:LEU:CD2	2:P:255:GLN:OE1	2.37	0.70
1:A:42:ASN:CB	5:A:398:HOH:O	2.30	0.69
1:A:77:LYS:CG	5:A:512:HOH:O	2.40	0.69
2:P:258:THR:CB	5:P:344:HOH:O	2.18	0.69
1:A:-3:ALA:O	5:A:349:HOH:O	2.12	0.67
1:A:77:LYS:HE2	5:A:561:HOH:O	1.95	0.66
2:P:260[A]:THR:CG2	5:P:404:HOH:O	1.69	0.65
2:P:260[A]:THR:CB	5:P:404:HOH:O	2.24	0.63
1:A:77:LYS:HG3	5:A:512:HOH:O	1.96	0.63
1:A:202[B]:MET:HE1	5:A:382:HOH:O	2.01	0.61
1:A:171:GLY:O	2:P:260[B]:THR:HG21	2.01	0.60
1:A:174:LEU:HD23	2:P:260[B]:THR:HG22	1.82	0.60
1:A:195[B]:LYS:CE	5:A:358:HOH:O	2.34	0.60
1:A:218:LEU:O	1:A:222[B]:LEU:HD13	2.07	0.55
1:A:89:GLU:OE2	5:A:346:HOH:O	2.18	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:ASN:CG	5:A:398:HOH:O	2.48	0.52
1:A:182[B]:GLU:OE2	2:P:257:SER:HB2	2.10	0.52
1:A:50[A]:ASN:HD21	2:P:262:ASN:HD21	1.59	0.51
2:P:262:ASN:ND2	5:P:395:HOH:O	2.38	0.51
1:A:60[A]:ARG:NH2	5:A:392:HOH:O	1.91	0.48
1:A:119[B]:PHE:CE2	1:A:123[B]:MET:HE3	2.49	0.48
1:A:50[A]:ASN:ND2	2:P:262:ASN:HD21	2.12	0.48
1:A:119[B]:PHE:CZ	1:A:123[B]:MET:HE1	2.49	0.47
1:A:174:LEU:HD23	2:P:260[B]:THR:CG2	2.43	0.47
2:P:260[B]:THR:CG2	5:P:404:HOH:O	2.60	0.47
1:A:153:GLU:HG3	5:A:535:HOH:O	2.15	0.47
1:A:119[B]:PHE:CZ	1:A:123[B]:MET:CE	2.98	0.46
1:A:19:TYR:CZ	1:A:50[B]:ASN:ND2	2.84	0.46
2:P:260[B]:THR:HG22	5:P:404:HOH:O	2.16	0.45
1:A:119[B]:PHE:HE2	1:A:123[B]:MET:HE1	1.74	0.45
1:A:162[B]:MET:HB3	1:A:162[B]:MET:HE2	1.93	0.45
1:A:218:LEU:C	1:A:218:LEU:HD23	2.37	0.44
1:A:222[B]:LEU:HD21	2:P:261:PRO:HD3	1.99	0.44
1:A:200[A]:GLU:CG	5:A:352:HOH:O	2.26	0.44
1:A:155[B]:MET:SD	1:A:159:LYS:HE3	2.56	0.44
1:A:222[B]:LEU:CD2	2:P:261:PRO:HD3	2.48	0.44
1:A:165:THR:HG23	1:A:204:ASP:HB3	2.01	0.43
1:A:77:LYS:HE3	5:A:512:HOH:O	2.19	0.43
1:A:52[B]:VAL:CG2	1:A:95:VAL:HG11	2.48	0.43
1:A:174:LEU:HD12	5:A:414:HOH:O	2.19	0.43
1:A:52[B]:VAL:CG1	1:A:92[B]:LEU:CD1	2.73	0.42
1:A:50[A]:ASN:HD21	2:P:262:ASN:ND2	2.16	0.42
1:A:86:GLU:HG2	5:A:238:HOH:O	2.21	0.41
1:A:206:HIS:HD2	5:A:538:HOH:O	2.04	0.40

All (10) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:311:HOH:O	5:A:452:HOH:O[4_555]	1.39	0.81
1:A:142:ARG:CD	5:A:366:HOH:O[3_455]	1.55	0.65
5:A:362:HOH:O	5:A:555:HOH:O[6_444]	1.88	0.32
1:A:259[B]:CSO:OD	5:A:483:HOH:O[6_444]	1.99	0.21
5:A:277:HOH:O	5:A:277:HOH:O[3_454]	2.05	0.15
5:A:382:HOH:O	5:A:415:HOH:O[3_555]	2.07	0.13

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:271:HOH:O	5:A:499:HOH:O[8_445]	2.12	0.08
1:A:259[B]:CSO:SG	5:A:516:HOH:O[6_444]	2.12	0.08
5:A:345:HOH:O	5:A:407:HOH:O[4_555]	2.14	0.06
5:A:464:HOH:O	5:A:532:HOH:O[3_454]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	269/236 (114%)	266 (99%)	3 (1%)	0	100	100
2	P	6/10 (60%)	6 (100%)	0	0	100	100
All	All	275/246 (112%)	272 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/197 (114%)	224 (100%)	0	100	100
2	P	7/9 (78%)	7 (100%)	0	100	100
All	All	231/206 (112%)	231 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	206	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CSO	A	259[A]	-	3,6,7	0.59	0	1,6,8	1.64	0
1	CSO	A	259[B]	-	3,6,7	0.53	0	1,6,8	1.80	0
2	SEP	P	259	2	8,9,10	0.74	0	8,12,14	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	259[A]	-	-	0/1/5/7	0/0/0/0
1	CSO	A	259[B]	-	-	0/1/5/7	0/0/0/0
2	SEP	P	259	2	-	0/6/8/10	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	259[B]	CSO	0	2

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	235/236 (99%)	0.31	18 (7%)	16 17	7, 12, 27, 49	5 (2%)
2	P	7/10 (70%)	3.48	5 (71%)	0 0	18, 20, 30, 44	0
All	All	242/246 (98%)	0.40	23 (9%)	10 12	7, 12, 28, 49	5 (2%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	P	255	GLN	8.5
1	A	73	GLY	6.7
1	A	74	SER	6.3
1	A	72	GLU	5.3
1	A	137	GLY	5.0
1	A	70	ASN	4.6
1	A	139	ASP	4.1
1	A	138	ASP	4.0
1	A	71	GLU	4.0
2	P	261	PRO	3.9
2	P	262	ASN	3.6
1	A	67[A]	GLN	3.6
1	A	75	GLU	3.6
2	P	256	ARG	3.4
1	A	-4	GLY	3.4
1	A	136	THR	3.1
1	A	69	SER	3.0
1	A	-3	ALA	2.8
1	A	77	LYS	2.6
2	P	260[A]	THR	2.2
1	A	231	THR	2.1
1	A	211	ASP	2.1
1	A	76	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	CSO	A	259[B]	7/8	0.97	0.10	-	9,10,22,26	4
2	SEP	P	259	10/11	0.98	0.08	-	11,13,15,16	0
1	CSO	A	259[A]	7/8	0.97	0.10	-	9,9,17,23	4

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	CL	A	263	1/1	1.00	0.06	-1.86	14,14,14,14	0
4	MG	A	266	1/1	0.99	0.08	-	10,10,10,10	0
4	MG	A	265	1/1	1.00	0.07	-	10,10,10,10	1
4	MG	A	264	1/1	0.98	0.16	-	18,18,18,18	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.