



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:35 PM GMT

PDB ID : 4IRU  
Title : Crystal Structure of lepB GAP core in a transition state mimetic complex with Rab1A and ALF3  
Authors : Mishra, A.K.; Delcampo, C.M.; Collins, R.E.; Roy, C.R.; Lambright, D.G.  
Deposited on : 2013-01-15  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1124 (3.24-3.16)
Clashscore	102246	1024 (3.22-3.18)
Ramachandran outliers	100387	1004 (3.22-3.18)
Sidechain outliers	100360	1003 (3.22-3.18)
RSRZ outliers	91569	1129 (3.24-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	299	<div> <div>67%</div> <div>29%</div> <div>..</div> </div>
1	C	299	<div> <div>75%</div> <div>22%</div> <div>..</div> </div>
1	E	299	<div> <div>17%</div> <div>53%</div> <div>39%</div> <div>6%</div> <div>..</div> </div>
2	B	184	<div> <div>2%</div> <div>60%</div> <div>30%</div> <div>..</div> <div>8%</div> </div>
2	D	184	<div> <div>65%</div> <div>23%</div> <div>.</div> <div>8%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	184	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	B	204	-	-	-	X
3	ACT	B	205	-	-	-	X
3	ACT	C	701	-	-	-	X
3	ACT	C	702	-	-	-	X
3	ACT	C	703	-	-	-	X
3	ACT	C	704	-	-	-	X
3	ACT	C	705	-	-	X	X
4	GOL	A	904	-	-	-	X
4	GOL	A	906	-	-	-	X
4	GOL	A	907	-	-	-	X
4	GOL	A	908	-	-	-	X
4	GOL	B	206	-	-	-	X
5	K	A	911	-	-	-	X
8	AF3	B	203	-	-	-	X

## 2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 11410 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LepB.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	296	Total	C	N	O	S	Se	0	0	0
			2365	1526	389	443	4	3			
1	C	296	Total	C	N	O	S	Se	0	0	0
			2365	1526	389	443	4	3			
1	E	295	Total	C	N	O	S	Se	0	0	0
			2359	1523	388	441	4	3			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	325	SER	-	EXPRESSION TAG	UNP Q5ZSM7
C	325	SER	-	EXPRESSION TAG	UNP Q5ZSM7
E	325	SER	-	EXPRESSION TAG	UNP Q5ZSM7

- Molecule 2 is a protein called Ras-related protein Rab-1A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	170	Total	C	N	O	S	Se	0	0	0
			1362	868	222	267	2	3			
2	D	170	Total	C	N	O	S	Se	0	0	0
			1362	868	222	267	2	3			
2	F	170	Total	C	N	O	S	Se	0	0	0
			1362	868	222	267	2	3			

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	MSE	-	EXPRESSION TAG	UNP P62820
B	-5	GLY	-	EXPRESSION TAG	UNP P62820
B	-4	HIS	-	EXPRESSION TAG	UNP P62820
B	-3	HIS	-	EXPRESSION TAG	UNP P62820
B	-2	HIS	-	EXPRESSION TAG	UNP P62820

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	HIS	-	EXPRESSION TAG	UNP P62820
B	0	HIS	-	EXPRESSION TAG	UNP P62820
B	1	HIS	-	EXPRESSION TAG	UNP P62820
B	2	GLY	-	EXPRESSION TAG	UNP P62820
B	3	SER	-	EXPRESSION TAG	UNP P62820
D	-6	MSE	-	EXPRESSION TAG	UNP P62820
D	-5	GLY	-	EXPRESSION TAG	UNP P62820
D	-4	HIS	-	EXPRESSION TAG	UNP P62820
D	-3	HIS	-	EXPRESSION TAG	UNP P62820
D	-2	HIS	-	EXPRESSION TAG	UNP P62820
D	-1	HIS	-	EXPRESSION TAG	UNP P62820
D	0	HIS	-	EXPRESSION TAG	UNP P62820
D	1	HIS	-	EXPRESSION TAG	UNP P62820
D	2	GLY	-	EXPRESSION TAG	UNP P62820
D	3	SER	-	EXPRESSION TAG	UNP P62820
F	-6	MSE	-	EXPRESSION TAG	UNP P62820
F	-5	GLY	-	EXPRESSION TAG	UNP P62820
F	-4	HIS	-	EXPRESSION TAG	UNP P62820
F	-3	HIS	-	EXPRESSION TAG	UNP P62820
F	-2	HIS	-	EXPRESSION TAG	UNP P62820
F	-1	HIS	-	EXPRESSION TAG	UNP P62820
F	0	HIS	-	EXPRESSION TAG	UNP P62820
F	1	HIS	-	EXPRESSION TAG	UNP P62820
F	2	GLY	-	EXPRESSION TAG	UNP P62820
F	3	SER	-	EXPRESSION TAG	UNP P62820

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0


- Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total K 1 1	0	0
5	A	4	Total K 4 4	0	0
5	C	1	Total K 1 1	0	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total Mg 1 1	0	0
6	D	1	Total Mg 1 1	0	0
6	F	1	Total Mg 1 1	0	0

- Molecule 7 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>11</sub>P<sub>2</sub>).



- 
- Diagram illustrating the AF3 molecule structure. The central atom is Aluminum (Al), labeled AL. It is bonded to three Fluorine (F) atoms, labeled F1, F2, and F3. The bonds are shown as lines connecting the central Al atom to the F atoms.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Al	F	0	0
			4	1	3		





*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	D	1	Total	Al	F	0	0
			4	1	3		
8	F	1	Total	Al	F	0	0
			4	1	3		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	8	Total	O	0	0
			8	8		
9	B	8	Total	O	0	0
			8	8		
9	C	12	Total	O	0	0
			12	12		
9	D	3	Total	O	0	0
			3	3		
9	F	3	Total	O	0	0
			3	3		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: LepB




#### • Molecule 1: LepB




#### • Molecule 1: LepB







• Molecule 2: Ras-related protein Rab-1A



• Molecule 2: Ras-related protein Rab-1A



• Molecule 2: Ras-related protein Rab-1A



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	139.40 Å   139.40 Å   384.52 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	43.01 – 3.20 42.97 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (43.01-3.20) 100.0 (42.97-3.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.54	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.87 (at 3.19 Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.230   ,   0.288 0.230   ,   0.290	Depositor DCC
$R_{free}$ test set	1602 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	54.4	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 51.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 31774 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	11410	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, GOL, MG, K, ACT, AF3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	0/2415	0.71	0/3262
1	C	0.55	0/2415	0.71	1/3262 (0.0%)
1	E	0.47	0/2408	0.67	1/3251 (0.0%)
2	B	0.46	0/1381	0.67	0/1859
2	D	0.42	0/1381	0.66	0/1859
2	F	0.45	0/1381	0.68	1/1859 (0.1%)
All	All	0.50	0/11381	0.69	3/15352 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	16	LEU	CA-CB-CG	5.26	127.41	115.30
1	C	332	LEU	CA-CB-CG	5.18	127.22	115.30
1	E	429	LEU	CA-CB-CG	5.06	126.94	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2365	0	2378	50	0
1	C	2365	0	2378	44	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2359	0	2372	110	2
2	B	1362	0	1358	40	0
2	D	1362	0	1358	41	0
2	F	1362	0	1358	80	0
3	A	8	0	6	0	0
3	B	8	0	6	0	0
3	C	24	0	18	4	0
3	D	8	0	6	0	0
4	A	36	0	48	1	0
4	B	6	0	8	3	0
4	C	6	0	8	0	0
5	A	4	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
6	B	1	0	0	0	0
6	D	1	0	0	0	0
6	F	1	0	0	0	0
7	B	28	0	12	4	0
7	D	28	0	12	8	0
7	F	28	0	12	4	0
8	B	4	0	0	1	0
8	D	4	0	0	1	0
8	F	4	0	0	0	0
9	A	8	0	0	0	0
9	B	8	0	0	0	0
9	C	12	0	0	0	0
9	D	3	0	0	0	0
9	F	3	0	0	0	0
All	All	11410	0	11338	353	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 353 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:53:ILE:HD11	2:B:166:MSE:SE	1.92	1.19
1:E:456:CYS:HA	1:E:465:VAL:HG22	1.44	1.00
1:E:479:ASN:O	1:E:483:ILE:HG12	1.62	0.99
1:C:393:LYS:HG3	1:C:524:PRO:HB3	1.46	0.97
2:F:118:ASN:HD21	2:F:175:ARG:HH21	1.15	0.95

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:395:ARG:O	1:E:395:ARG:O[6_555]	1.49	0.71
1:E:396:ILE:O	1:E:396:ILE:O[6_555]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/299 (98%)	267 (91%)	23 (8%)	4 (1%)	14	57
1	C	294/299 (98%)	273 (93%)	19 (6%)	2 (1%)	26	72
1	E	291/299 (97%)	231 (79%)	44 (15%)	16 (6%)	2	18
2	B	168/184 (91%)	150 (89%)	16 (10%)	2 (1%)	16	60
2	D	168/184 (91%)	149 (89%)	16 (10%)	3 (2%)	11	51
2	F	168/184 (91%)	132 (79%)	28 (17%)	8 (5%)	3	22
All	All	1383/1449 (95%)	1202 (87%)	146 (11%)	35 (2%)	7	41

5 of 35 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	445	PHE
1	E	596	ARG
1	E	598	ASP
2	F	24	LYS
2	F	46	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	259/257 (101%)	234 (90%)	25 (10%)	10	39
1	C	259/257 (101%)	240 (93%)	19 (7%)	17	57
1	E	258/257 (100%)	230 (89%)	28 (11%)	8	33
2	B	150/156 (96%)	139 (93%)	11 (7%)	17	57
2	D	150/156 (96%)	142 (95%)	8 (5%)	28	69
2	F	150/156 (96%)	132 (88%)	18 (12%)	6	28
All	All	1226/1239 (99%)	1117 (91%)	109 (9%)	12	44

5 of 109 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	522	LYS
2	D	138	THR
2	F	85	HIS
1	C	526	PHE
2	D	94	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	160	ASN
1	E	441	ASN
2	F	118	ASN
2	D	157	ASN
2	F	70	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 35 ligands modelled in this entry, 9 are monoatomic - leaving 26 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	ACT	A	901	-	1,3,3	2.12	1 (100%)	0,3,3	0.00	-
3	ACT	A	902	5	1,3,3	1.79	0	0,3,3	0.00	-
4	GOL	A	903	-	5,5,5	0.45	0	5,5,5	0.81	0
4	GOL	A	904	-	5,5,5	0.32	0	5,5,5	0.43	0
4	GOL	A	905	-	5,5,5	0.43	0	5,5,5	0.51	0
4	GOL	A	906	-	5,5,5	0.59	0	5,5,5	0.81	0
4	GOL	A	907	-	5,5,5	0.36	0	5,5,5	0.29	0
4	GOL	A	908	-	5,5,5	0.26	0	5,5,5	0.29	0
7	GDP	B	202	6	23,30,30	1.34	3 (13%)	30,47,47	1.82	6 (20%)
8	AF3	B	203	-	0,3,3	0.00	-	0,3,3	0.00	-
3	ACT	B	204	-	1,3,3	1.65	0	0,3,3	0.00	-
3	ACT	B	205	-	1,3,3	1.54	0	0,3,3	0.00	-
4	GOL	B	206	-	5,5,5	0.48	0	5,5,5	0.45	0
3	ACT	C	701	-	1,3,3	2.14	1 (100%)	0,3,3	0.00	-
3	ACT	C	702	-	1,3,3	1.83	0	0,3,3	0.00	-
3	ACT	C	703	-	1,3,3	2.07	1 (100%)	0,3,3	0.00	-
3	ACT	C	704	-	1,3,3	1.55	0	0,3,3	0.00	-
3	ACT	C	705	-	1,3,3	1.45	0	0,3,3	0.00	-
3	ACT	C	706	-	1,3,3	1.70	0	0,3,3	0.00	-
4	GOL	C	707	-	5,5,5	0.44	0	5,5,5	0.17	0
7	GDP	D	202	6	23,30,30	1.44	3 (13%)	30,47,47	1.74	6 (20%)
8	AF3	D	203	-	0,3,3	0.00	-	0,3,3	0.00	-
3	ACT	D	204	-	1,3,3	1.65	0	0,3,3	0.00	-
3	ACT	D	205	-	1,3,3	1.69	0	0,3,3	0.00	-
7	GDP	F	201	6	23,30,30	1.34	2 (8%)	30,47,47	1.72	5 (16%)
8	AF3	F	202	-	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACT	A	901	-	-	0/0/0/0	0/0/0/0
3	ACT	A	902	5	-	0/0/0/0	0/0/0/0
4	GOL	A	903	-	-	0/4/4/4	0/0/0/0
4	GOL	A	904	-	-	0/4/4/4	0/0/0/0
4	GOL	A	905	-	-	0/4/4/4	0/0/0/0
4	GOL	A	906	-	-	0/4/4/4	0/0/0/0
4	GOL	A	907	-	-	0/4/4/4	0/0/0/0
4	GOL	A	908	-	-	0/4/4/4	0/0/0/0
7	GDP	B	202	6	-	0/12/32/32	0/3/3/3
8	AF3	B	203	-	-	0/0/0/0	0/0/0/0
3	ACT	B	204	-	-	0/0/0/0	0/0/0/0
3	ACT	B	205	-	-	0/0/0/0	0/0/0/0
4	GOL	B	206	-	-	0/4/4/4	0/0/0/0
3	ACT	C	701	-	-	0/0/0/0	0/0/0/0
3	ACT	C	702	-	-	0/0/0/0	0/0/0/0
3	ACT	C	703	-	-	0/0/0/0	0/0/0/0
3	ACT	C	704	-	-	0/0/0/0	0/0/0/0
3	ACT	C	705	-	-	0/0/0/0	0/0/0/0
3	ACT	C	706	-	-	0/0/0/0	0/0/0/0
4	GOL	C	707	-	-	0/4/4/4	0/0/0/0
7	GDP	D	202	6	-	0/12/32/32	0/3/3/3
8	AF3	D	203	-	-	0/0/0/0	0/0/0/0
3	ACT	D	204	-	-	0/0/0/0	0/0/0/0
3	ACT	D	205	-	-	0/0/0/0	0/0/0/0
7	GDP	F	201	6	-	0/12/32/32	0/3/3/3
8	AF3	F	202	-	-	0/0/0/0	0/0/0/0

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	D	202	GDP	O4'-C1'	2.03	1.43	1.41
3	C	703	ACT	CH3-C	2.07	1.51	1.48
3	A	901	ACT	CH3-C	2.12	1.51	1.48
3	C	701	ACT	CH3-C	2.14	1.51	1.48
7	B	202	GDP	O4'-C1'	2.25	1.44	1.41

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
7	B	202	GDP	C5-C6-N1	-4.45	117.50	123.59
7	D	202	GDP	C4-C5-N7	-4.44	105.40	109.48
7	D	202	GDP	C5-C6-N1	-4.16	117.89	123.59
7	B	202	GDP	C4-C5-N7	-4.02	105.78	109.48
7	F	201	GDP	C5-C6-N1	-3.75	118.46	123.59

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 26 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	907	GOL	1	0
7	B	202	GDP	4	0
8	B	203	AF3	1	0
4	B	206	GOL	3	0
3	C	705	ACT	4	0
7	D	202	GDP	8	0
8	D	203	AF3	1	0
7	F	201	GDP	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	293/299 (97%)	-0.41	3 (1%) 84 75	15, 36, 87, 158	0
1	C	293/299 (97%)	-0.43	2 (0%) 89 83	14, 36, 76, 112	0
1	E	292/299 (97%)	1.11	51 (17%) 2 1	55, 116, 178, 213	0
2	B	167/184 (90%)	-0.21	4 (2%) 62 47	17, 46, 96, 128	0
2	D	167/184 (90%)	-0.07	2 (1%) 81 69	22, 57, 100, 127	0
2	F	167/184 (90%)	2.17	71 (42%) 0 0	59, 133, 185, 213	0
All	All	1379/1449 (95%)	0.29	133 (9%) 10 6	14, 57, 162, 213	0

The worst 5 of 133 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	149	PRO	13.5
2	F	53	ILE	12.2
2	F	120	LEU	12.0
2	F	52	THR	8.2
2	F	148	ILE	8.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	ACT	C	705	4/4	0.91	0.74	14.36	88,88,94,100	0
4	GOL	B	206	6/6	0.87	0.60	6.99	58,74,76,77	0
3	ACT	B	205	4/4	0.89	0.27	6.54	63,66,68,75	0
3	ACT	C	701	4/4	0.90	0.39	5.37	59,61,65,67	0
3	ACT	C	703	4/4	0.77	0.35	4.69	82,88,90,94	0
4	GOL	A	907	6/6	0.86	0.36	4.18	80,94,97,98	0
3	ACT	C	702	4/4	0.90	0.30	3.98	60,63,65,65	0
4	GOL	A	908	6/6	0.84	0.45	3.83	88,96,97,99	0
4	GOL	A	904	6/6	0.90	0.44	3.64	61,68,73,78	0
5	K	A	911	1/1	0.80	0.22	2.94	90,90,90,90	0
4	GOL	A	906	6/6	0.87	0.28	2.67	43,47,51,53	0
3	ACT	B	204	4/4	0.86	0.32	2.54	84,87,89,94	0
3	ACT	C	704	4/4	0.79	0.38	2.41	65,75,79,79	0
8	AF3	B	203	4/4	0.97	0.24	2.04	31,32,33,33	0
3	ACT	A	902	4/4	0.88	0.22	1.20	66,68,72,76	0
7	GDP	B	202	28/28	0.97	0.20	0.64	19,32,43,47	0
3	ACT	D	204	4/4	0.88	0.24	0.41	64,66,69,72	0
8	AF3	D	203	4/4	0.98	0.18	-0.18	29,30,30,30	0
7	GDP	D	202	28/28	0.97	0.17	-0.26	22,29,38,41	0
5	K	A	912	1/1	0.77	0.23	-0.34	85,85,85,85	0
4	GOL	A	903	6/6	0.87	0.19	-0.37	44,50,51,52	0
6	MG	B	201	1/1	0.90	0.18	-0.62	32,32,32,32	0
7	GDP	F	201	28/28	0.88	0.27	-1.20	84,111,172,173	0
8	AF3	F	202	4/4	0.90	0.28	-1.33	56,60,61,66	0
6	MG	D	201	1/1	0.97	0.16	-1.39	36,36,36,36	0
6	MG	F	200	1/1	0.98	0.10	-5.10	45,45,45,45	0
4	GOL	C	707	6/6	0.83	0.36	-	73,79,85,86	0
3	ACT	C	706	4/4	0.79	0.28	-	98,100,102,103	0
3	ACT	A	901	4/4	0.91	0.20	-	51,54,56,57	0
3	ACT	D	205	4/4	0.89	0.25	-	64,69,71,73	0
5	K	A	910	1/1	0.85	0.23	-	101,101,101,101	0
5	K	A	909	1/1	0.89	0.15	-	85,85,85,85	0
4	GOL	A	905	6/6	0.72	0.42	-	94,97,102,104	0
5	K	C	708	1/1	0.96	0.14	-	72,72,72,72	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	K	B	207	1/1	0.96	0.13	-	52,52,52,52	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.