



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 08:07 PM GMT

PDB ID : 1IUG  
Title : The crystal structure of aspartate aminotransferase which belongs to subgroup IV from *Thermus thermophilus*  
Authors : Katsura, Y.; Shirouzu, M.; Yamaguchi, H.; Ishitani, R.; Nureki, O.; Kuramitsu, S.; Hayashi, H.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2002-03-04  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry. We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org). A user guide is available at <http://wwpdb.org/validation/2016/XrayValidationReportHelp> with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	 78% 17% ..
1	B	352	 78% 17% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	B	3693	-	-	X	-
2	PO4	B	3694	-	-	-	X

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 5540 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called putative aspartate aminotransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
1	A	348	Total 2669	C 1717	N 460	O 484	P 1	S 7	0	0	0
1	B	348	Total 2669	C 1717	N 460	O 484	P 1	S 7	0	0	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	B	1	Total 5	O 4	P 1	0	0
2	B	1	Total 5	O 4	P 1	0	0


- Molecule 3 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	97	Total 97	O 97	0	0
3	B	95	Total 95	O 95	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: putative aspartate aminotransferase



- Molecule 1: putative aspartate aminotransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.85Å 135.33Å 133.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 34.67 – 2.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-2.20) 97.5 (34.67-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.06 (at 2.20Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.199 , 0.236 0.203 , 0.205	Depositor DCC
$R_{free}$ test set	2119 reflections (5.06%)	DCC
Wilson B-factor (Å <sup>2</sup> )	32.8	Xtrriage
Anisotropy	0.364	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 38.2	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	0 of 41892 reflections	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5540	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.34	0/2701	0.69	0/3663
1	B	0.43	2/2701 (0.1%)	0.68	1/3663 (0.0%)
All	All	0.39	2/5402 (0.0%)	0.68	1/7326 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	90	PHE	CB-CG	-11.09	1.32	1.51
1	B	322	ARG	C-N	-5.62	1.21	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	322	ARG	C-N-CA	5.17	134.62	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2669	0	2721	57	0
1	B	2669	0	2720	66	0
2	B	10	0	0	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	97	0	0	3	0
3	B	95	0	0	5	0
All	All	5540	0	5441	122	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (122) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:ARG:CG	1:A:89:ARG:HH11	1.69	1.06
1:A:89:ARG:HG3	1:A:89:ARG:HH11	0.92	1.05
1:A:89:ARG:HG3	1:A:89:ARG:NH1	1.64	1.01
1:A:103:ARG:HH11	1:A:103:ARG:HG3	1.38	0.87
1:B:28:ARG:HH22	2:B:3693:PO4:P	2.00	0.85
1:A:44:ARG:HG3	1:A:44:ARG:HH11	1.46	0.80
1:B:170:ALA:HB3	1:B:174:MET:HG3	1.63	0.78
1:B:40:ARG:HH11	1:B:40:ARG:HG3	1.47	0.78
1:A:170:ALA:HB3	1:A:174:MET:HG3	1.69	0.73
1:A:165:LEU:HD22	1:A:187:LEU:HD23	1.70	0.73
1:A:103:ARG:NH1	1:A:103:ARG:HG3	2.02	0.73
1:A:44:ARG:HG3	1:A:44:ARG:NH1	2.03	0.71
1:B:27:HIS:NE2	1:B:28:ARG:NE	2.40	0.69
1:A:40:ARG:HG3	1:A:40:ARG:HH11	1.57	0.69
1:A:267:VAL:HG13	1:A:342:ARG:HD3	1.74	0.69
1:B:26:HIS:CE1	1:B:28:ARG:HD2	2.27	0.69
1:A:13:HIS:HD2	1:A:15:LYS:H	1.41	0.69
1:B:85:LYS:HE2	1:B:89:ARG:NH2	2.09	0.68
1:A:86:PHE:HA	1:A:89:ARG:HB2	1.75	0.68
1:A:279:ARG:HG3	3:A:2078:HOH:O	1.95	0.67
1:A:33:ARG:HH11	1:A:33:ARG:HG3	1.60	0.66
1:A:85:LYS:NZ	2:B:3693:PO4:O2	2.29	0.65
1:A:48:ARG:HB2	1:A:170:ALA:HA	1.78	0.65
1:A:69:LYS:HD2	1:A:209:ARG:NE	2.11	0.64
1:A:119:LYS:HB3	1:A:122:TYR:HE1	1.64	0.61
1:A:162:THR:HG22	1:A:185:LLP:HG2	1.84	0.59
1:A:185:LLP:H4'1	3:B:2102:HOH:O	2.01	0.59
1:B:103:ARG:HG3	1:B:103:ARG:HH11	1.68	0.59
1:B:69:LYS:HD2	1:B:209:ARG:NE	2.18	0.58
1:B:85:LYS:CE	1:B:89:ARG:HH21	2.15	0.58
1:B:33:ARG:HH11	1:B:33:ARG:HG3	1.68	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:LEU:HG	1:B:24:GLN:HB2	1.86	0.58
1:A:213:LEU:HD22	3:B:2116:HOH:O	2.03	0.58
1:B:29:THR:O	1:B:33:ARG:HG2	2.05	0.56
1:A:274:ARG:HD3	1:A:275:PRO:HD2	1.88	0.56
1:B:26:HIS:HE1	1:B:28:ARG:HD2	1.70	0.56
1:B:33:ARG:NH1	1:B:33:ARG:HG3	2.21	0.56
1:A:113:ARG:HH11	1:A:113:ARG:HG3	1.72	0.55
1:B:37:LEU:O	1:B:40:ARG:HB2	2.07	0.55
1:B:36:PHE:O	1:B:40:ARG:HG2	2.07	0.54
1:A:325:LEU:HD21	1:A:337:VAL:HG21	1.89	0.54
1:B:85:LYS:CE	1:B:89:ARG:NH2	2.71	0.54
1:A:26:HIS:O	1:A:29:THR:HG22	2.08	0.53
1:B:28:ARG:NH2	2:B:3693:PO4:P	2.74	0.53
1:B:217:ARG:HG2	1:B:217:ARG:HH11	1.74	0.53
1:B:35:VAL:HA	1:B:38:LYS:CE	2.38	0.53
1:B:48:ARG:HB2	1:B:170:ALA:HA	1.90	0.52
1:B:27:HIS:CD2	1:B:28:ARG:HE	2.28	0.52
1:B:103:ARG:NH1	1:B:103:ARG:HG3	2.23	0.52
1:A:24:GLN:HB2	1:B:12:LEU:HG	1.92	0.52
1:B:342:ARG:HH11	1:B:342:ARG:HG2	1.75	0.51
1:A:33:ARG:NH1	1:A:33:ARG:HG3	2.20	0.51
1:B:217:ARG:HG3	3:B:2044:HOH:O	2.11	0.51
1:B:342:ARG:CG	1:B:342:ARG:HH11	2.24	0.51
1:B:85:LYS:HE3	1:B:89:ARG:HH21	1.74	0.51
1:A:301:ALA:O	1:A:305:ARG:HG2	2.12	0.50
1:B:217:ARG:HH11	1:B:217:ARG:CG	2.24	0.50
1:A:44:ARG:HD3	1:A:51:GLY:O	2.12	0.50
1:A:214:ASP:HB3	1:A:217:ARG:HH11	1.77	0.50
1:B:257:LYS:HD3	1:B:260:GLN:OE1	2.12	0.49
1:A:214:ASP:HB3	1:A:217:ARG:NH1	2.28	0.49
1:A:83:TYR:HB3	1:A:106:TYR:HB2	1.94	0.49
1:B:219:LEU:O	1:B:223:LYS:HG2	2.13	0.49
1:B:206:LEU:O	1:B:208:PRO:HD3	2.12	0.49
1:B:128:VAL:HA	1:B:159:ASP:HB3	1.93	0.49
1:A:162:THR:CG2	1:A:185:LLP:HG2	2.42	0.48
1:B:26:HIS:CE1	1:B:28:ARG:HB2	2.48	0.48
1:B:82:VAL:HB	1:B:103:ARG:HE	1.79	0.48
1:A:288:TYR:HA	1:A:320:VAL:HG12	1.96	0.48
1:B:35:VAL:HA	1:B:38:LYS:CD	2.43	0.47
1:B:13:HIS:HD2	1:B:15:LYS:H	1.61	0.47
1:B:69:LYS:HD2	1:B:209:ARG:HE	1.79	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:3693:PO4:O4	3:B:2102:HOH:O	2.19	0.47
1:B:162:THR:HG22	1:B:185:LLP:HG2	1.96	0.47
1:B:267:VAL:HG13	1:B:342:ARG:HD3	1.97	0.47
1:B:165:LEU:HD22	1:B:187:LEU:HD23	1.97	0.46
1:A:85:LYS:NZ	1:A:313:GLN:OE1	2.37	0.46
1:A:69:LYS:HE3	1:A:97:ALA:HA	1.97	0.46
1:A:119:LYS:HB3	1:A:122:TYR:CE1	2.48	0.46
1:B:88:GLU:HG3	1:B:88:GLU:O	2.15	0.46
1:A:170:ALA:O	1:A:174:MET:HE2	2.15	0.45
1:A:29:THR:HG23	1:A:32:ALA:H	1.81	0.45
1:B:222:GLN:HG2	1:B:226:GLU:O	2.16	0.45
1:A:11:ARG:HG2	3:A:2169:HOH:O	2.16	0.45
1:B:27:HIS:H	1:B:27:HIS:HD1	1.63	0.45
1:B:11:ARG:HG3	1:B:11:ARG:NH1	2.31	0.45
1:B:145:ARG:HD2	1:B:145:ARG:C	2.38	0.44
1:B:4:LEU:HD22	1:B:10:VAL:HA	2.00	0.44
1:A:206:LEU:O	1:A:208:PRO:HD3	2.17	0.44
1:A:128:VAL:HA	1:A:159:ASP:HB3	1.98	0.44
1:A:338:ALA:O	1:A:342:ARG:HG2	2.18	0.44
1:B:35:VAL:O	1:B:38:LYS:HD3	2.17	0.44
1:B:162:THR:CG2	1:B:185:LLP:HG2	2.48	0.43
1:B:40:ARG:HG3	1:B:40:ARG:NH1	2.21	0.43
1:B:40:ARG:HH11	1:B:40:ARG:CG	2.24	0.43
1:A:140:LEU:HD23	1:A:174:MET:HE3	2.00	0.43
1:A:85:LYS:O	1:A:88:GLU:HB3	2.19	0.43
1:B:11:ARG:NH1	3:B:2184:HOH:O	2.50	0.43
1:A:161:VAL:HG22	1:A:181:SER:HA	1.99	0.43
1:B:27:HIS:NE2	1:B:28:ARG:CZ	2.81	0.42
1:B:48:ARG:HH11	1:B:48:ARG:HG3	1.84	0.42
1:B:171:LEU:HA	1:B:171:LEU:HD12	1.88	0.42
1:B:13:HIS:HD2	1:B:15:LYS:HB2	1.85	0.42
1:A:342:ARG:HH11	1:A:342:ARG:HG2	1.84	0.42
1:A:342:ARG:HH11	1:A:342:ARG:CG	2.32	0.42
1:B:13:HIS:CD2	1:B:15:LYS:H	2.37	0.42
1:A:89:ARG:CG	1:A:89:ARG:NH1	2.40	0.42
1:A:88:GLU:O	1:A:91:TYR:HB3	2.20	0.42
1:B:217:ARG:NH1	1:B:217:ARG:CG	2.81	0.42
1:B:297:ARG:HH11	1:B:297:ARG:HG3	1.85	0.42
1:B:69:LYS:HE3	1:B:97:ALA:HA	2.02	0.42
1:A:257:LYS:HA	1:A:257:LYS:HD3	1.83	0.42
1:B:301:ALA:O	1:B:305:ARG:HG2	2.19	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:LYS:NZ	3:A:2186:HOH:O	2.53	0.41
1:A:297:ARG:HD2	1:A:348:ILE:HG22	2.01	0.41
1:A:293:VAL:HA	1:A:294:PRO:HD3	1.78	0.41
1:B:298:VAL:HG22	1:B:348:ILE:HD13	2.02	0.41
1:B:129:HIS:CG	1:B:140:LEU:HD22	2.56	0.41
1:B:259:TRP:CH2	1:B:331:ARG:HD3	2.56	0.41
1:A:257:LYS:HD3	1:A:260:GLN:OE1	2.21	0.40
1:B:293:VAL:HA	1:B:294:PRO:HD3	1.75	0.40
1:B:67:LEU:C	1:B:67:LEU:HD23	2.42	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/352 (98%)	339 (98%)	6 (2%)	0	100	100
1	B	345/352 (98%)	338 (98%)	7 (2%)	0	100	100
All	All	690/704 (98%)	677 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	264/267 (99%)	237 (90%)	27 (10%)	9	8
1	B	264/267 (99%)	242 (92%)	22 (8%)	14	13
All	All	528/534 (99%)	479 (91%)	49 (9%)	11	10

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	ARG
1	A	38	LYS
1	A	44	ARG
1	A	85	LYS
1	A	88	GLU
1	A	89	ARG
1	A	90	PHE
1	A	103	ARG
1	A	129	HIS
1	A	145	ARG
1	A	165	LEU
1	A	201	ARG
1	A	209	ARG
1	A	213	LEU
1	A	217	ARG
1	A	220	LYS
1	A	223	LYS
1	A	247	LEU
1	A	274	ARG
1	A	279	ARG
1	A	297	ARG
1	A	299	LYS
1	A	313	GLN
1	A	317	LYS
1	A	331	ARG
1	A	342	ARG
1	A	345	LEU
1	B	11	ARG
1	B	28	ARG
1	B	38	LYS
1	B	44	ARG
1	B	83	TYR
1	B	85	LYS
1	B	90	PHE
1	B	103	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	145	ARG
1	B	201	ARG
1	B	209	ARG
1	B	217	ARG
1	B	220	LYS
1	B	224	GLU
1	B	237	LEU
1	B	245	GLU
1	B	299	LYS
1	B	313	GLN
1	B	317	LYS
1	B	331	ARG
1	B	342	ARG
1	B	345	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	HIS
1	B	13	HIS
1	B	24	GLN
1	B	26	HIS
1	B	253	HIS
1	B	313	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	LLP	A	185	1	23,24,25	1.71	4 (17%)	28,32,34	1.93	6 (21%)
1	LLP	B	185	1	23,24,25	1.80	4 (17%)	28,32,34	2.15	8 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	185	1	-	0/15/17/19	0/1/1/1
1	LLP	B	185	1	-	0/15/17/19	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	185	LLP	C3-C2	-2.99	1.38	1.40
1	A	185	LLP	C3-C2	-2.41	1.39	1.40
1	A	185	LLP	P-OP3	-2.16	1.46	1.54
1	B	185	LLP	P-OP3	-2.13	1.47	1.54
1	A	185	LLP	C4'-NZ	3.34	1.37	1.27
1	B	185	LLP	C4'-NZ	3.84	1.39	1.27
1	A	185	LLP	C4-C4'	4.91	1.55	1.46
1	B	185	LLP	C4-C4'	4.96	1.55	1.46

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	185	LLP	C3-C4-C4'	-2.80	116.54	120.16
1	B	185	LLP	OP4-P-OP1	-2.41	101.01	107.14
1	A	185	LLP	OP4-P-OP1	-2.19	101.57	107.14
1	A	185	LLP	C5'-C5-C4	2.14	125.06	121.47
1	B	185	LLP	C5'-C5-C4	2.70	126.01	121.47
1	B	185	LLP	C5-C4-C4'	2.86	125.63	121.52
1	A	185	LLP	CE-NZ-C4'	3.21	128.22	118.97
1	B	185	LLP	CE-NZ-C4'	3.45	128.92	118.97
1	A	185	LLP	OP4-C5'-C5	3.63	115.00	108.99
1	B	185	LLP	C2'-C2-C3	3.81	125.64	121.04
1	A	185	LLP	C2'-C2-C3	4.12	126.00	121.04
1	B	185	LLP	OP4-C5'-C5	4.30	116.10	108.99
1	A	185	LLP	CD-CE-NZ	5.08	119.30	110.98
1	B	185	LLP	CD-CE-NZ	5.61	120.16	110.98

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	185	LLP	3	0
1	B	185	LLP	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	B	3693	1	4,4,4	0.64	0	6,6,6	0.28	0
2	PO4	B	3694	-	4,4,4	0.64	0	6,6,6	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	B	3693	1	-	0/0/0/0	0/0/0/0
2	PO4	B	3694	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	3693	PO4	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	347/352 (98%)	-0.29	5 (1%) 78 77	21, 33, 52, 67	0
1	B	347/352 (98%)	-0.23	3 (0%) 85 85	20, 33, 53, 70	0
All	All	694/704 (98%)	-0.26	8 (1%) 81 80	20, 33, 52, 70	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	292	GLY	2.8
1	B	291	GLU	2.7
1	A	349	LEU	2.5
1	B	297	ARG	2.4
1	A	249	ARG	2.2
1	A	291	GLU	2.2
1	A	348	ILE	2.2
1	A	274	ARG	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	LLP	B	185	24/25	0.98	0.14	-	20,23,25,27	0
1	LLP	A	185	24/25	0.98	0.16	-	18,24,28,30	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PO4	B	3694	5/5	0.86	0.21	2.45	78,79,80,80	0
2	PO4	B	3693	5/5	0.80	0.17	0.33	87,88,89,89	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.