



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:36 PM GMT

PDB ID : 4IV3
Title : Crystal structure of recombinant foot-and-mouth-disease virus A22-H2093C empty capsid
Authors : Porta, C.; Kotecha, A.; Burman, A.; Jackson, T.; Ren, J.; Loureiro, S.; Jones, I.M.; Fry, E.E.; Stuart, D.I.; Charleston, B.
Deposited on : 2013-01-22
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	211	<div> <div>2%</div> <div>74%</div> <div>10%</div> <div>16%</div> </div>
2	B	218	<div> <div>3%</div> <div>83%</div> <div>12%</div> <div>5%</div> </div>
3	C	221	<div> <div>%</div> <div>86%</div> <div>13%</div> <div>•</div> </div>
4	D	85	<div> <div>5%</div> <div>93%</div> <div>•</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4848 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Capsid protein VP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	178	Total	C	N	O	S	0	0	0
			1398	887	253	254	4			

- Molecule 2 is a protein called Capsid protein VP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	207	Total	C	N	O	S	0	0	0
			1645	1050	284	305	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	93	CYS	HIS	ENGINEERED MUTATION	UNP Q6PN23

- Molecule 3 is a protein called Capsid protein VP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	221	Total	C	N	O	S	0	0	0
			1707	1087	277	334	9			

- Molecule 4 is a protein called Capsid protein VP4.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	6	Total	C	N	O	0	0	0
			42	29	6	7			

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	19	Total	O	0	0
			19	19		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	16	Total	O	0	0
			16	16		
5	C	21	Total	O	0	0
			21	21		

- Molecule 1: Capsid protein VP1



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	328.02Å 341.49Å 363.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.56 – 2.90 49.56 – 2.90	Depositor EDS
% Data completeness (in resolution range)	79.3 (49.56-2.90) 79.3 (49.56-2.90)	Depositor EDS
R_{merge}	0.41	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.26 (at 2.91Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.235 , 0.239 0.234 , 0.238	Depositor DCC
R_{free} test set	17689 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	36.2	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 49.3	EDS
Estimated twinning fraction	0.014 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	5 of 352479 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	4848	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/1430	0.66	0/1950
2	B	0.35	0/1692	0.64	0/2305
3	C	0.34	0/1755	0.64	0/2400
4	D	0.61	0/42	0.51	0/54
All	All	0.34	0/4919	0.64	0/6709

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1398	0	1401	14	0
2	B	1645	0	1606	12	1
3	C	1707	0	1635	21	1
4	D	42	0	43	1	0
5	A	19	0	0	0	0
5	B	16	0	0	0	0
5	C	21	0	0	0	0
All	All	4848	0	4685	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:214:LEU:HD23	3:C:127:PRO:HG2	1.69	0.74
1:A:40:VAL:HG21	1:A:62:VAL:HB	1.71	0.71
3:C:76:LYS:HE3	3:C:133:PRO:HG2	1.75	0.68
3:C:68:THR:HG22	3:C:70:GLU:HG3	1.74	0.68
2:B:81:LEU:HD11	2:B:85:THR:HG21	1.76	0.66
3:C:66:THR:HG21	3:C:189:GLN:OE1	2.00	0.62
1:A:195:VAL:HG22	1:A:201:HIS:HB2	1.83	0.61
3:C:76:LYS:HE2	3:C:184:TRP:CD1	2.37	0.59
3:C:71:GLN:O	3:C:72:ARG:HG3	2.03	0.58
3:C:79:VAL:HG22	3:C:185:VAL:HG23	1.86	0.57
3:C:181:VAL:HG12	3:C:182:GLN:NE2	2.19	0.56
2:B:116:PHE:HB3	2:B:190:ASN:HD22	1.70	0.56
1:A:101:THR:HG23	3:C:16:THR:HG21	1.90	0.53
1:A:198:GLN:O	1:A:198:GLN:HG3	2.10	0.52
3:C:213:ARG:HG3	3:C:214:LEU:HG	1.93	0.51
2:B:38:THR:O	2:B:38:THR:HG22	2.12	0.50
3:C:66:THR:HG21	3:C:189:GLN:NE2	2.28	0.49
3:C:66:THR:HG21	3:C:189:GLN:CD	2.33	0.48
3:C:130:VAL:CG1	3:C:184:TRP:HZ3	2.25	0.48
2:B:18:ARG:HG3	2:B:23:THR:HG22	1.97	0.47
1:A:108:LEU:HD13	1:A:108:LEU:C	2.35	0.47
1:A:108:LEU:HD13	1:A:109:LYS:N	2.30	0.47
2:B:12:ASP:OD1	2:B:13:ARG:HG3	2.15	0.47
2:B:100:TYR:HB2	2:B:211:ALA:HB3	1.97	0.46
4:D:84:LEU:O	4:D:84:LEU:HG	2.15	0.45
1:A:37:ASP:OD1	1:A:180:LYS:HE2	2.16	0.45
2:B:68:ASP:O	2:B:73:LYS:HE2	2.16	0.45
2:B:17:THR:OG1	2:B:157:HIS:HE1	2.00	0.45
3:C:130:VAL:HG13	3:C:184:TRP:HZ3	1.82	0.45
2:B:114:ASN:HD21	2:B:193:SER:HA	1.81	0.44
1:A:71:TYR:HB2	1:A:185:TYR:HB2	2.00	0.44
3:C:65:VAL:HG13	3:C:199:THR:HG22	2.00	0.44
3:C:34:ARG:HH11	3:C:34:ARG:HG3	1.83	0.43
2:B:43:VAL:HG11	2:B:209:HIS:CE1	2.53	0.43
2:B:80:LYS:HD3	2:B:129:TRP:CZ2	2.54	0.43
3:C:133:PRO:HA	3:C:134:PRO:HD3	1.88	0.43
1:A:101:THR:HG22	1:A:101:THR:O	2.18	0.43
1:A:206:ILE:O	1:A:206:ILE:HG13	2.20	0.42
3:C:90:TYR:CE2	3:C:94:ILE:HD11	2.55	0.42
3:C:120:ARG:NH1	3:C:149:ASP:OD1	2.53	0.41
1:A:25:GLN:HG2	1:A:27:ARG:HG2	2.01	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:72:ARG:HB3	3:C:72:ARG:NH1	2.36	0.41
1:A:38:ARG:HD2	1:A:61:LEU:HD23	2.03	0.41
1:A:25:GLN:OE1	1:A:25:GLN:N	2.53	0.40
1:A:112:PHE:O	3:C:10:GLY:HA2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:27:GLN:OE1	3:C:151:GLY:N[2_555]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	174/211 (82%)	166 (95%)	8 (5%)	0	100	100
2	B	205/218 (94%)	191 (93%)	14 (7%)	0	100	100
3	C	219/221 (99%)	201 (92%)	17 (8%)	1 (0%)	34	71
4	D	4/85 (5%)	3 (75%)	1 (25%)	0	100	100
All	All	602/735 (82%)	561 (93%)	40 (7%)	1 (0%)	52	84

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	58	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	150/172 (87%)	145 (97%)	5 (3%)	45	80
2	B	182/193 (94%)	179 (98%)	3 (2%)	70	91
3	C	185/185 (100%)	182 (98%)	3 (2%)	70	91
4	D	3/67 (4%)	2 (67%)	1 (33%)	0	1
All	All	520/617 (84%)	508 (98%)	12 (2%)	58	87

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	ARG
1	A	58	GLN
1	A	89	VAL
1	A	181	ARG
1	A	199	ASP
2	B	40	GLU
2	B	64	LYS
2	B	131	GLU
3	C	58	ASP
3	C	65	VAL
3	C	179	THR
4	D	80	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	GLN
1	A	198	GLN
2	B	103	ASN
2	B	157	HIS
2	B	190	ASN
3	C	71	GLN
3	C	221	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	178/211 (84%)	-0.66	5 (2%) 56 50	15, 25, 88, 153	0
2	B	207/218 (94%)	-0.64	7 (3%) 49 41	12, 25, 79, 159	0
3	C	221/221 (100%)	-0.70	2 (0%) 85 84	14, 26, 77, 111	0
4	D	6/85 (7%)	3.00	4 (66%) 0 0	96, 118, 135, 154	0
All	All	612/735 (83%)	-0.63	18 (2%) 55 49	12, 26, 88, 159	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	39	GLN	6.1
4	D	80	PHE	5.4
2	B	40	GLU	5.2
1	A	198	GLN	5.2
4	D	83	LEU	3.8
2	B	218	GLU	3.5
1	A	210	LYS	3.3
2	B	41	ASP	3.2
4	D	85	ALA	3.1
2	B	38	THR	2.7
1	A	156	ALA	2.7
2	B	12	ASP	2.5
4	D	84	LEU	2.4
3	C	67	ARG	2.4
1	A	209	ALA	2.4
3	C	178	THR	2.3
2	B	13	ARG	2.1
1	A	134	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.