



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:50 AM GMT

PDB ID : 2IWG
Title : COMPLEX BETWEEN THE PRYSPRY DOMAIN OF TRIM21 AND IGG
FC
Authors : James, L.C.; Keeble, A.H.; Rhodes, D.A.; Trowsdale, J.
Deposited on : 2006-06-30
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1352 (2.38-2.34)
Clashscore	102246	1456 (2.38-2.34)
Ramachandran outliers	100387	1435 (2.38-2.34)
Sidechain outliers	100360	1436 (2.38-2.34)
RSRZ outliers	91569	1358 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	<div> <div>2%</div> <div>65%</div> <div>29%</div> <div>6%</div> </div>
1	D	207	<div> <div>%</div> <div>62%</div> <div>33%</div> <div>5%</div> </div>
2	B	181	<div> <div>58%</div> <div>35%</div> <div>6%</div> </div>
2	E	181	<div> <div>%</div> <div>64%</div> <div>31%</div> <div>5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MAN	A	1446	X	-	-	-
3	MAN	D	1446	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6590 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IG GAMMA-1 CHAIN C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	0	0	0
			1658	1056	279	317	6			
1	D	207	Total	C	N	O	S	0	0	0
			1658	1056	279	317	6			


- Molecule 2 is a protein called 52 KDA RO PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	181	Total	C	N	O	S	0	0	0
			1455	936	248	264	7			
2	E	181	Total	C	N	O	S	0	0	0
			1455	936	248	264	7			

- Molecule 3 is a polymer of unknown type called SUGAR (7-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	7	Total	C	N	O	0	0
			86	48	3	35		
3	D	7	Total	C	N	O	0	0
			86	48	3	35		

- Molecule 4 is SUGAR (2,6-ANHYDRO-1-DEOXY-D-GALACTITOL) (three-letter code: FU4) (formula: C₆H₁₂O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		
4	D	1	Total	C	O	0	0
			10	6	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	48	Total	O	0	0
			48	48		
5	B	31	Total	O	0	0
			31	31		
5	D	42	Total	O	0	0
			42	42		
5	E	51	Total	O	0	0
			51	51		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: IG GAMMA-1 CHAIN C



• Molecule 1: IG GAMMA-1 CHAIN C




• Molecule 2: 52 KDA RO PROTEIN



• Molecule 2: 52 KDA RO PROTEIN





4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	112.49Å 112.49Å 194.61Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	97.59 – 2.35 97.42 – 2.35	Depositor EDS
% Data completeness (in resolution range)	86.1 (97.59-2.35) 86.1 (97.42-2.35)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 2.34Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.215 , 0.253 0.213 , 0.248	Depositor DCC
R_{free} test set	2533 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	52.5	Xtriage
Anisotropy	0.502	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
Estimated twinning fraction	0.477 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 49869 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6590	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FU4, GAL, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.63	16/1704 (0.9%)	1.16	7/2322 (0.3%)
1	D	1.65	22/1704 (1.3%)	1.20	10/2322 (0.4%)
2	B	1.74	28/1505 (1.9%)	1.44	17/2050 (0.8%)
2	E	1.69	23/1505 (1.5%)	1.47	17/2050 (0.8%)
All	All	1.68	89/6418 (1.4%)	1.31	51/8744 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	D	0	1
3	A	1	0
3	D	1	0
All	All	2	2

The worst 5 of 89 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	135	VAL	CB-CG1	-9.28	1.33	1.52
2	B	135	VAL	CB-CG1	-8.98	1.33	1.52
2	E	163	PHE	CG-CD2	8.63	1.51	1.38
1	D	407	TYR	CG-CD1	8.62	1.50	1.39
1	A	296	TYR	CE2-CZ	8.24	1.49	1.38

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	41	ARG	NE-CZ-NH2	-12.66	113.97	120.30
2	B	41	ARG	NE-CZ-NH1	12.62	126.61	120.30
2	E	64	ASP	CB-CG-OD2	12.30	129.37	118.30
2	B	41	ARG	NE-CZ-NH2	-12.27	114.17	120.30
2	E	41	ARG	NE-CZ-NH1	11.73	126.16	120.30

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	1446	MAN	C1
3	D	1446	MAN	C1

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	264	VAL	Peptide
1	D	264	VAL	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1658	0	1625	38	0
1	D	1658	0	1625	39	0
2	B	1455	0	1381	37	0
2	E	1455	0	1381	36	0
3	A	86	0	73	7	0
3	D	86	0	73	7	0
4	A	10	0	12	3	0
4	D	10	0	12	4	0
5	A	48	0	0	2	0
5	B	31	0	0	3	0
5	D	42	0	0	2	0
5	E	51	0	0	3	0
All	All	6590	0	6182	160	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 160 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1445:NAG:HN2	4:A:1451:FU4:H1C1	1.11	1.08
3:D:1445:NAG:HN2	4:D:1451:FU4:H1C1	1.12	1.07
1:A:265:ASP:HB3	1:A:299:THR:HB	1.30	1.06
1:D:265:ASP:HB3	1:D:299:THR:HB	1.39	1.00
3:D:1444:NAG:O6	4:D:1451:FU4:H2	1.71	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/207 (99%)	192 (94%)	11 (5%)	2 (1%)	19	20
1	D	205/207 (99%)	194 (95%)	9 (4%)	2 (1%)	19	20
2	B	179/181 (99%)	163 (91%)	13 (7%)	3 (2%)	11	9
2	E	179/181 (99%)	164 (92%)	12 (7%)	3 (2%)	11	9
All	All	768/776 (99%)	713 (93%)	45 (6%)	10 (1%)	15	13

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	102	LYS
2	B	143	HIS
2	E	102	LYS
2	E	143	HIS
2	B	154	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/193 (100%)	182 (94%)	11 (6%)	25	30
1	D	193/193 (100%)	185 (96%)	8 (4%)	37	48
2	B	158/158 (100%)	145 (92%)	13 (8%)	14	15
2	E	158/158 (100%)	148 (94%)	10 (6%)	22	25
All	All	702/702 (100%)	660 (94%)	42 (6%)	24	28

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	115	LEU
2	B	172	LYS
2	E	115	LEU
2	B	123	GLN
2	B	151	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	38	ASN
1	D	390	ASN
2	E	25	GLN
2	B	33	GLN
1	D	438	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

14 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAG	A	1444	1,3	14,14,15	1.28	2 (14%)	15,19,21	2.02	3 (20%)
3	NAG	A	1445	3	14,14,15	0.87	0	15,19,21	1.58	4 (26%)
3	MAN	A	1446	3	11,11,12	0.88	1 (9%)	14,15,17	2.09	5 (35%)
3	MAN	A	1447	3	11,11,12	0.81	0	14,15,17	2.43	6 (42%)
3	NAG	A	1448	3	14,14,15	0.71	0	15,19,21	1.62	1 (6%)
3	MAN	A	1449	3	11,11,12	1.78	4 (36%)	14,15,17	2.39	6 (42%)
3	GAL	A	1450	3	11,11,12	0.84	1 (9%)	14,15,17	2.06	4 (28%)
3	NAG	D	1444	1,3	14,14,15	1.06	1 (7%)	15,19,21	2.03	4 (26%)
3	NAG	D	1445	3	14,14,15	0.79	0	15,19,21	1.49	3 (20%)
3	MAN	D	1446	3	11,11,12	0.91	1 (9%)	14,15,17	1.87	5 (35%)
3	MAN	D	1447	3	11,11,12	0.86	0	14,15,17	2.24	4 (28%)
3	NAG	D	1448	3	14,14,15	0.67	0	15,19,21	1.73	1 (6%)
3	MAN	D	1449	3	11,11,12	1.45	2 (18%)	14,15,17	2.52	5 (35%)
3	GAL	D	1450	3	11,11,12	1.06	0	14,15,17	2.29	6 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1444	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1445	3	-	0/6/23/26	0/1/1/1
3	MAN	A	1446	3	1/1/4/5	0/2/19/22	0/1/1/1
3	MAN	A	1447	3	-	0/2/19/22	0/1/1/1
3	NAG	A	1448	3	-	0/6/23/26	0/1/1/1
3	MAN	A	1449	3	-	0/2/19/22	0/1/1/1
3	GAL	A	1450	3	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	D	1444	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	1445	3	-	0/6/23/26	0/1/1/1
3	MAN	D	1446	3	1/1/4/5	0/2/19/22	0/1/1/1
3	MAN	D	1447	3	-	0/2/19/22	0/1/1/1
3	NAG	D	1448	3	-	0/6/23/26	0/1/1/1
3	MAN	D	1449	3	-	0/2/19/22	0/1/1/1
3	GAL	D	1450	3	-	0/2/19/22	0/1/1/1

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	1444	NAG	C1-C2	2.01	1.55	1.52
3	D	1446	MAN	O5-C1	2.01	1.47	1.43
3	A	1449	MAN	C1-C2	2.07	1.57	1.52
3	A	1444	NAG	C8-C7	2.19	1.54	1.50
3	A	1446	MAN	O5-C1	2.20	1.47	1.43

The worst 5 of 57 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1449	MAN	C1-C2-C3	-5.27	103.30	109.54
3	A	1449	MAN	C1-C2-C3	-4.89	103.76	109.54
3	A	1446	MAN	O3-C3-C4	-4.63	99.91	110.34
3	A	1444	NAG	O4-C4-C5	-4.36	97.69	109.24
3	D	1444	NAG	O4-C4-C5	-4.32	97.79	109.24

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	D	1446	MAN	C1
3	A	1446	MAN	C1

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1444	NAG	1	0
3	A	1445	NAG	5	0
3	A	1446	MAN	1	0
3	A	1447	MAN	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	1444	NAG	2	0
3	D	1445	NAG	4	0
3	D	1446	MAN	1	0
3	D	1447	MAN	1	0

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	FU4	A	1451	-	10,10,10	1.66	3 (30%)	14,14,14	2.95	4 (28%)
4	FU4	D	1451	-	10,10,10	1.42	2 (20%)	14,14,14	2.88	6 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FU4	A	1451	-	-	0/0/17/17	0/1/1/1
4	FU4	D	1451	-	-	0/0/17/17	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	1451	FU4	C6-C5	2.11	1.56	1.51
4	D	1451	FU4	C2-C3	2.39	1.55	1.52
4	A	1451	FU4	C6-C5	2.40	1.57	1.51
4	A	1451	FU4	C1-C2	2.58	1.58	1.52
4	A	1451	FU4	C2-C3	2.79	1.56	1.52

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	D	1451	FU4	O3-C3-C2	-2.47	105.53	110.00
4	D	1451	FU4	O2-C2-C3	-2.21	105.67	110.12
4	A	1451	FU4	O3-C3-C2	-2.17	106.08	110.00
4	D	1451	FU4	C6-C5-C4	2.04	117.10	113.08
4	D	1451	FU4	O4-C4-C5	2.06	114.68	109.84

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1451	FU4	3	0
4	D	1451	FU4	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	207/207 (100%)	0.30	5 (2%) 62 74	48, 78, 90, 90	0
1	D	207/207 (100%)	0.38	3 (1%) 78 87	48, 78, 90, 90	0
2	B	181/181 (100%)	0.29	0 100 100	48, 64, 82, 89	0
2	E	181/181 (100%)	0.26	1 (0%) 90 95	49, 64, 82, 89	0
All	All	776/776 (100%)	0.31	9 (1%) 81 89	48, 70, 90, 90	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	296	TYR	3.7
1	D	241	PHE	2.9
1	A	243	PHE	2.4
1	A	298	SER	2.3
1	D	237	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GAL	D	1450	11/12	0.94	0.22	1.89	83,85,90,90	0
3	GAL	A	1450	11/12	0.98	0.13	-0.98	83,85,90,90	0
3	NAG	A	1448	14/15	0.96	0.14	-	76,87,90,90	0
3	MAN	A	1447	11/12	0.93	0.13	-	89,90,90,90	0
3	MAN	A	1449	11/12	0.85	0.13	-	89,90,90,90	0
3	MAN	D	1446	11/12	0.95	0.07	-	88,89,90,90	0
3	MAN	D	1449	11/12	0.84	0.18	-	89,90,90,90	0
3	MAN	A	1446	11/12	0.88	0.10	-	88,89,90,90	0
3	NAG	A	1445	14/15	0.96	0.10	-	88,89,90,90	0
3	NAG	A	1444	14/15	0.90	0.12	-	87,90,90,90	0
3	NAG	D	1448	14/15	0.97	0.18	-	76,88,90,90	0
3	MAN	D	1447	11/12	0.92	0.17	-	89,90,90,90	0
3	NAG	D	1445	14/15	0.94	0.13	-	88,89,90,90	0
3	NAG	D	1444	14/15	0.87	0.14	-	87,90,90,90	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	FU4	D	1451	10/10	0.91	0.16	-	89,90,90,90	0
4	FU4	A	1451	10/10	0.92	0.16	-	89,90,90,90	0

6.5 Other polymers [i](#)

There are no such residues in this entry.