



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 09:36 AM GMT

PDB ID : 3IWK
Title : Crystal structure of aminoaldehyde dehydrogenase 1 from Pisum sativum (PsAMADH1)
Authors : Kopecny, D.; Morera, S.; Briozzo, P.
Deposited on : 2009-09-02
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	503	<div> <div></div> <div>83%13% ..</div> </div>
1	B	503	<div> <div></div> <div>79%16% ..</div> </div>
1	C	503	<div> <div></div> <div>81%15% ..</div> </div>
1	D	503	<div> <div>2%</div> <div>83%14% ..</div> </div>
1	E	503	<div> <div></div> <div>81%16% ..</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	503	
1	G	503	
1	H	503	
1	I	503	
1	J	503	
1	K	503	
1	L	503	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NA	A	504	-	-	-	X
2	NA	C	504	-	-	-	X
2	NA	E	504	-	-	-	X
2	NA	I	504	-	-	-	X
3	NAD	B	505	-	-	-	X
3	NAD	C	505	-	-	-	X
3	NAD	D	505	-	-	-	X
3	NAD	E	505	-	-	-	X
3	NAD	F	505	-	-	-	X
3	NAD	H	505	-	-	-	X
3	NAD	I	505	-	-	-	X
3	NAD	K	505	-	-	-	X
4	GOL	B	2001	-	-	-	X
4	GOL	B	2002	-	-	-	X
4	GOL	C	2003	-	-	-	X
4	GOL	F	2006	-	-	-	X
4	GOL	G	2007	-	-	-	X
4	GOL	G	2009	-	-	-	X
4	GOL	G	2011	-	-	-	X
4	GOL	H	2010	-	-	-	X
4	GOL	I	2012	-	-	-	X
4	GOL	J	2014	-	-	-	X
4	GOL	K	2016	-	-	-	X
4	GOL	K	2018	-	-	-	X
4	GOL	L	2015	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	L	2017	-	-	-	X
5	MRD	D	2004	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 48369 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminoaldehyde dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	496	Total	C	N	O	S	0	1	0
			3811	2430	639	726	16			
1	B	496	Total	C	N	O	S	0	1	0
			3813	2432	641	724	16			
1	C	495	Total	C	N	O	S	0	0	0
			3797	2421	637	723	16			
1	D	496	Total	C	N	O	S	0	0	0
			3806	2427	639	724	16			
1	E	496	Total	C	N	O	S	0	0	0
			3806	2427	639	724	16			
1	F	495	Total	C	N	O	S	0	1	0
			3804	2426	639	723	16			
1	G	496	Total	C	N	O	S	0	0	0
			3806	2427	639	724	16			
1	H	497	Total	C	N	O	S	0	1	0
			3820	2435	643	726	16			
1	I	496	Total	C	N	O	S	0	2	0
			3821	2437	644	724	16			
1	J	497	Total	C	N	O	S	0	0	0
			3812	2430	640	726	16			
1	K	496	Total	C	N	O	S	0	2	0
			3821	2437	644	724	16			
1	L	496	Total	C	N	O	S	0	1	0
			3810	2429	640	725	16			

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Na	0	0
			1	1		
2	J	1	Total	Na	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total	Na	0	0
			1	1		
2	K	1	Total	Na	0	0
			1	1		
2	E	1	Total	Na	0	0
			1	1		
2	H	1	Total	Na	0	0
			1	1		
2	B	1	Total	Na	0	0
			1	1		
2	I	1	Total	Na	0	0
			1	1		
2	C	1	Total	Na	0	0
			1	1		
2	A	1	Total	Na	0	0
			1	1		
2	L	1	Total	Na	0	0
			1	1		
2	F	1	Total	Na	0	0
			1	1		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O P	0	0
			44	21	7	14 2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	E	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	F	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	G	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	H	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	I	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	J	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	K	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	L	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		
4	F	1	Total	C	O	0	0
			6	3	3		
4	G	1	Total	C	O	0	0
			6	3	3		
4	G	1	Total	C	O	0	0
			6	3	3		
4	G	1	Total	C	O	0	0
			6	3	3		
4	H	1	Total	C	O	0	0
			6	3	3		
4	I	1	Total	C	O	0	0
			6	3	3		
4	J	1	Total	C	O	0	0
			6	3	3		
4	J	1	Total	C	O	0	0
			6	3	3		
4	K	1	Total	C	O	0	0
			6	3	3		
4	K	1	Total	C	O	0	0
			6	3	3		
4	K	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is (4R)-2-METHYLPENTANE-2,4-DIOL (three-letter code: MRD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			8	6	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	145	Total	O	0	0
			145	145		
6	B	134	Total	O	0	0
			134	134		
6	C	148	Total	O	0	0
			148	148		
6	D	117	Total	O	0	0
			117	117		
6	E	154	Total	O	0	0
			154	154		
6	F	160	Total	O	0	0
			160	160		
6	G	180	Total	O	0	0
			180	180		
6	H	197	Total	O	0	0
			197	197		
6	I	199	Total	O	0	0
			199	199		
6	J	170	Total	O	0	0
			170	170		
6	K	195	Total	O	0	0
			195	195		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	L	198	Total 198	O 198	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Aminoaldehyde dehydrogenase




• Molecule 1: Aminoaldehyde dehydrogenase




• Molecule 1: Aminoaldehyde dehydrogenase







• Molecule 1: Aminoaldehyde dehydrogenase




• Molecule 1: Aminoaldehyde dehydrogenase




• Molecule 1: Aminoaldehyde dehydrogenase




• Molecule 1: Aminoaldehyde dehydrogenase






• Molecule 1: Aminoaldehyde dehydrogenase

Chain H: 82% 15% ..




• Molecule 1: Aminoaldehyde dehydrogenase

Chain I: 82% 14% ..




• Molecule 1: Aminoaldehyde dehydrogenase


Chain J: 84% 12% ..



- Molecule 1: Aminoaldehyde dehydrogenase



- Molecule 1: Aminoaldehyde dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	86.41Å 216.87Å 205.76Å 90.00° 98.02° 90.00°	Depositor
Resolution (Å)	47.84 – 2.40 46.81 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.84-2.40) 99.8 (46.81-2.40)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.61 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.200 , 0.244 0.200 , 0.244	Depositor DCC
R_{free} test set	14543 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	33.0	Xtriage
Anisotropy	0.056	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 290857 reflections (0.000%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	48369	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MRD, GOL, NAD, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/3894	0.53	0/5286
1	B	0.38	0/3897	0.53	0/5290
1	C	0.37	0/3877	0.54	0/5264
1	D	0.36	0/3886	0.54	0/5275
1	E	0.39	0/3886	0.56	1/5275 (0.0%)
1	F	0.39	0/3888	0.53	0/5279
1	G	0.38	0/3886	0.54	1/5275 (0.0%)
1	H	0.38	0/3903	0.56	1/5297 (0.0%)
1	I	0.38	0/3908	0.55	0/5304
1	J	0.38	0/3892	0.53	0/5283
1	K	0.38	0/3908	0.54	0/5304
1	L	0.39	0/3894	0.55	0/5287
All	All	0.38	0/46719	0.54	3/63419 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	73	LEU	CA-CB-CG	6.92	131.21	115.30
1	E	95	LEU	CA-CB-CG	6.09	129.30	115.30
1	G	302	VAL	CB-CA-C	-5.15	101.61	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3811	0	3841	44	0
1	B	3813	0	3844	59	0
1	C	3797	0	3824	50	0
1	D	3806	0	3837	44	0
1	E	3806	0	3837	48	0
1	F	3804	0	3831	34	0
1	G	3806	0	3837	39	0
1	H	3820	0	3855	41	0
1	I	3821	0	3857	38	0
1	J	3812	0	3842	35	0
1	K	3821	0	3857	36	0
1	L	3810	0	3836	40	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
2	I	1	0	0	0	0
2	J	1	0	0	0	0
2	K	1	0	0	0	0
2	L	1	0	0	0	0
3	A	44	0	26	5	0
3	B	44	0	26	2	0
3	C	44	0	26	0	0
3	D	27	0	12	0	0
3	E	44	0	26	1	0
3	F	44	0	26	3	0
3	G	44	0	26	2	0
3	H	44	0	26	2	0
3	I	44	0	26	2	0
3	J	44	0	26	1	0
3	K	44	0	26	2	0
3	L	44	0	26	2	0
4	B	12	0	16	2	0
4	C	6	0	8	0	0
4	F	6	0	8	0	0
4	G	18	0	24	0	0
4	H	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	I	6	0	8	0	0
4	J	12	0	16	1	0
4	K	18	0	24	2	0
4	L	30	0	40	3	0
5	D	8	0	14	2	0
6	A	145	0	0	1	0
6	B	134	0	0	1	0
6	C	148	0	0	2	0
6	D	117	0	0	1	0
6	E	154	0	0	0	0
6	F	160	0	0	3	0
6	G	180	0	0	3	0
6	H	197	0	0	1	0
6	I	199	0	0	0	0
6	J	170	0	0	0	0
6	K	195	0	0	3	0
6	L	198	0	0	0	0
All	All	48369	0	46562	490	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 490 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:323:ILE:HD11	1:J:374:PHE:HD2	1.19	1.07
1:I:456:GLN:HB3	1:J:136:ILE:HD11	1.39	1.05
1:E:24:ARG:HH11	1:E:24:ARG:HG3	1.29	0.97
1:E:323:ILE:HD11	1:E:374:PHE:HD2	1.34	0.92
1:J:323:ILE:HD11	1:J:374:PHE:CD2	2.03	0.92

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	495/503 (98%)	472 (95%)	20 (4%)	3 (1%)	30	43
1	B	495/503 (98%)	473 (96%)	20 (4%)	2 (0%)	39	56
1	C	493/503 (98%)	471 (96%)	18 (4%)	4 (1%)	24	35
1	D	494/503 (98%)	472 (96%)	20 (4%)	2 (0%)	39	56
1	E	494/503 (98%)	476 (96%)	17 (3%)	1 (0%)	52	69
1	F	494/503 (98%)	477 (97%)	16 (3%)	1 (0%)	52	69
1	G	494/503 (98%)	473 (96%)	17 (3%)	4 (1%)	24	35
1	H	496/503 (99%)	477 (96%)	15 (3%)	4 (1%)	24	35
1	I	496/503 (99%)	478 (96%)	15 (3%)	3 (1%)	30	43
1	J	495/503 (98%)	474 (96%)	19 (4%)	2 (0%)	39	56
1	K	496/503 (99%)	475 (96%)	19 (4%)	2 (0%)	39	56
1	L	495/503 (98%)	480 (97%)	14 (3%)	1 (0%)	52	69
All	All	5937/6036 (98%)	5698 (96%)	210 (4%)	29 (0%)	34	48

5 of 29 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	451	GLN
1	B	451	GLN
1	C	60	ARG
1	C	61	LYS
1	C	65	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	411/416 (99%)	378 (92%)	33 (8%)	15	23
1	B	411/416 (99%)	377 (92%)	34 (8%)	14	21

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	409/416 (98%)	372 (91%)	37 (9%)	12	17
1	D	410/416 (99%)	382 (93%)	28 (7%)	20	31
1	E	410/416 (99%)	380 (93%)	30 (7%)	17	27
1	F	410/416 (99%)	385 (94%)	25 (6%)	23	36
1	G	410/416 (99%)	378 (92%)	32 (8%)	16	24
1	H	412/416 (99%)	379 (92%)	33 (8%)	15	23
1	I	412/416 (99%)	375 (91%)	37 (9%)	12	17
1	J	411/416 (99%)	381 (93%)	30 (7%)	17	27
1	K	412/416 (99%)	385 (93%)	27 (7%)	21	32
1	L	411/416 (99%)	381 (93%)	30 (7%)	17	27
All	All	4929/4992 (99%)	4553 (92%)	376 (8%)	16	25

5 of 376 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	140	MET
1	G	379	ILE
1	L	9	GLN
1	F	217	LEU
1	G	84	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	212	ASN
1	G	368	HIS
1	L	29	ASN
1	F	292	GLN
1	G	9	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 44 ligands modelled in this entry, 12 are monoatomic - leaving 32 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAD	A	505	-	38,48,48	2.58	7 (18%)	47,73,73	3.63	6 (12%)
4	GOL	B	2001	-	5,5,5	0.35	0	5,5,5	0.32	0
4	GOL	B	2002	-	5,5,5	0.35	0	5,5,5	0.29	0
3	NAD	B	505	-	38,48,48	2.60	8 (21%)	47,73,73	3.56	6 (12%)
4	GOL	C	2003	-	5,5,5	0.37	0	5,5,5	0.25	0
3	NAD	C	505	-	38,48,48	2.56	7 (18%)	47,73,73	3.58	5 (10%)
5	MRD	D	2004	-	6,7,7	0.29	0	7,10,10	0.42	0
3	NAD	D	505	-	23,29,48	3.31	7 (30%)	27,45,73	4.30	2 (7%)
3	NAD	E	505	-	38,48,48	2.59	7 (18%)	47,73,73	3.73	8 (17%)
4	GOL	F	2006	-	5,5,5	0.39	0	5,5,5	0.13	0
3	NAD	F	505	-	38,48,48	2.54	7 (18%)	47,73,73	3.60	7 (14%)
4	GOL	G	2007	-	5,5,5	0.34	0	5,5,5	0.24	0
4	GOL	G	2009	-	5,5,5	0.36	0	5,5,5	0.49	0
4	GOL	G	2011	-	5,5,5	0.37	0	5,5,5	0.28	0
3	NAD	G	505	-	38,48,48	2.61	8 (21%)	47,73,73	3.54	6 (12%)
4	GOL	H	2010	-	5,5,5	0.37	0	5,5,5	0.50	0
3	NAD	H	505	-	38,48,48	2.61	7 (18%)	47,73,73	3.62	7 (14%)
4	GOL	I	2012	-	5,5,5	0.30	0	5,5,5	0.50	0
3	NAD	I	505	-	38,48,48	2.61	7 (18%)	47,73,73	3.61	6 (12%)
4	GOL	J	2013	-	5,5,5	0.36	0	5,5,5	0.25	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	J	2014	-	5,5,5	0.34	0	5,5,5	0.38	0
3	NAD	J	505	-	38,48,48	2.58	7 (18%)	47,73,73	3.63	6 (12%)
4	GOL	K	2016	-	5,5,5	0.34	0	5,5,5	0.38	0
4	GOL	K	2018	-	5,5,5	0.33	0	5,5,5	0.23	0
4	GOL	K	2019	-	5,5,5	0.33	0	5,5,5	0.27	0
3	NAD	K	505	-	38,48,48	2.59	7 (18%)	47,73,73	3.54	6 (12%)
4	GOL	L	2000	-	5,5,5	0.35	0	5,5,5	0.20	0
4	GOL	L	2005	-	5,5,5	0.38	0	5,5,5	0.31	0
4	GOL	L	2008	-	5,5,5	0.39	0	5,5,5	0.17	0
4	GOL	L	2015	-	5,5,5	0.35	0	5,5,5	0.16	0
4	GOL	L	2017	-	5,5,5	0.32	0	5,5,5	0.27	0
3	NAD	L	505	-	38,48,48	2.62	8 (21%)	47,73,73	3.53	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	505	-	-	0/22/62/62	0/5/5/5
4	GOL	B	2001	-	-	0/4/4/4	0/0/0/0
4	GOL	B	2002	-	-	0/4/4/4	0/0/0/0
3	NAD	B	505	-	-	0/22/62/62	0/5/5/5
4	GOL	C	2003	-	-	0/4/4/4	0/0/0/0
3	NAD	C	505	-	-	0/22/62/62	0/5/5/5
5	MRD	D	2004	-	-	0/5/5/5	0/0/0/0
3	NAD	D	505	-	-	0/12/32/62	0/3/3/5
3	NAD	E	505	-	-	0/22/62/62	0/5/5/5
4	GOL	F	2006	-	-	0/4/4/4	0/0/0/0
3	NAD	F	505	-	-	0/22/62/62	0/5/5/5
4	GOL	G	2007	-	-	0/4/4/4	0/0/0/0
4	GOL	G	2009	-	-	0/4/4/4	0/0/0/0
4	GOL	G	2011	-	-	0/4/4/4	0/0/0/0
3	NAD	G	505	-	-	0/22/62/62	0/5/5/5
4	GOL	H	2010	-	-	0/4/4/4	0/0/0/0
3	NAD	H	505	-	-	0/22/62/62	0/5/5/5
4	GOL	I	2012	-	-	0/4/4/4	0/0/0/0
3	NAD	I	505	-	-	0/22/62/62	0/5/5/5
4	GOL	J	2013	-	-	0/4/4/4	0/0/0/0
4	GOL	J	2014	-	-	0/4/4/4	0/0/0/0
3	NAD	J	505	-	-	0/22/62/62	0/5/5/5
4	GOL	K	2016	-	-	0/4/4/4	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	K	2018	-	-	0/4/4/4	0/0/0/0
4	GOL	K	2019	-	-	0/4/4/4	0/0/0/0
3	NAD	K	505	-	-	0/22/62/62	0/5/5/5
4	GOL	L	2000	-	-	0/4/4/4	0/0/0/0
4	GOL	L	2005	-	-	0/4/4/4	0/0/0/0
4	GOL	L	2008	-	-	0/4/4/4	0/0/0/0
4	GOL	L	2015	-	-	0/4/4/4	0/0/0/0
4	GOL	L	2017	-	-	0/4/4/4	0/0/0/0
3	NAD	L	505	-	-	0/22/62/62	0/5/5/5

The worst 5 of 87 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	L	505	NAD	PN-O2N	2.03	1.63	1.54
3	A	505	NAD	C6N-N1N	2.05	1.41	1.35
3	D	505	NAD	PN-O5D	2.05	1.63	1.55
3	J	505	NAD	C6N-N1N	2.05	1.41	1.35
3	B	505	NAD	C3N-C7N	2.08	1.53	1.50

The worst 5 of 72 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	505	NAD	N3A-C2A-N1A	-22.30	111.82	128.89
3	C	505	NAD	N3A-C2A-N1A	-22.26	111.85	128.89
3	E	505	NAD	N3A-C2A-N1A	-22.14	111.94	128.89
3	A	505	NAD	N3A-C2A-N1A	-22.11	111.96	128.89
3	J	505	NAD	N3A-C2A-N1A	-22.01	112.04	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

17 monomers are involved in 32 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	505	NAD	5	0
4	B	2002	GOL	2	0
3	B	505	NAD	2	0
5	D	2004	MRD	2	0
3	E	505	NAD	1	0
3	F	505	NAD	3	0
3	G	505	NAD	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	H	505	NAD	2	0
3	I	505	NAD	2	0
4	J	2014	GOL	1	0
3	J	505	NAD	1	0
4	K	2016	GOL	1	0
4	K	2018	GOL	1	0
3	K	505	NAD	2	0
4	L	2005	GOL	2	0
4	L	2008	GOL	1	0
3	L	505	NAD	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	496/503 (98%)	-0.11	7 (1%) 78 77	24, 34, 50, 65	0
1	B	496/503 (98%)	-0.23	4 (0%) 87 87	21, 34, 50, 60	0
1	C	495/503 (98%)	-0.22	5 (1%) 84 83	20, 32, 51, 65	0
1	D	496/503 (98%)	0.02	11 (2%) 65 64	22, 40, 65, 71	0
1	E	496/503 (98%)	-0.21	5 (1%) 84 83	22, 33, 47, 56	0
1	F	495/503 (98%)	-0.34	3 (0%) 90 90	21, 32, 44, 52	0
1	G	496/503 (98%)	-0.17	6 (1%) 81 81	21, 31, 45, 55	0
1	H	497/503 (98%)	-0.28	2 (0%) 93 93	20, 31, 42, 51	0
1	I	496/503 (98%)	-0.31	4 (0%) 87 87	19, 31, 44, 51	0
1	J	497/503 (98%)	-0.27	6 (1%) 81 81	21, 33, 50, 58	0
1	K	496/503 (98%)	-0.25	5 (1%) 84 83	19, 32, 45, 51	0
1	L	496/503 (98%)	-0.38	1 (0%) 95 95	19, 30, 45, 50	0
All	All	5952/6036 (98%)	-0.23	59 (0%) 84 83	19, 32, 49, 71	0

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	63	GLY	5.0
1	H	6	SER	5.0
1	C	501	SER	5.0
1	K	502	LYS	4.5
1	A	502	LYS	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	GOL	K	2016	6/6	0.85	0.30	8.74	45,46,46,46	0
5	MRD	D	2004	8/8	0.80	0.37	8.20	54,55,55,55	0
4	GOL	I	2012	6/6	0.87	0.27	7.55	47,47,47,47	0
2	NA	C	504	1/1	0.64	0.34	6.81	65,65,65,65	0
4	GOL	H	2010	6/6	0.84	0.38	5.87	59,60,60,60	0
4	GOL	F	2006	6/6	0.89	0.27	5.65	50,50,51,51	0
4	GOL	J	2014	6/6	0.87	0.27	5.48	54,55,55,56	0
2	NA	I	504	1/1	0.54	0.23	4.91	61,61,61,61	0
4	GOL	B	2002	6/6	0.89	0.27	4.72	60,61,61,61	0
2	NA	E	504	1/1	0.89	0.24	4.60	47,47,47,47	0
4	GOL	L	2017	6/6	0.77	0.24	4.54	63,64,64,64	0
4	GOL	C	2003	6/6	0.80	0.27	4.52	50,52,52,53	0
4	GOL	G	2011	6/6	0.79	0.31	4.35	65,65,66,66	0
4	GOL	G	2009	6/6	0.85	0.27	3.97	43,46,46,47	0
2	NA	A	504	1/1	0.88	0.20	3.94	60,60,60,60	0
4	GOL	K	2018	6/6	0.84	0.36	3.68	48,49,49,50	0
4	GOL	G	2007	6/6	0.80	0.17	3.28	61,62,62,62	0
3	NAD	K	505	44/44	0.78	0.22	3.02	60,73,76,77	0
3	NAD	E	505	44/44	0.73	0.24	2.65	58,75,78,78	0
3	NAD	C	505	44/44	0.78	0.22	2.64	61,74,79,79	0
3	NAD	F	505	44/44	0.77	0.23	2.60	61,76,79,79	0
3	NAD	I	505	44/44	0.79	0.23	2.52	67,81,86,86	0
3	NAD	D	505	27/44	0.78	0.23	2.42	67,73,79,79	0
4	GOL	L	2015	6/6	0.52	0.19	2.39	61,62,62,62	0
3	NAD	H	505	44/44	0.80	0.24	2.31	46,69,76,77	0
3	NAD	B	505	44/44	0.78	0.21	2.15	58,75,79,79	0
4	GOL	B	2001	6/6	0.81	0.18	2.04	58,59,59,60	0
4	GOL	L	2008	6/6	0.89	0.18	1.74	46,46,47,47	0
3	NAD	J	505	44/44	0.81	0.20	1.69	65,77,79,79	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAD	A	505	44/44	0.78	0.22	1.66	53,71,73,73	0
3	NAD	L	505	44/44	0.81	0.18	1.41	47,66,68,68	0
3	NAD	G	505	44/44	0.81	0.20	1.04	57,69,73,74	0
2	NA	L	504	1/1	0.75	0.15	0.54	61,61,61,61	0
4	GOL	L	2005	6/6	0.92	0.21	0.26	66,66,67,67	0
2	NA	G	504	1/1	0.95	0.14	0.21	43,43,43,43	0
2	NA	J	504	1/1	0.96	0.09	-0.80	41,41,41,41	0
2	NA	D	504	1/1	0.87	0.09	-0.92	43,43,43,43	0
2	NA	H	504	1/1	0.97	0.09	-1.31	30,30,30,30	0
2	NA	K	504	1/1	0.89	0.07	-1.74	37,37,37,37	0
2	NA	B	504	1/1	0.96	0.06	-2.13	41,41,41,41	0
2	NA	F	504	1/1	0.97	0.05	-2.26	34,34,34,34	0
4	GOL	J	2013	6/6	0.86	0.18	-	58,58,59,59	0
4	GOL	L	2000	6/6	0.72	0.26	-	66,66,66,67	0
4	GOL	K	2019	6/6	0.81	0.20	-	56,56,57,57	0

6.5 Other polymers

There are no such residues in this entry.