



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 29, 2016 – 02:26 PM EDT

PDB ID : 5IWP  
Title : Structure of Transient Receptor Potential (TRP) channel TRPV6 in the presence of calcium  
Authors : Saotome, K.; Singh, A.K.; Yelshanskaya, M.V.; Sobolevsky, A.I.  
Deposited on : 2016-03-22  
Resolution : 3.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027790  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027790


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1010 (3.82-3.50)
Clashscore	102246	1125 (3.82-3.50)
Ramachandran outliers	100387	1079 (3.82-3.50)
Sidechain outliers	100360	1078 (3.82-3.50)
RSRZ outliers	91569	1017 (3.82-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	672	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4735 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transient receptor potential cation channel subfamily V member 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	591	Total	C	N	O	S	0	0	0
			4717	3049	792	844	32			


There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	62	TYR	ILE	engineered mutation	UNP Q9R186
A	92	ASN	LEU	engineered mutation	UNP Q9R186
A	96	GLN	MET	engineered mutation	UNP Q9R186
A	495	GLN	LEU	engineered mutation	UNP Q9R186
A	670	VAL	-	expression tag	UNP Q9R186
A	671	PRO	-	expression tag	UNP Q9R186
A	672	ARG	-	expression tag	UNP Q9R186

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Ca	0	0
			3	3		

- Molecule 3 is 6-(5-METHYL-2-OXO-IMIDAZOLIDIN-4-YL)-HEXANOIC ACID (three-letter code: DTB) (formula: C<sub>10</sub>H<sub>18</sub>N<sub>2</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			15	10	2	3		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transient receptor potential cation channel subfamily V member 6



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	144.35Å 144.35Å 113.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.56 – 3.65 49.56 – 3.65	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.56-3.65) 99.9 (49.56-3.65)	Depositor EDS
$R_{merge}$	0.01	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.88 (at 3.67Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.273 , 0.283 0.299 , 0.289	Depositor DCC
$R_{free}$ test set	690 reflections (4.99%)	DCC
Wilson B-factor (Å <sup>2</sup> )	142.3	Xtrriage
Anisotropy	0.374	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 110.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.83	EDS
Total number of atoms	4735	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	138.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DTB, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.28	1/4824 (0.0%)	0.57	12/6553 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	372	LEU	CA-C	-5.13	1.39	1.52

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	68	GLU	N-CA-C	-8.86	87.09	111.00
1	A	359	THR	N-CA-CB	8.40	126.26	110.30
1	A	67	PHE	N-CA-C	-7.78	90.01	111.00
1	A	359	THR	N-CA-C	-6.99	92.12	111.00
1	A	67	PHE	CB-CA-C	6.85	124.11	110.40
1	A	375	ALA	N-CA-C	-6.79	92.65	111.00
1	A	442	ARG	NE-CZ-NH2	-6.71	116.95	120.30
1	A	68	GLU	N-CA-CB	6.65	122.57	110.60
1	A	378	THR	N-CA-CB	-6.63	97.70	110.30
1	A	378	THR	N-CA-C	6.54	128.66	111.00
1	A	358	ARG	CB-CA-C	-5.22	99.96	110.40
1	A	584	VAL	CB-CA-C	-5.08	101.76	111.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	351	LEU	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4717	0	4734	102	0
2	A	3	0	0	0	0
3	A	15	0	17	4	0
All	All	4735	0	4751	104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:376:TYR:O	1:A:377:VAL:HG22	1.41	1.19
1:A:375:ALA:HA	1:A:376:TYR:O	1.68	0.92
1:A:376:TYR:O	1:A:377:VAL:CG2	2.25	0.81
1:A:315:LEU:O	1:A:315:LEU:HD12	1.81	0.80
1:A:578:GLY:HA2	1:A:581:HIS:HD2	1.49	0.78
1:A:114:LEU:O	1:A:151:VAL:HG22	1.93	0.68
1:A:351:LEU:HB3	1:A:352:LYS:HA	1.75	0.67
1:A:578:GLY:HA2	1:A:581:HIS:CD2	2.30	0.65
1:A:115:TYR:CE1	3:A:704:DTB:HCR	2.32	0.65
1:A:375:ALA:HA	1:A:376:TYR:C	2.18	0.64
1:A:420:LEU:H	1:A:421:GLY:HA3	1.60	0.64
1:A:366:LEU:HD12	1:A:366:LEU:H	1.63	0.63
1:A:183:ILE:HD13	1:A:187:ALA:HB3	1.81	0.63
1:A:519:LEU:HD11	1:A:543:PRO:HB3	1.83	0.61
1:A:354:ARG:NE	1:A:367:LEU:O	2.34	0.60
1:A:373:GLN:HB2	1:A:375:ALA:HB3	1.83	0.60
1:A:365:THR:HG22	1:A:367:LEU:H	1.67	0.59

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:378:THR:O	1:A:378:THR:HG23	2.04	0.58
1:A:599:THR:HG23	1:A:602:MET:HE2	1.86	0.58
1:A:543:PRO:O	1:A:554:TYR:OH	2.15	0.57
1:A:505:SER:HG	1:A:525:TYR:HH	1.53	0.57
1:A:540:ILE:HG22	1:A:542:GLY:H	1.71	0.56
1:A:190:ARG:NH2	1:A:229:LYS:O	2.41	0.54
3:A:704:DTB:HCD1	3:A:704:DTB:HCT2	1.91	0.53
1:A:243:PHE:HA	1:A:258:LEU:HD13	1.90	0.52
1:A:367:LEU:HB3	1:A:369:GLN:HG3	1.91	0.52
1:A:115:TYR:CZ	3:A:704:DTB:HCR	2.45	0.52
1:A:343:THR:HG21	1:A:461:TRP:HE1	1.74	0.52
1:A:590:GLU:O	1:A:594:ALA:N	2.32	0.51
1:A:364:ASN:ND2	1:A:549:ASP:OD1	2.37	0.51
1:A:437:VAL:O	1:A:441:MET:N	2.39	0.51
1:A:282:ILE:HG23	1:A:291:LEU:HD23	1.93	0.51
1:A:506:ALA:O	1:A:510:ILE:HG12	2.10	0.50
1:A:584:VAL:HG22	1:A:584:VAL:O	2.10	0.50
1:A:284:SER:HB2	1:A:285:SER:HB2	1.94	0.50
1:A:351:LEU:CB	1:A:352:LYS:HA	2.40	0.49
1:A:591:LEU:O	1:A:595:GLN:HG3	2.12	0.49
1:A:72:VAL:HG22	1:A:105:LEU:HD11	1.94	0.49
1:A:190:ARG:HD2	1:A:231:LEU:HD13	1.93	0.49
1:A:443:LEU:O	1:A:443:LEU:HD12	2.12	0.49
1:A:230:SER:HB3	1:A:233:LEU:HD23	1.94	0.49
1:A:315:LEU:C	1:A:315:LEU:HD12	2.32	0.49
1:A:630:LEU:HD12	1:A:631:ARG:N	2.28	0.49
1:A:304:ARG:O	1:A:307:LEU:HG	2.13	0.49
1:A:442:ARG:O	1:A:443:LEU:HB3	2.13	0.48
1:A:425:HIS:NE2	1:A:602:MET:SD	2.74	0.48
1:A:40:GLN:O	1:A:44:ILE:HG12	2.14	0.48
1:A:73:HIS:HE2	1:A:104:GLU:HG3	1.80	0.47
1:A:316:VAL:HA	1:A:319:LYS:HD3	1.96	0.47
1:A:337:LEU:HA	1:A:340:ILE:HG22	1.97	0.47
1:A:451:VAL:HG13	1:A:553:MET:SD	2.54	0.47
1:A:28:SER:OG	1:A:29:TRP:N	2.48	0.46
1:A:441:MET:CE	1:A:452:PRO:HG3	2.45	0.46
1:A:461:TRP:O	1:A:464:VAL:HB	2.15	0.46
1:A:343:THR:HG22	1:A:457:LEU:HD22	1.98	0.46
1:A:372:LEU:HB2	1:A:373:GLN:HA	1.97	0.46
1:A:115:TYR:CE1	1:A:151:VAL:HG21	2.51	0.46
1:A:236:ASN:OD1	1:A:237:ASN:N	2.47	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:PHE:HA	1:A:68:GLU:HA	1.77	0.46
1:A:88:LEU:HD23	1:A:89:TYR:CE1	2.51	0.46
1:A:505:SER:O	1:A:509:ILE:HG12	2.16	0.46
1:A:315:LEU:CD1	1:A:319:LYS:HD2	2.47	0.45
1:A:51:LEU:O	1:A:51:LEU:HD12	2.17	0.45
1:A:220:LEU:HD21	1:A:232:GLU:HG3	2.00	0.44
1:A:73:HIS:NE2	1:A:104:GLU:HG3	2.32	0.44
1:A:352:LYS:HB2	1:A:353:PRO:HD2	1.98	0.44
1:A:373:GLN:HB2	1:A:375:ALA:N	2.33	0.44
1:A:625:GLY:HA2	1:A:626:ASP:HA	1.63	0.44
1:A:441:MET:HE2	1:A:452:PRO:HG3	2.00	0.43
1:A:291:LEU:O	1:A:291:LEU:HD12	2.17	0.43
1:A:315:LEU:HD11	1:A:319:LYS:HD2	2.00	0.43
1:A:350:PRO:HG2	1:A:442:ARG:HH12	1.82	0.43
1:A:483:LYS:O	1:A:487:GLY:N	2.50	0.43
1:A:523:TYR:CE1	1:A:527:MET:HG2	2.54	0.43
1:A:27:GLU:HA	1:A:28:SER:HA	1.73	0.43
1:A:534:GLU:HB3	1:A:539:ILE:CG2	2.49	0.43
1:A:614:ARG:HD3	1:A:628:TRP:CE2	2.54	0.43
1:A:427:ILE:HG23	1:A:459:LEU:HG	2.00	0.42
1:A:621:GLU:N	1:A:622:TYR:O	2.52	0.42
1:A:64:LEU:O	1:A:68:GLU:O	2.36	0.42
1:A:252:ILE:HD13	1:A:306:ILE:HD13	2.01	0.42
1:A:244:LYS:HB2	1:A:244:LYS:HE3	1.74	0.42
1:A:459:LEU:O	1:A:463:ASN:HB2	2.20	0.42
1:A:601:VAL:O	1:A:605:ARG:HG3	2.19	0.42
1:A:49:LEU:O	1:A:49:LEU:HD12	2.20	0.42
1:A:539:ILE:HG23	1:A:540:ILE:N	2.33	0.42
1:A:465:MET:SD	1:A:495:GLN:HG2	2.59	0.42
1:A:315:LEU:O	1:A:319:LYS:HG3	2.19	0.42
1:A:40:GLN:HA	1:A:77:ALA:HB3	2.01	0.42
3:A:704:DTB:HCT2	3:A:704:DTB:CD	2.50	0.41
1:A:462:CYS:O	1:A:465:MET:HG2	2.20	0.41
1:A:179:VAL:O	1:A:183:ILE:HG12	2.20	0.41
1:A:366:LEU:HD12	1:A:366:LEU:N	2.31	0.41
1:A:396:VAL:O	1:A:400:LEU:N	2.51	0.41
1:A:328:PHE:CZ	1:A:467:PHE:HA	2.55	0.41
1:A:543:PRO:HB2	1:A:554:TYR:CZ	2.55	0.41
1:A:189:ILE:HD11	1:A:223:ASP:HA	2.02	0.41
1:A:260:GLN:HA	1:A:263:LYS:HE3	2.03	0.41
1:A:366:LEU:HD13	1:A:514:GLU:HB3	2.03	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:511:PHE:HA	1:A:514:GLU:HG2	2.02	0.41
1:A:110:MET:HB2	1:A:115:TYR:O	2.21	0.40
1:A:373:GLN:HA	1:A:374:GLU:HB2	2.03	0.40
1:A:535:LEU:CD2	1:A:542:GLY:HA2	2.51	0.40
1:A:441:MET:HG2	1:A:449:GLU:HA	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	585/672 (87%)	537 (92%)	44 (8%)	4 (1%)	26 72

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	376	TYR
1	A	377	VAL
1	A	379	PRO
1	A	583	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	508/584 (87%)	489 (96%)	19 (4%)	41 77

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	LEU
1	A	78	MET
1	A	139	ARG
1	A	279	LEU
1	A	349	ARG
1	A	354	ARG
1	A	359	THR
1	A	366	LEU
1	A	371	LEU
1	A	376	TYR
1	A	420	LEU
1	A	442	ARG
1	A	443	LEU
1	A	514	GLU
1	A	519	LEU
1	A	541	ASP
1	A	622	TYR
1	A	630	LEU
1	A	631	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	DTB	A	704	-	12,15,15	3.32	4 (33%)	14,19,19	2.91	6 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DTB	A	704	-	-	0/6/20/20	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	704	DTB	CS-N1	-3.43	1.43	1.46
3	A	704	DTB	CR-N2	-2.21	1.42	1.45
3	A	704	DTB	CN-N1	7.01	1.45	1.35
3	A	704	DTB	CN-N2	7.95	1.47	1.35

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	704	DTB	CR-N2-CN	-5.57	108.25	112.49
3	A	704	DTB	CS-N1-CN	-4.54	108.33	112.63
3	A	704	DTB	CT-CS-N1	-3.06	108.42	111.56
3	A	704	DTB	CD-CE-CR	-2.06	109.68	114.03
3	A	704	DTB	CS-CR-N2	4.70	107.66	102.30
3	A	704	DTB	CR-CS-N1	4.99	107.69	102.43

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	704	DTB	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	591/672 (87%)	0.21	43 (7%) 18 10	80, 131, 187, 213	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	360	ASN	5.1
1	A	578	GLY	5.1
1	A	607	LEU	4.8
1	A	358	ARG	4.5
1	A	374	GLU	4.4
1	A	355	ILE	4.4
1	A	361	PRO	4.2
1	A	609	ARG	4.2
1	A	577	MET	4.0
1	A	606	LYS	4.0
1	A	610	CYS	3.9
1	A	608	PRO	3.8
1	A	615	SER	3.7
1	A	284	SER	3.6
1	A	575	ALA	3.5
1	A	377	VAL	3.5
1	A	249	GLU	3.4
1	A	605	ARG	3.4
1	A	614	ARG	3.2
1	A	571	ASN	3.1
1	A	603	LEU	3.1
1	A	612	TRP	3.0
1	A	359	THR	3.0
1	A	574	ILE	2.8
1	A	68	GLU	2.8
1	A	382	ASP	2.8
1	A	611	LEU	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	141	ALA	2.7
1	A	283	ASP	2.5
1	A	154	TYR	2.5
1	A	613	PRO	2.5
1	A	570	LEU	2.4
1	A	62	TYR	2.4
1	A	383	LEU	2.4
1	A	376	TYR	2.3
1	A	372	LEU	2.3
1	A	380	LYS	2.3
1	A	143	VAL	2.3
1	A	616	GLY	2.2
1	A	327	TYR	2.2
1	A	393	VAL	2.1
1	A	604	GLU	2.1
1	A	357	ASN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	DTB	A	704	15/15	0.81	0.38	0.82	77,77,77,77	0
2	CA	A	702	1/1	0.78	0.32	-	25,25,25,25	1
2	CA	A	701	1/1	0.91	0.23	-	25,25,25,25	1
2	CA	A	703	1/1	0.97	0.32	-	25,25,25,25	1



## 6.5 Other polymers [i](#)

There are no such residues in this entry.