



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:49 AM GMT

PDB ID : 2IXE
Title : CRYSTAL STRUCTURE OF THE ATPASE DOMAIN OF TAP1 WITH ATP
(D645N MUTANT)
Authors : Procko, E.; Ferrin-O'Connell, I.; Ng, S.-L.; Gaudet, R.
Deposited on : 2006-07-07
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	
1	D	271	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4203 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ANTIGEN PEPTIDE TRANSPORTER 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	S	0	2	0
			1902	1203	336	356	7			
1	D	248	Total	C	N	O	S	0	5	0
			1900	1205	337	351	7			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	464	MET	-	EXPRESSION TAG	UNP P36370
A	726	ALA	-	EXPRESSION TAG	UNP P36370
A	727	ALA	-	EXPRESSION TAG	UNP P36370
A	728	ALA	-	EXPRESSION TAG	UNP P36370
A	729	HIS	-	EXPRESSION TAG	UNP P36370
A	730	HIS	-	EXPRESSION TAG	UNP P36370
A	731	HIS	-	EXPRESSION TAG	UNP P36370
A	732	HIS	-	EXPRESSION TAG	UNP P36370
A	733	HIS	-	EXPRESSION TAG	UNP P36370
A	734	HIS	-	EXPRESSION TAG	UNP P36370
A	645	ASN	ASP	ENGINEERED MUTATION	UNP P36370
D	464	MET	-	EXPRESSION TAG	UNP P36370
D	726	ALA	-	EXPRESSION TAG	UNP P36370
D	727	ALA	-	EXPRESSION TAG	UNP P36370
D	728	ALA	-	EXPRESSION TAG	UNP P36370
D	729	HIS	-	EXPRESSION TAG	UNP P36370
D	730	HIS	-	EXPRESSION TAG	UNP P36370
D	731	HIS	-	EXPRESSION TAG	UNP P36370
D	732	HIS	-	EXPRESSION TAG	UNP P36370
D	733	HIS	-	EXPRESSION TAG	UNP P36370
D	734	HIS	-	EXPRESSION TAG	UNP P36370
D	645	ASN	ASP	ENGINEERED MUTATION	UNP P36370

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	159	Total	O	0	0
			159	159		
5	D	173	Total	O	0	0
			173	173		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ANTIGEN PEPTIDE TRANSPORTER 1



• Molecule 1: ANTIGEN PEPTIDE TRANSPORTER 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	80.43 Å 80.43 Å 168.12 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.55 – 2.00 37.25 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (72.55-2.00) 99.7 (37.25-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 2.00 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.199 , 0.241 0.204 , 0.249	Depositor DCC
R_{free} test set	2705 reflections (7.66%)	DCC
Wilson B-factor (Å ²)	32.8	Xtriage
Anisotropy	0.230	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 38006 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4203	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PO4, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/1946	0.54	0/2648
1	D	0.35	0/1951	0.54	0/2648
All	All	0.35	0/3897	0.54	0/5296

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1902	0	1875	10	0
1	D	1900	0	1906	7	0
2	A	31	0	12	0	0
2	D	31	0	12	0	0
3	A	1	0	0	0	0
3	D	1	0	0	0	0
4	A	5	0	0	0	0
5	A	159	0	0	2	0
5	D	173	0	0	1	0
All	All	4203	0	3805	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:478:GLY:O	1:D:672:THR:HG21	1.86	0.76
1:A:570:ARG:HD3	1:A:574:GLU:OE1	1.91	0.71
1:A:618[A]:ASN:HA	1:A:625[A]:ARG:HH21	1.56	0.69
1:A:529:GLN:HB2	1:A:531:LEU:HD13	1.84	0.60
1:D:680:LEU:HD21	1:D:716:MET:HG2	1.86	0.57
1:A:478:GLY:O	1:A:672:THR:HG21	2.07	0.54
1:A:679:GLN:OE1	5:A:2141:HOH:O	2.20	0.48
1:A:567:LEU:O	1:A:616:THR:HA	2.13	0.48
1:A:632:ARG:HD2	5:A:2109:HOH:O	2.14	0.47
1:D:559:ALA:HB3	1:D:638:PRO:HG3	1.98	0.46
1:A:529:GLN:CB	1:A:531:LEU:HD13	2.47	0.45
1:D:480:VAL:HG11	1:D:528:LEU:HD21	1.98	0.44
1:D:645:ASN:ND2	5:D:2112:HOH:O	2.50	0.44
1:D:533:GLN:NE2	1:D:547:VAL:CG1	2.81	0.43
1:D:590:THR:HG23	1:D:599:HIS:CD2	2.56	0.41
1:A:528:LEU:CD1	1:A:674:LEU:HD22	2.52	0.40
1:A:566:LEU:O	1:A:632:ARG:NH2	2.48	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	251/271 (93%)	248 (99%)	2 (1%)	1 (0%)	39	33
1	D	247/271 (91%)	246 (100%)	1 (0%)	0	100	100
All	All	498/542 (92%)	494 (99%)	3 (1%)	1 (0%)	52	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	615	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/219 (90%)	192 (97%)	6 (3%)	48	47
1	D	201/219 (92%)	190 (94%)	11 (6%)	27	21
All	All	399/438 (91%)	382 (96%)	17 (4%)	37	30

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	509	LYS
1	A	573	ARG
1	A	581	THR
1	A	640	LEU
1	A	672	THR
1	A	680	LEU
1	D	471	LEU
1	D	474	LEU
1	D	501	LEU
1	D	535[A]	THR
1	D	535[B]	THR
1	D	588	GLU
1	D	607	GLN
1	D	639	ARG
1	D	672	THR
1	D	709	ARG
1	D	718	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	557	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	619	GLN
1	D	494	ASN
1	D	533	GLN
1	D	599	HIS
1	D	607	GLN
1	D	619	GLN
1	D	679	GLN
1	D	700	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ATP	A	1	3	24,33,33	0.96	1 (4%)	31,52,52	1.87	5 (16%)
4	PO4	A	3	-	4,4,4	0.46	0	6,6,6	0.27	0
2	ATP	D	1	3	24,33,33	0.97	1 (4%)	31,52,52	1.90	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	A	1	3	-	0/18/38/38	0/3/3/3
4	PO4	A	3	-	-	0/0/0/0	0/0/0/0
2	ATP	D	1	3	-	0/18/38/38	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	ATP	C5-C4	2.87	1.47	1.40
2	A	1	ATP	C5-C4	2.96	1.47	1.40

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	ATP	N3-C2-N1	-7.86	122.88	128.89
2	A	1	ATP	N3-C2-N1	-7.66	123.03	128.89
2	A	1	ATP	C4-C5-N7	-3.27	106.47	109.48
2	D	1	ATP	PA-O3A-PB	-2.83	124.79	132.73
2	D	1	ATP	C4-C5-N7	-2.69	107.00	109.48
2	A	1	ATP	PA-O3A-PB	-2.58	125.48	132.73
2	A	1	ATP	O2B-PB-O3B	2.01	114.23	105.09
2	A	1	ATP	O3G-PG-O2G	2.16	115.61	107.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	251/271 (92%)	1.04	36 (14%)	4 4	38, 45, 52, 58	0
1	D	248/271 (91%)	1.25	52 (20%)	1 1	30, 44, 52, 55	0
All	All	499/542 (92%)	1.14	88 (17%)	2 2	30, 44, 52, 58	0

All (88) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	602	ILE	6.0
1	D	568	PHE	5.9
1	D	613	VAL	5.7
1	D	605	PHE	5.5
1	A	704	LEU	5.2
1	A	669	ALA	5.0
1	D	603	SER	4.8
1	A	718	GLU	4.6
1	A	709	ARG	4.4
1	A	670	SER	4.4
1	A	711	GLY	4.3
1	D	607	GLN	4.2
1	D	721	ALA	4.2
1	A	469	GLY	4.2
1	A	715	SER	4.1
1	D	606	PRO	4.0
1	A	710	GLY	3.9
1	D	676	ILE	3.8
1	D	572	PHE	3.7
1	D	578	TYR	3.6
1	D	586	MET	3.4
1	A	714	ARG	3.4
1	D	583	THR	3.3
1	D	570	ARG	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	474	LEU	3.2
1	D	566	LEU	3.2
1	A	668	TRP	3.0
1	A	706	LEU	3.0
1	D	643	LEU	3.0
1	D	608	GLY	2.9
1	A	618[A]	ASN	2.9
1	D	673	VAL	2.9
1	A	475	ASN	2.9
1	A	717	VAL	2.8
1	A	707	MET	2.8
1	A	660	ARG	2.8
1	A	708	GLU	2.8
1	A	716	MET	2.8
1	D	646	ALA	2.7
1	A	667	GLU	2.6
1	D	674	LEU	2.6
1	D	599	HIS	2.6
1	D	600	ASP	2.6
1	D	620	LEU	2.5
1	D	561	VAL	2.5
1	D	710	GLY	2.5
1	A	583	THR	2.5
1	D	513	LEU	2.5
1	D	585	THR	2.4
1	A	643	LEU	2.4
1	A	680	LEU	2.4
1	D	567	LEU	2.4
1	D	692	LEU	2.4
1	A	641	LEU	2.4
1	D	690	LEU	2.4
1	A	506	TYR	2.4
1	D	468[A]	SER	2.4
1	A	491	ASN	2.4
1	A	551	HIS	2.3
1	D	584	PRO	2.3
1	D	647	THR	2.3
1	D	514	VAL	2.3
1	A	705	GLN	2.3
1	D	571	SER	2.2
1	D	609	TYR	2.2
1	A	674	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	675	LEU	2.2
1	D	511	THR	2.2
1	D	551[A]	HIS	2.2
1	D	619	GLN	2.2
1	D	656	LEU	2.1
1	A	703	HIS	2.1
1	D	505	LEU	2.1
1	A	673	VAL	2.1
1	D	680	LEU	2.1
1	A	698	CYS	2.1
1	D	579	GLY	2.1
1	A	671	ARG	2.1
1	D	610	ASP	2.1
1	D	604	GLY	2.1
1	D	714	ARG	2.1
1	D	558	VAL	2.1
1	A	650	LEU	2.1
1	D	467	LEU	2.1
1	D	582	ARG	2.0
1	D	569	GLY	2.0
1	A	676	ILE	2.0
1	D	524	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ATP	A	1	31/31	0.95	0.17	-0.25	30,43,46,47	0
2	ATP	D	1	31/31	0.98	0.14	-1.00	24,36,39,39	0
4	PO4	A	3	5/5	0.79	0.26	-	96,96,97,97	0
3	MG	A	2	1/1	0.95	0.15	-	32,32,32,32	0
3	MG	D	2	1/1	0.97	0.20	-	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.