



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 08:08 PM GMT

PDB ID : 1IYH  
Title : Crystal structure of hematopoietic prostaglandin D synthase  
Authors : Inoue, T.  
Deposited on : 2002-08-26  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	198	<div> <div>3%</div> <div>85%</div> <div>14%</div> </div>
1	B	198	<div> <div>%</div> <div>88%</div> <div>12%</div> </div>
1	C	198	<div> <div>6%</div> <div>80%</div> <div>19%</div> </div>
1	D	198	<div> <div>5%</div> <div>85%</div> <div>15%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7616 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEMATOPOIETIC PROSTAGLANDIN D SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	198	Total	C	N	O	S	0	0	0
			1638	1058	273	298	9			
1	B	198	Total	C	N	O	S	0	0	0
			1638	1058	273	298	9			
1	C	198	Total	C	N	O	S	0	0	0
			1638	1058	273	298	9			
1	D	198	Total	C	N	O	S	0	0	0
			1638	1058	273	298	9			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	144	MET	ASN	CONFLICT	UNP O60760
B	344	MET	ASN	CONFLICT	UNP O60760
C	544	MET	ASN	CONFLICT	UNP O60760
D	744	MET	ASN	CONFLICT	UNP O60760

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is GLUTATHIONE (three-letter code: GSH) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>3</sub>O<sub>6</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			20	10	3	6	1		
3	B	1	Total	C	N	O	S	0	0
			20	10	3	6	1		
3	C	1	Total	C	N	O	S	0	0
			20	10	3	6	1		
3	D	1	Total	C	N	O	S	0	0
			20	10	3	6	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	279	Total	O	0	0
			279	279		
4	B	268	Total	O	0	0
			268	268		
4	C	234	Total	O	0	0
			234	234		
4	D	201	Total	O	0	0
			201	201		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: HEMATOPOIETIC PROSTAGLANDIN D SYNTHASE




#### • Molecule 1: HEMATOPOIETIC PROSTAGLANDIN D SYNTHASE



#### • Molecule 1: HEMATOPOIETIC PROSTAGLANDIN D SYNTHASE



#### • Molecule 1: HEMATOPOIETIC PROSTAGLANDIN D SYNTHASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.11Å 47.37Å 184.70Å 90.00° 97.44° 90.00°	Depositor
Resolution (Å)	30.94 – 1.70 30.94 – 1.70	Depositor EDS
% Data completeness (in resolution range)	86.4 (30.94-1.70) 86.5 (30.94-1.70)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.65 (at 1.71Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.205 , 0.238 0.206 , 0.238	Depositor DCC
$R_{free}$ test set	8029 reflections (10.02%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.8	Xtriage
Anisotropy	0.170	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 48.6	EDS
Estimated twinning fraction	0.012 for h,-k,-h-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 80142 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7616	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.77 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.4792e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GSH, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.32	0/1681	0.55	0/2283
1	B	0.32	0/1681	0.57	0/2283
1	C	0.31	0/1681	0.53	0/2283
1	D	0.30	0/1681	0.51	0/2283
All	All	0.31	0/6724	0.54	0/9132

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1638	0	1622	28	0
1	B	1638	0	1622	19	0
1	C	1638	0	1622	32	0
1	D	1638	0	1622	19	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	20	0	15	0	0
3	B	20	0	15	0	0
3	C	20	0	15	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	20	0	15	0	0
4	A	279	0	0	15	0
4	B	268	0	0	4	0
4	C	234	0	0	7	0
4	D	201	0	0	3	0
All	All	7616	0	6548	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181:VAL:HG22	4:A:1934:HOH:O	1.75	0.84
1:C:576:THR:HG23	1:D:765:PRO:HG3	1.62	0.80
1:C:593:ARG:HH11	1:C:593:ARG:HG3	1.54	0.71
1:B:319:LEU:HA	1:B:323:ASN:HD22	1.58	0.68
1:B:338:ARG:HD2	4:B:2168:HOH:O	1.93	0.68
1:A:151:PHE:HA	4:A:1957:HOH:O	1.92	0.67
1:C:442:ILE:O	1:C:446:LEU:HD13	1.93	0.67
1:A:151:PHE:HD1	4:A:1957:HOH:O	1.78	0.65
1:C:589:ASN:HA	1:C:592:LYS:NZ	2.15	0.62
1:B:281:THR:O	1:B:285:GLN:HG3	2.01	0.60
1:A:9:PHE:CE1	1:A:51:ILE:HD11	2.37	0.59
1:B:209:PHE:CE1	1:B:251:ILE:HD11	2.40	0.57
1:A:119:LEU:HA	1:A:123:ASN:HD22	1.69	0.57
1:D:703:PRO:HB2	1:D:706:GLU:HB2	1.88	0.56
1:C:402:PRO:HG2	1:C:404:TYR:CZ	2.41	0.56
1:C:524:ALA:HB3	1:C:525:PRO:HD3	1.87	0.56
1:A:116:PHE:O	1:A:120:LEU:HD22	2.06	0.56
1:C:410:ASN:HD22	1:C:433:ARG:HH11	1.53	0.55
1:A:50:LYS:HE2	4:A:2019:HOH:O	2.06	0.55
1:A:181:VAL:CG2	4:A:1957:HOH:O	2.54	0.55
1:A:39:TRP:HB3	1:A:40:PRO:HD3	1.89	0.55
1:C:503:PRO:HB2	1:C:506:GLU:HB2	1.89	0.55
1:C:440:PRO:HG2	4:C:1705:HOH:O	2.07	0.55
1:C:439:TRP:N	1:C:440:PRO:HD2	2.22	0.55
1:C:593:ARG:HG3	1:C:593:ARG:NH1	2.20	0.54
1:C:480:ASN:ND2	4:C:1815:HOH:O	2.40	0.54
1:D:778:ARG:O	1:D:782:GLN:HG3	2.07	0.54
1:D:639:TRP:HB3	1:D:640:PRO:HD3	1.88	0.54

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2060:HOH:O	1:C:462:HIS:HE1	1.92	0.53
1:B:281:THR:OG1	1:B:284:GLU:HG3	2.09	0.52
1:D:659:LEU:HD12	4:D:1978:HOH:O	2.08	0.52
1:A:193:ARG:HD3	4:A:2115:HOH:O	2.09	0.52
1:C:402:PRO:HG2	1:C:404:TYR:CE1	2.45	0.52
1:A:181:VAL:CG1	4:A:1957:HOH:O	2.58	0.52
1:C:589:ASN:HA	1:C:592:LYS:HZ3	1.75	0.51
1:D:651:ILE:HB	1:D:652:PRO:HA	1.93	0.51
1:D:621:ILE:HD13	1:D:672:THR:HG23	1.94	0.50
1:D:714:GLN:HB3	4:D:1958:HOH:O	2.12	0.50
1:D:724:ALA:HB3	1:D:725:PRO:HD3	1.94	0.50
1:A:14:ARG:NH1	4:A:2132:HOH:O	2.42	0.49
1:B:205:LYS:HB2	4:B:2120:HOH:O	2.12	0.49
1:B:262:HIS:HE1	4:D:1869:HOH:O	1.95	0.49
1:B:235:GLU:HA	1:B:235:GLU:OE2	2.13	0.49
1:B:202:PRO:HB3	1:B:257:ASP:OD1	2.12	0.49
1:C:578:ARG:O	1:C:582:GLN:HG3	2.13	0.49
1:A:103:PRO:HB2	1:A:106:GLU:HB2	1.95	0.49
1:A:108:LYS:HB2	1:A:111:VAL:HG12	1.94	0.49
1:A:14:ARG:HD3	4:A:2132:HOH:O	2.13	0.48
1:A:151:PHE:N	4:A:1957:HOH:O	2.45	0.48
1:A:62:HIS:HE1	4:C:1652:HOH:O	1.96	0.48
1:B:341:LEU:HB3	1:B:342:ILE:HD12	1.96	0.48
1:B:319:LEU:HD23	1:B:323:ASN:HD22	1.79	0.47
1:A:62:HIS:O	1:A:63:GLN:HB2	2.13	0.47
1:C:596:GLN:HA	4:C:1730:HOH:O	2.14	0.47
1:B:319:LEU:HD23	1:B:323:ASN:ND2	2.29	0.46
1:A:150:ASP:C	4:A:1957:HOH:O	2.54	0.46
1:A:116:PHE:O	1:A:120:LEU:CD2	2.64	0.46
1:A:151:PHE:CA	4:A:1957:HOH:O	2.57	0.46
1:C:451:ILE:HB	1:C:452:PRO:HA	1.97	0.46
1:B:314:GLN:O	1:B:318:GLU:HG3	2.16	0.45
1:C:462:HIS:O	1:C:463:GLN:HB2	2.16	0.45
1:B:380:LYS:HD3	4:B:2142:HOH:O	2.16	0.45
1:C:588:ALA:O	1:C:592:LYS:HG2	2.16	0.45
1:C:414:ARG:NH2	4:C:1768:HOH:O	2.49	0.45
1:B:341:LEU:CB	1:B:342:ILE:HD12	2.46	0.44
1:D:688:VAL:HG13	1:D:749:ALA:HB2	2.00	0.44
1:C:462:HIS:HD2	4:C:1636:HOH:O	2.00	0.44
1:A:51:ILE:HB	1:A:52:PRO:HA	1.99	0.44
1:A:88:VAL:HG13	1:A:149:ALA:HB2	1.98	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:GLU:HG2	1:A:180:LYS:HE2	1.99	0.43
1:C:572:PRO:HB3	1:D:798:LYS:HA	1.99	0.43
1:D:742:ILE:HD12	1:D:742:ILE:N	2.34	0.43
1:B:307:LYS:HB2	4:B:2044:HOH:O	2.18	0.43
1:A:167:LEU:HG	1:A:168:LEU:HD13	2.00	0.43
1:D:741:LEU:C	1:D:742:ILE:HD12	2.39	0.43
1:C:409:PHE:CE1	1:C:451:ILE:HD11	2.54	0.43
1:C:414:ARG:HD2	3:C:1600:GSH:SG2	2.58	0.43
1:C:421:ILE:HD13	1:C:472:THR:HG23	2.00	0.43
1:D:775:VAL:O	1:D:779:LYS:HG2	2.18	0.42
1:C:450:LYS:HB3	3:C:1600:GSH:HG13	2.02	0.42
1:C:519:LEU:HD22	4:C:1658:HOH:O	2.19	0.42
1:D:739:GLU:HG3	1:D:780:LYS:HE3	2.02	0.42
1:C:542:ILE:N	1:C:542:ILE:HD12	2.35	0.41
1:B:269:ARG:O	1:B:273:LYS:HG3	2.21	0.41
1:A:69:ARG:O	1:A:73:LYS:HG3	2.20	0.41
1:A:146:VAL:HG23	4:A:1934:HOH:O	2.21	0.41
1:B:262:HIS:O	1:B:263:GLN:HB2	2.21	0.41
1:D:753:TRP:CD1	1:D:777:LEU:HD13	2.55	0.41
1:B:251:ILE:HB	1:B:252:PRO:HA	2.02	0.41
1:C:408:TYR:CG	1:C:409:PHE:N	2.89	0.41
1:D:741:LEU:HG	1:D:777:LEU:HD11	2.02	0.41
1:C:541:LEU:CB	1:C:542:ILE:HD12	2.51	0.41
1:D:662:HIS:O	1:D:663:GLN:HB2	2.21	0.41
1:C:576:THR:CG2	1:D:765:PRO:HG3	2.42	0.40
1:A:181:VAL:HG21	4:A:1957:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	196/198 (99%)	191 (97%)	5 (3%)	0	100	100
1	B	196/198 (99%)	193 (98%)	3 (2%)	0	100	100
1	C	196/198 (99%)	192 (98%)	4 (2%)	0	100	100
1	D	196/198 (99%)	191 (97%)	5 (3%)	0	100	100
All	All	784/792 (99%)	767 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/178 (100%)	176 (99%)	2 (1%)	80	69
1	B	178/178 (100%)	177 (99%)	1 (1%)	90	85
1	C	178/178 (100%)	171 (96%)	7 (4%)	39	16
1	D	178/178 (100%)	175 (98%)	3 (2%)	68	51
All	All	712/712 (100%)	699 (98%)	13 (2%)	66	49

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	120	LEU
1	A	168	LEU
1	B	344	MET
1	C	403	ASN
1	C	438	ASP
1	C	509	GLN
1	C	514	GLN
1	C	544	MET
1	C	593	ARG
1	C	596	GLN
1	D	635	GLU
1	D	744	MET
1	D	777	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (24) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	36	GLN
1	A	62	HIS
1	A	80	ASN
1	A	114	GLN
1	A	123	ASN
1	A	170	ASN
1	A	171	HIS
1	B	228	GLN
1	B	262	HIS
1	B	323	ASN
1	B	370	ASN
1	C	403	ASN
1	C	410	ASN
1	C	462	HIS
1	C	480	ASN
1	C	509	GLN
1	C	570	ASN
1	D	628	GLN
1	D	662	HIS
1	D	680	ASN
1	D	709	GLN
1	D	770	ASN
1	D	796	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GSH	A	1200	-	13,19,19	0.85	1 (7%)	15,24,24	1.25	1 (6%)
3	GSH	B	1400	-	13,19,19	0.97	1 (7%)	15,24,24	1.27	2 (13%)
3	GSH	C	1600	-	13,19,19	0.77	1 (7%)	15,24,24	1.29	1 (6%)
3	GSH	D	1800	-	13,19,19	0.70	0	15,24,24	1.31	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GSH	A	1200	-	-	0/18/24/24	0/0/0/0
3	GSH	B	1400	-	-	0/18/24/24	0/0/0/0
3	GSH	C	1600	-	-	0/18/24/24	0/0/0/0
3	GSH	D	1800	-	-	0/18/24/24	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1600	GSH	CG1-CD1	2.01	1.55	1.51
3	A	1200	GSH	CG1-CD1	2.41	1.56	1.51
3	B	1400	GSH	CG1-CD1	2.88	1.57	1.51

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1600	GSH	CB1-CG1-CD1	-3.60	104.72	113.27
3	D	1800	GSH	CB1-CG1-CD1	-3.54	104.84	113.27
3	A	1200	GSH	CB1-CG1-CD1	-3.36	105.28	113.27
3	B	1400	GSH	CB1-CG1-CD1	-3.20	105.66	113.27

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	B	1400	GSH	CB2-CA2-C2	-2.69	103.59	109.66
3	D	1800	GSH	CB2-CA2-C2	-2.13	104.85	109.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1600	GSH	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	198/198 (100%)	0.18	6 (3%) 54 58	12, 20, 30, 37	0
1	B	198/198 (100%)	0.12	2 (1%) 84 87	11, 19, 27, 35	0
1	C	198/198 (100%)	0.36	11 (5%) 28 30	14, 22, 37, 40	0
1	D	198/198 (100%)	0.62	9 (4%) 37 41	14, 25, 41, 44	0
All	All	792/792 (100%)	0.32	28 (3%) 48 52	11, 21, 37, 44	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	637	ALA	4.6
1	D	722	TYR	4.6
1	D	636	GLN	4.3
1	A	111	VAL	4.1
1	B	344	MET	3.9
1	A	110	ASP	3.7
1	C	441	GLU	3.5
1	C	522	TYR	3.5
1	C	593	ARG	3.3
1	D	710	ASP	3.2
1	C	438	ASP	3.2
1	D	641	GLU	3.0
1	A	144	MET	3.0
1	C	435	GLU	2.9
1	C	437	ALA	2.9
1	A	107	LYS	2.8
1	C	440	PRO	2.8
1	D	602	PRO	2.6
1	D	640	PRO	2.5
1	C	544	MET	2.5
1	B	292	VAL	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	478	ALA	2.4
1	D	635	GLU	2.3
1	A	105	ALA	2.2
1	C	510	ASP	2.2
1	D	674	ASN	2.2
1	A	145	SER	2.0
1	C	596	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GSH	D	1800	20/20	0.83	0.15	1.00	26,28,31,31	0
3	GSH	A	1200	20/20	0.90	0.12	0.87	20,23,28,28	0
3	GSH	B	1400	20/20	0.92	0.11	0.73	18,20,23,23	0
3	GSH	C	1600	20/20	0.86	0.13	0.61	26,27,29,29	0
2	MG	A	1901	1/1	0.99	0.06	-2.82	22,22,22,22	0
2	MG	B	1902	1/1	0.99	0.02	-4.50	23,23,23,23	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.