



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 10, 2016 – 03:05 PM EDT

PDB ID : 5IZM
Title : The crystal structure of human eEFSec in complex with GDPNP
Authors : Dobosz-Bartoszek, M.; Otwinowski, Z.; Simonovic, M.
Deposited on : 2016-03-25
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1476 (3.50-3.30)
Clashscore	102246	1611 (3.50-3.30)
Ramachandran outliers	100387	1571 (3.50-3.30)
Sidechain outliers	100360	1571 (3.50-3.30)
RSRZ outliers	91569	1485 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	616	
1	B	616	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6632 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Selenocysteine-specific elongation factor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	460	3147	2026	544	554	9	14	0	1	0
1	B	495	3419	2212	579	604	9	15	0	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MSE	-	initiating methionine	UNP P57772
A	-18	GLY	-	expression tag	UNP P57772
A	-17	SER	-	expression tag	UNP P57772
A	-16	SER	-	expression tag	UNP P57772
A	-15	HIS	-	expression tag	UNP P57772
A	-14	HIS	-	expression tag	UNP P57772
A	-13	HIS	-	expression tag	UNP P57772
A	-12	HIS	-	expression tag	UNP P57772
A	-11	HIS	-	expression tag	UNP P57772
A	-10	HIS	-	expression tag	UNP P57772
A	-9	SER	-	expression tag	UNP P57772
A	-8	SER	-	expression tag	UNP P57772
A	-7	GLY	-	expression tag	UNP P57772
A	-6	LEU	-	expression tag	UNP P57772
A	-5	VAL	-	expression tag	UNP P57772
A	-4	PRO	-	expression tag	UNP P57772
A	-3	ARG	-	expression tag	UNP P57772
A	-2	GLY	-	expression tag	UNP P57772
A	-1	SER	-	expression tag	UNP P57772
A	0	HIS	-	expression tag	UNP P57772
A	1	MSE	-	expression tag	UNP P57772
B	-19	MSE	-	initiating methionine	UNP P57772
B	-18	GLY	-	expression tag	UNP P57772
B	-17	SER	-	expression tag	UNP P57772
B	-16	SER	-	expression tag	UNP P57772

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	HIS	-	expression tag	UNP P57772
B	-14	HIS	-	expression tag	UNP P57772
B	-13	HIS	-	expression tag	UNP P57772
B	-12	HIS	-	expression tag	UNP P57772
B	-11	HIS	-	expression tag	UNP P57772
B	-10	HIS	-	expression tag	UNP P57772
B	-9	SER	-	expression tag	UNP P57772
B	-8	SER	-	expression tag	UNP P57772
B	-7	GLY	-	expression tag	UNP P57772
B	-6	LEU	-	expression tag	UNP P57772
B	-5	VAL	-	expression tag	UNP P57772
B	-4	PRO	-	expression tag	UNP P57772
B	-3	ARG	-	expression tag	UNP P57772
B	-2	GLY	-	expression tag	UNP P57772
B	-1	SER	-	expression tag	UNP P57772
B	0	HIS	-	expression tag	UNP P57772
B	1	MSE	-	expression tag	UNP P57772

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mn 1 1	0	0
2	A	1	Total Mn 1 1	0	0

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter code: GNP) (formula: C₁₀H₁₇N₆O₁₃P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			
3	A	1	Total	32	10	6	13	3	0	0
3	B	1	Total	32	10	6	13	3	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Selenocysteine-specific elongation factor



- Molecule 1: Selenocysteine-specific elongation factor



ALA	GLU	ARG	SER	GLU	PRO	S572	Q573	H574	V575	L579	T580	F581	K582	V585	F586	D587	T588	H589	M592	V593	Q594	S595	PRO																														
I488	G490	R491	S492	L493	F494	LYS	LYS	E497	T498	M499	I500	Q501	E502	F503	V504	L510	S511	T512	G513	L518	D519	S520	G523	GLN	SER	GLY	K527	F528	R529	I530	H531	I532	P533	L544	THR	PRO	ALA	LEU	LYS	ARG	ALA	ALA	ALA	GLY	ARG	GLY	GLU	ALA	THR	ARG	GLN	GLU	SER
LYS	ALA	GLY	GLN	ALA	THR	GLU	GLY	HIS	CNS	P407	V414	V420	T421	L425	C426	L427	V428	I429	S431	R432	L433	D434	A435	ASP	ILE	H438	C442	R443	L444	A445	F446	H447	L450	G453	Y459	A460	L464	P465	Y470	K471	L472	G477	L478	V479	E480	M483							
E320	K321	I322	F323	Y324	F325	L329	Q330	T331	I337	T338	V344	M345	G346	R347	L348	M349	F350	F351	S352	P353	A354	P355	F358	D359	Q360	E361	P362	M368	F369	S370	Q371	E372	Y373	E377	Q378	Y379	L380	S381	LYS	ASP	LEU	THR	PRO	ALA	VAL	THR	ASP	ASN	ASP	GLU	ALA	ASP	LYS
T217	P220	F223	F224	L225	M226	F232	S233	S234	T239	V240	M241	I245	L246	S247	T250	S251	E257	I258	P259	A260	L261	K265	K266	V267	M270	Q271	M272	I277	A280	L286	T291	Q292	F293	D294	P295	K296	L297	L298	C304	L309	H310	T311	V312	V319									
G46	I47	T48	F53	L68	PRO	GLU	PHE	GLN	ALA	ALA	PRO	GLU	GLU	PRO	GLY	E83	P84	V88	T89	L90	V91	P94	G95	S98	L99	I100	I104	G105	G106	I109	I110	D111	L112	M113	M114	L115	T120	K121	G122	M123	Q124	T125	Q126	S127	A128	E129	G130	L131					
V132	I133	G134	Q135	I136	V144	L145	M146	K147	I148	D149	L150	L151	P152	E153	GLY	LYS	R156	I160	D161	R177	G178	A179	P180	I182	P183	V184	K187	L99	G189	G190	PRO	GLU	ALA	PRO	GLU	THR	GLU	ALA	F199	Q200	G201	I202	P203	E207	L208	L209	T210	S211	Q212	I213	S214	I215	P216

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	92.32Å 112.40Å 327.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.16 – 3.40 46.16 – 3.38	Depositor EDS
% Data completeness (in resolution range)	78.4 (46.16-3.40) 84.9 (46.16-3.38)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 3.40Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.235 , 0.285 0.254 , 0.311	Depositor DCC
R_{free} test set	1068 reflections (5.13%)	DCC
Wilson B-factor (Å ²)	41.6	Xtrriage
Anisotropy	0.039	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , 33.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.79	EDS
Total number of atoms	6632	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GNP, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/3193	0.77	12/4335 (0.3%)
1	B	0.37	0/3467	0.72	9/4708 (0.2%)
All	All	0.39	0/6660	0.75	21/9043 (0.2%)

There are no bond length outliers.

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	262	LYS	CB-CA-C	-7.40	95.59	110.40
1	B	225	LEU	N-CA-CB	-6.58	97.25	110.40
1	A	592	MSE	CB-CA-C	-6.37	97.66	110.40
1	B	445	ALA	CB-CA-C	-6.24	100.75	110.10
1	A	511	SER	N-CA-C	6.23	127.83	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3147	0	2871	206	0
1	B	3419	0	3163	219	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	32	0	13	7	0
3	B	32	0	13	2	0
All	All	6632	0	6060	424	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 33.

The worst 5 of 424 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:272:MSE:CB	1:B:286:LEU:HD23	1.20	1.64
1:B:272:MSE:CE	1:B:286:LEU:HD21	1.17	1.58
1:B:272:MSE:CG	1:B:286:LEU:HD23	1.45	1.43
1:B:272:MSE:HB2	1:B:286:LEU:CD2	1.55	1.37
1:B:272:MSE:CB	1:B:286:LEU:CD2	2.05	1.34

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/616 (72%)	350 (79%)	81 (18%)	10 (2%)	8	45
1	B	475/616 (77%)	377 (79%)	87 (18%)	11 (2%)	8	45
All	All	916/1232 (74%)	727 (79%)	168 (18%)	21 (2%)	8	45

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	590	LYS
1	A	296	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	359	ASP
1	A	439	THR
1	A	511	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/507 (55%)	276 (99%)	2 (1%)	88	95
1	B	312/507 (62%)	309 (99%)	3 (1%)	82	93
All	All	590/1014 (58%)	585 (99%)	5 (1%)	86	94

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	ILE
1	A	507	LYS
1	B	233	SER
1	B	286	LEU
1	B	594	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	330	GLN
1	B	7	ASN
1	B	9	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GNP	A	1002	2	29,34,34	5.15	11 (37%)	28,54,54	1.64	7 (25%)
3	GNP	B	1002	2	29,34,34	5.14	11 (37%)	28,54,54	1.64	7 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GNP	A	1002	2	-	0/16/38/38	0/3/3/3
3	GNP	B	1002	2	-	0/16/38/38	0/3/3/3

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1002	GNP	C5-C6	-10.81	1.33	1.53
3	A	1002	GNP	C5-C6	-10.75	1.33	1.53
3	A	1002	GNP	C4-N9	-10.74	1.33	1.47
3	B	1002	GNP	C4-N9	-10.71	1.33	1.47
3	B	1002	GNP	C8-N9	-4.35	1.33	1.47

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1002	GNP	PA-O3A-PB	-3.51	119.96	132.71
3	A	1002	GNP	PA-O3A-PB	-3.51	119.99	132.71
3	B	1002	GNP	O2G-PG-O1G	-3.23	105.07	113.58
3	A	1002	GNP	O2G-PG-O1G	-3.23	105.09	113.58
3	B	1002	GNP	C4'-O4'-C1'	-2.52	103.96	109.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1002	GNP	7	0
3	B	1002	GNP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	446/616 (72%)	-0.03	12 (2%) 58 53	6, 31, 70, 104	0
1	B	480/616 (77%)	-0.19	3 (0%) 90 86	8, 28, 55, 76	0
All	All	926/1232 (75%)	-0.11	15 (1%) 74 69	6, 30, 64, 104	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	359	ASP	3.2
1	A	514	GLU	2.8
1	A	536	LEU	2.8
1	A	515	LEU	2.6
1	A	534	GLY	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GNP	B	1002	32/32	0.89	0.26	0.73	24,54,66,96	0
3	GNP	A	1002	32/32	0.90	0.27	0.59	17,47,78,96	0
2	MN	B	1001	1/1	0.97	0.13	-1.78	15,15,15,15	0
2	MN	A	1001	1/1	0.97	0.07	-3.31	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.