



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 5, 2016 – 02:56 PM EDT

PDB ID : 5J1I
Title : Structure of the spectrin repeats 7, 8, and 9 of the plakin domain of plectin
Authors : ORTEGA, E.; DE PEREDA, J.M.
Deposited on : 2016-03-29
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939



i

X-RAY DIFFRACTION

A.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

Mol	Chain	Length	Quality of chain
1	A	373	 <div> <div></div> <div>3%</div> <div>90%</div> <div>7%</div> </div>
1	B	373	 <div> <div></div> <div>6%</div> <div>89%</div> <div>8%</div> </div>

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 5179 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Plectin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	347	Total	C	N	O	S	0	0	1
			2550	1581	473	491	5			
1	B	345	Total	C	N	O	S	0	0	0
			2629	1626	486	511	6			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1000	GLY	-	expression tag	UNP Q15149
A	1001	SER	-	expression tag	UNP Q15149
A	1002	HIS	-	expression tag	UNP Q15149
A	1003	MET	-	expression tag	UNP Q15149
B	1000	GLY	-	expression tag	UNP Q15149
B	1001	SER	-	expression tag	UNP Q15149
B	1002	HIS	-	expression tag	UNP Q15149
B	1003	MET	-	expression tag	UNP Q15149


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Plectin



• Molecule 1: Plectin



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	75.24Å 90.70Å 154.54Å 90.00° 99.13° 90.00°	Depositor
Resolution (Å)	48.88 – 2.80 48.88 – 2.80	Depositor EDS
% Data completeness (in resolution range)	55.6 (48.88-2.80) 55.7 (48.88-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.81 (at 2.81Å)	Xtriage
Refinement program	PHENIX (dev_2299: ???)	Depositor
R, R_{free}	0.296 , 0.322 0.291 , 0.316	Depositor DCC
R_{free} test set	690 reflections (4.90%)	DCC
Wilson B-factor (Å ²)	64.6	Xtriage
Anisotropy	0.080	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	5179	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/2574	0.49	0/3488
1	B	0.29	0/2654	0.45	0/3587
All	All	0.30	0/5228	0.47	0/7075

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2550	0	2392	7	0
1	B	2629	0	2513	6	0
All	All	5179	0	4905	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1138:GLN:OE1	1:B:1162:LEU:HD13	1.99	0.62
1:A:1355:THR:O	1:A:1359:GLN:HG2	2.09	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1236:LEU:HD13	1:A:1302:TYR:CZ	2.45	0.51
1:B:1152:LEU:HD11	1:B:1229:LEU:HB3	1.93	0.50
1:A:1132:LEU:HD12	1:A:1211:LEU:HD22	1.94	0.49
1:A:1224:VAL:O	1:A:1228:GLU:HG2	2.12	0.49
1:A:1140:LYS:HG2	1:A:1317:LYS:HG3	1.97	0.47
1:B:1066:GLY:HA2	1:B:1069:ARG:NH1	2.31	0.45
1:B:1127:GLY:O	1:B:1131:VAL:HG23	2.18	0.44
1:A:1009:CYS:HB2	1:A:1077:VAL:HG11	2.00	0.42
1:B:1251:GLN:O	1:B:1255:ARG:HG3	2.20	0.41
1:A:1140:LYS:HE3	1:A:1320:LEU:HB2	2.03	0.40
1:B:1340:ILE:O	1:B:1344:VAL:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/373 (91%)	338 (100%)	1 (0%)	0	100	100
1	B	335/373 (90%)	334 (100%)	1 (0%)	0	100	100
All	All	674/746 (90%)	672 (100%)	2 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230 / 319 (72%)	230 (100%)	0	100	100
1	B	250 / 319 (78%)	250 (100%)	0	100	100
All	All	480 / 638 (75%)	480 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	347/373 (93%)	0.32	11 (3%) 51 39	46, 71, 126, 146	0
1	B	345/373 (92%)	0.35	21 (6%) 25 15	58, 81, 108, 130	0
All	All	692/746 (92%)	0.34	32 (4%) 36 25	46, 78, 117, 146	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1078	LEU	4.7
1	A	1094	LEU	4.1
1	A	1009	CYS	4.0
1	A	1270	ALA	3.9
1	B	1085	PRO	3.4
1	B	1150	PRO	3.4
1	A	1090	LEU	3.3
1	B	1091	ARG	3.3
1	B	1215	TRP	3.2
1	A	1365	SER	3.2
1	B	1222	THR	2.9
1	B	1226	GLN	2.9
1	B	1094	LEU	2.9
1	B	1132	LEU	2.8
1	A	1084	SER	2.7
1	B	1078	LEU	2.6
1	A	1098	LEU	2.5
1	B	1013	ILE	2.5
1	A	1077	VAL	2.5
1	B	1166	ALA	2.5
1	B	1019	ILE	2.5
1	B	1227	ARG	2.4
1	B	1250	LEU	2.3
1	B	1089	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1177	ARG	2.2
1	B	1098	LEU	2.2
1	A	1032	HIS	2.2
1	A	1070	LEU	2.2
1	B	1118	ILE	2.1
1	B	1246	LEU	2.1
1	B	1179	GLU	2.1
1	B	1088	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.