



## wwPDB EM Map/Model Validation Report ⓘ

Apr 10, 2016 – 01:50 PM BST

PDB ID : 3J5P  
EMDB ID: : EMD-5778  
Title : Structure of TRPV1 ion channel determined by single particle electron cryo-microscopy  
Authors : Liao, M.; Cao, E.; Julius, D.; Cheng, Y.  
Deposited on : 2013-10-28  
Resolution : 3.27 Å(reported)

This is a wwPDB EM Map/Model Validation Report for a publicly released PDB/EMDB entry.  
For rigid body fitted models, validation errors reported here could stem from errors in the original structure(s) used in the fitting.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/EMValidationReportHelp>

---

MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk27241


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	114402	924
Ramachandran outliers	111179	726
Sidechain outliers	111093	686

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	598	
1	B	598	
1	C	598	
1	D	598	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 18636 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Transient receptor potential cation channel subfamily V member 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	B	592	Total	C	N	O	S	0	0
			4659	3027	765	842	25		
1	A	592	Total	C	N	O	S	0	0
			4659	3027	765	842	25		
1	C	592	Total	C	N	O	S	0	0
			4659	3027	765	842	25		
1	D	592	Total	C	N	O	S	0	0
			4659	3027	765	842	25		


### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transient receptor potential cation channel subfamily V member 1




- Molecule 1: Transient receptor potential cation channel subfamily V member 1






• Molecule 1: Transient receptor potential cation channel subfamily V member 1

Chain C: 37% 43% 19%



• Molecule 1: Transient receptor potential cation channel subfamily V member 1

Chain D: 



## 4 Experimental information

Property	Value	Source
Reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	Depositor
Number of images	35645	Depositor
Resolution determination method	Gold standard FSC at 0.143 cut-off	Depositor
CTF correction method	Each particle	Depositor
Microscope	FEI POLARA 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	21	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	31000	Depositor
Image detector	K2 Summit in counting mode	Depositor

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 2$	RMSZ	# $ Z  > 2$
1	A	0.52	9/4706 (0.2%)	0.66	6/6370 (0.1%)
1	B	0.52	9/4706 (0.2%)	0.66	6/6370 (0.1%)
1	C	0.52	9/4706 (0.2%)	0.66	6/6370 (0.1%)
1	D	0.52	9/4706 (0.2%)	0.66	6/6370 (0.1%)
All	All	0.52	36/18824 (0.2%)	0.66	24/25480 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6
1	B	0	6
1	C	0	6
1	D	0	6
All	All	0	24

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	162	MET	CG-SD	7.74	2.01	1.81
1	D	162	MET	CG-SD	7.72	2.01	1.81
1	C	162	MET	CG-SD	7.71	2.01	1.81
1	A	162	MET	CG-SD	7.71	2.01	1.81
1	A	214	MET	CG-SD	6.28	1.97	1.81

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	460	LEU	C-N-CD	5.71	140.39	128.40
1	A	460	LEU	C-N-CD	5.70	140.37	128.40
1	C	500	ARG	C-N-CD	5.69	140.35	128.40

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	460	LEU	C-N-CD	5.68	140.34	128.40
1	B	500	ARG	C-N-CD	5.68	140.33	128.40

There are no chirality outliers.

5 of 24 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	243	PRO	Peptide
1	B	244	GLY	Peptide
1	B	247	PHE	Peptide
1	B	248	GLY	Peptide
1	B	376	PRO	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4659	0	4609	782	0
1	B	4659	0	4609	778	0
1	C	4659	0	4609	780	0
1	D	4659	0	4609	773	0
All	All	18636	0	18436	2888	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 78.

The worst 5 of 2888 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:374:TYR:CE2	1:A:235:PHE:CE1	1.86	1.62
1:C:374:TYR:CE2	1:D:235:PHE:CE1	1.89	1.61
1:C:374:TYR:CD2	1:D:235:PHE:CZ	1.90	1.59
1:B:235:PHE:CZ	1:D:374:TYR:CD2	1.91	1.58
1:B:374:TYR:CD2	1:A:235:PHE:CZ	1.88	1.58

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	577/598 (96%)	544 (94%)	33 (6%)	0	100	100
1	B	577/598 (96%)	544 (94%)	33 (6%)	0	100	100
1	C	577/598 (96%)	544 (94%)	33 (6%)	0	100	100
1	D	577/598 (96%)	544 (94%)	33 (6%)	0	100	100
All	All	2308/2392 (96%)	2176 (94%)	132 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	493/519 (95%)	359 (73%)	134 (27%)	0	2
1	B	493/519 (95%)	359 (73%)	134 (27%)	0	2
1	C	493/519 (95%)	359 (73%)	134 (27%)	0	2
1	D	493/519 (95%)	359 (73%)	134 (27%)	0	2
All	All	1972/2076 (95%)	1436 (73%)	536 (27%)	2	2

5 of 536 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	632	SER
1	C	325	LEU
1	D	545	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	655	PHE
1	C	136	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	423	GLN
1	C	269	GLN
1	D	419	ASN
1	A	498	GLN
1	A	691	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.