



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:42 PM GMT

PDB ID : 4J6N
Title : Crystal structure of calcium²⁺-free wild-type CD23 lectin domain (crystal form E)
Authors : Dhaliwal, B.; Pang, M.O.Y.; Sutton, B.J.
Deposited on : 2013-02-11
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2228 (2.90-2.82)
Clashscore	102246	2499 (2.90-2.82)
Ramachandran outliers	100387	2439 (2.90-2.82)
Sidechain outliers	100360	2442 (2.90-2.82)
RSRZ outliers	91569	2236 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	143	
1	B	143	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2186 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Low affinity immunoglobulin epsilon Fc receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	135	Total	C	N	O	S	0	2	0
			1093	689	193	199	12			
1	B	135	Total	C	N	O	S	0	2	0
			1093	689	193	199	12			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Low affinity immunoglobulin epsilon Fc receptor



- Molecule 1: Low affinity immunoglobulin epsilon Fc receptor



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	64.26 Å 73.07 Å 57.04 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.25 – 2.85 48.25 – 2.85	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.25-2.85) 99.8 (48.25-2.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 2.86 Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.252 , 0.317 0.307 , 0.308	Depositor DCC
R_{free} test set	604 reflections (10.00%)	DCC
Wilson B-factor (Å ²)	29.8	Xtriage
Anisotropy	1.009	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 57.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	0 of 6663 reflections	Xtriage
F_o, F_c correlation	0.77	EDS
Total number of atoms	2186	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/1133	0.82	2/1535 (0.1%)
1	B	0.43	0/1133	0.70	0/1535
All	All	0.44	0/2266	0.76	2/3070 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	254	SER	CB-CA-C	10.75	130.52	110.10
1	A	254	SER	N-CA-C	-6.76	92.74	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1093	0	1024	29	0
1	B	1093	0	1024	13	0
All	All	2186	0	2048	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:227:ASP:O	1:A:229:LYS:N	1.85	1.10
1:A:227:ASP:O	1:A:230:GLY:N	1.91	1.04
1:A:252:SER:HB2	1:A:254:SER:O	1.64	0.96
1:A:254:SER:HB3	1:A:258:ASP:OD1	1.82	0.80
1:A:227:ASP:C	1:A:229:LYS:H	1.80	0.80
1:B:191:CYS:HG	1:B:282:CYS:HG	1.30	0.77
1:A:252:SER:CB	1:A:254:SER:O	2.33	0.76
1:A:183:GLN:HB2	1:A:186:HIS:CD2	2.21	0.76
1:A:223:LEU:HB3	1:A:260:VAL:HB	1.74	0.69
1:A:191:CYS:HG	1:A:282:CYS:HG	1.38	0.68
1:B:183:GLN:HB2	1:B:186:HIS:CD2	2.31	0.66
1:A:227:ASP:OD1	1:A:228:LEU:N	2.31	0.63
1:A:176:TYR:HB3	1:A:282:CYS:HB2	1.81	0.61
1:B:176:TYR:HB3	1:B:282:CYS:HB2	1.82	0.60
1:B:186:HIS:H	1:B:186:HIS:CD2	2.21	0.59
1:A:226:LEU:HB2	1:A:229:LYS:HE3	1.86	0.57
1:A:227:ASP:HB3	1:A:256:GLY:HA2	1.87	0.57
1:B:223:LEU:HB3	1:B:260:VAL:HB	1.88	0.56
1:A:227:ASP:C	1:A:230:GLY:H	2.08	0.55
1:A:186:HIS:CD2	1:A:186:HIS:H	2.24	0.54
1:B:246:ALA:HB1	1:B:247:PRO:HD2	1.89	0.53
1:A:160:CYS:HB3	1:A:286:ALA:HB3	1.91	0.52
1:A:160:CYS:HB2	1:A:172:ARG:HG2	1.92	0.51
1:A:246:ALA:HB1	1:A:247:PRO:HD2	1.92	0.51
1:A:181:THR:HG23	1:A:276:LYS:HB3	1.94	0.50
1:A:227:ASP:O	1:A:229:LYS:C	2.49	0.50
1:B:186:HIS:N	1:B:186:HIS:CD2	2.80	0.49
1:A:252:SER:C	1:A:254:SER:H	2.16	0.49
1:B:181:THR:HG23	1:B:276:LYS:HB3	1.95	0.48
1:A:227:ASP:O	1:A:229:LYS:CA	2.62	0.47
1:A:159:VAL:O	1:A:289:THR:HB	2.15	0.47
1:A:186:HIS:CD2	1:A:186:HIS:N	2.83	0.46
1:B:186:HIS:H	1:B:186:HIS:HD2	1.63	0.46
1:A:226:LEU:O	1:A:229:LYS:HG2	2.15	0.45
1:B:184:TRP:HD1	1:B:220:TRP:CE3	2.34	0.45
1:B:224:ARG:HH21	1:B:235:VAL:HA	1.83	0.43
1:B:159:VAL:O	1:B:289:THR:HB	2.18	0.43
1:A:163:CYS:SG	1:A:174:CYS:SG	3.01	0.41
1:A:183:GLN:HB2	1:A:186:HIS:CG	2.55	0.41
1:A:223:LEU:HD23	1:A:260:VAL:HG21	2.02	0.41
1:A:170:PHE:HB3	1:A:175:TYR:HE1	1.86	0.40
1:B:162:THR:HG22	1:B:163:CYS:H	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	135/143 (94%)	115 (85%)	15 (11%)	5 (4%)	4	14
1	B	135/143 (94%)	117 (87%)	17 (13%)	1 (1%)	26	59
All	All	270/286 (94%)	232 (86%)	32 (12%)	6 (2%)	8	27

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	228	LEU
1	A	227	ASP
1	A	255	GLN
1	A	257	GLU
1	A	250	PRO
1	B	250	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/120 (98%)	103 (88%)	14 (12%)	6	16
1	B	117/120 (98%)	107 (92%)	10 (8%)	13	34
All	All	234/240 (98%)	210 (90%)	24 (10%)	9	24

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	163	CYS
1	A	186	HIS
1	A	219	SER
1	A	224	ARG
1	A	226	LEU
1	A	227	ASP
1	A	231	GLU
1	A	233	ILE
1	A	241	ASP
1	A	253	ARG
1	A	257	GLU
1	A	258	ASP
1	A	273	CYS
1	A	289	THR
1	B	159	VAL
1	B	161	ASN
1	B	162	THR
1	B	163	CYS
1	B	219	SER
1	B	226	LEU
1	B	228	LEU
1	B	241	ASP
1	B	253	ARG
1	B	289	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	183	GLN
1	A	186	HIS
1	B	186	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	135/143 (94%)	0.60	14 (10%) 8 5	14, 32, 82, 155	0
1	B	135/143 (94%)	1.92	48 (35%) 0 0	27, 69, 123, 173	0
All	All	270/286 (94%)	1.26	62 (22%) 1 0	14, 49, 114, 173	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	292	ALA	17.9
1	B	291	PRO	15.0
1	A	292	ALA	10.0
1	B	290	PRO	9.0
1	B	289	THR	8.1
1	B	288	CYS	6.9
1	A	291	PRO	6.5
1	B	158	PHE	6.5
1	B	257	GLU	5.9
1	A	158	PHE	5.5
1	B	256	GLY	5.2
1	A	159	VAL	4.8
1	B	252	SER	4.8
1	B	210	LEU	4.6
1	B	253	ARG	4.1
1	A	290	PRO	4.1
1	A	228	LEU	4.1
1	B	171	GLN	4.0
1	A	255	GLN	4.0
1	B	216	HIS	3.9
1	B	228	LEU	3.6
1	B	247	PRO	3.6
1	A	230	GLY	3.5
1	B	254	SER	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	226	LEU	3.4
1	B	209	PHE	3.4
1	B	227	ASP	3.2
1	B	169	ASN	3.1
1	B	240	VAL	3.1
1	B	229	LYS	3.1
1	A	256	GLY	3.1
1	B	264	GLY	3.0
1	B	159	VAL	3.0
1	B	282	CYS	3.0
1	B	286	ALA	2.9
1	B	211	THR	2.9
1	B	285	LEU	2.9
1	B	186	HIS	2.9
1	B	176	TYR	2.9
1	B	174	CYS	2.8
1	B	167	TRP	2.8
1	B	202	HIS	2.7
1	B	170	PHE	2.6
1	B	255	GLN	2.6
1	A	254	SER	2.6
1	B	208	ASP	2.6
1	B	239	HIS	2.6
1	B	192	ASP	2.6
1	B	162	THR	2.5
1	B	251	THR	2.4
1	B	248	GLY	2.4
1	B	175	TYR	2.3
1	B	160	CYS	2.3
1	B	284	ARG	2.2
1	A	227	ASP	2.2
1	A	226	LEU	2.2
1	B	287	THR	2.2
1	B	168[A]	ILE	2.2
1	A	183	GLN	2.2
1	A	160	CYS	2.1
1	B	265	SER	2.1
1	B	219	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.