



## wwPDB EM Map/Model Validation Report ⓘ

Apr 10, 2016 – 02:03 PM BST

PDB ID : 3J9V  
EMDB ID: : EMD-6286  
Title : Yeast V-ATPase state 3  
Authors : Zhao, J.; Benlekbir, S.; Rubinstein, J.L.  
Deposited on : 2015-02-23  
Resolution : 8.30 Å(reported)  
Based on PDB ID : 4DL0, 4RND, 1HO8, 1U7L

This is a wwPDB EM Map/Model Validation Report for a publicly released PDB/EMDB entry.  
For rigid body fitted models, validation errors reported here could stem from errors in the original structure(s) used in the fitting.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/EMValidationReportHelp>

---

MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk27241


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 8.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






















Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	114402	924
Ramachandran outliers	111179	726
Sidechain outliers	111093	686

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	b	840	27% 9% • 63%
2	O	392	79% 17% •
3	M	256	61% 18% •• 18%
4	N	118	80% 16% ••
5	A	616	72% 21% ••
5	C	616	70% 22% ••
5	E	616	68% 22% 5% ••
6	B	517	65% 19% 5% 12%
6	D	517	65% 20% • 12%

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	F	517	
7	Q	345	
8	H	114	
8	J	114	
8	L	114	
9	G	233	
9	I	233	
9	K	233	
10	P	478	
11	R	160	
11	S	160	
11	T	160	
11	U	160	
11	V	160	
11	W	160	
11	X	160	
11	Y	160	
11	Z	160	
11	a	160	

## 2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 57659 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called V-type proton ATPase subunit a, vacuolar isoform.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	b	312	Total	C	N	O	S	0	0
			2540	1614	434	489	3		

- Molecule 2 is a protein called V-type proton ATPase subunit C.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	O	392	Total	C	N	O	S	0	0
			3122	2005	516	596	5		

- Molecule 3 is a protein called V-type proton ATPase subunit D.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	M	210	Total	C	N	O	S	0	0
			1691	1061	305	321	4		

- Molecule 4 is a protein called V-type proton ATPase subunit F.

Mol	Chain	Residues	Atoms				AltConf	Trace
4	N	115	Total	C	N	O	0	0
			928	589	157	182		

- Molecule 5 is a protein called V-type proton ATPase catalytic subunit A.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	593	Total	C	N	O	S	0	0
			4578	2904	760	894	20		
5	A	593	Total	C	N	O	S	0	0
			4578	2904	760	894	20		
5	C	593	Total	C	N	O	S	0	0
			4578	2904	760	894	20		

- Molecule 6 is a protein called V-type proton ATPase subunit B.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	457	Total	C	N	O	S	0	0
			3585	2266	612	695	12		
6	B	457	Total	C	N	O	S	0	0
			3585	2266	612	695	12		
6	D	457	Total	C	N	O	S	0	0
			3585	2266	612	695	12		

- Molecule 7 is a protein called V-type proton ATPase subunit d.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	Q	345	Total	C	N	O	S	0	0
			2802	1779	454	555	14		

- Molecule 8 is a protein called V-type proton ATPase subunit G.

Mol	Chain	Residues	Atoms				AltConf	Trace
8	J	105	Total	C	N	O	0	0
			824	517	144	163		
8	L	105	Total	C	N	O	0	0
			824	517	144	163		
8	H	105	Total	C	N	O	0	0
			824	517	144	163		

- Molecule 9 is a protein called V-type proton ATPase subunit E.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	I	217	Total	C	N	O	S	0	0
			1731	1083	296	347	5		
9	K	217	Total	C	N	O	S	0	0
			1731	1083	296	347	5		
9	G	217	Total	C	N	O	S	0	0
			1731	1083	296	347	5		


- Molecule 10 is a protein called V-type proton ATPase subunit H.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	P	461	Total	C	N	O	S	0	0
			3712	2373	623	704	12		


- Molecule 11 is a protein called V-type proton ATPase subunit c.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	R	150	Total 1071	C 704	N 173	O 187	S 7	0	0
11	U	150	Total 1071	C 704	N 173	O 187	S 7	0	0
11	X	150	Total 1071	C 704	N 173	O 187	S 7	0	0
11	Y	150	Total 1071	C 704	N 173	O 187	S 7	0	0
11	W	150	Total 1071	C 704	N 173	O 187	S 7	0	0
11	Z	150	Total 1071	C 704	N 173	O 187	S 7	0	0
11	V	150	Total 1071	C 704	N 173	O 187	S 7	0	0
11	a	150	Total 1071	C 704	N 173	O 187	S 7	0	0
11	S	150	Total 1071	C 704	N 173	O 187	S 7	0	0
11	T	150	Total 1071	C 704	N 173	O 187	S 7	0	0







• Molecule 3: V-type proton ATPase subunit D



• Molecule 4: V-type proton ATPase subunit F



• Molecule 5: V-type proton ATPase catalytic subunit A









• Molecule 5: V-type proton ATPase catalytic subunit A

Chain A: 72% 21%




• Molecule 5: V-type proton ATPase catalytic subunit A


Chain C: 70% 22%



• Molecule 6: V-type proton ATPase subunit B


Chain F: 63% 21% 12%






• Molecule 6: V-type proton ATPase subunit B

Chain B: 65% 19% 5% 12%




• Molecule 6: V-type proton ATPase subunit B


Chain D: 65% 20% 12%




• Molecule 7: V-type proton ATPase subunit d




- Molecule 8: V-type proton ATPase subunit G




- Molecule 8: V-type proton ATPase subunit G




- Molecule 8: V-type proton ATPase subunit G




- Molecule 9: V-type proton ATPase subunit E




- Molecule 9: V-type proton ATPase subunit E




- Molecule 9: V-type proton ATPase subunit E




- Molecule 10: V-type proton ATPase subunit H




- Molecule 11: V-type proton ATPase subunit c



- Molecule 11: V-type proton ATPase subunit c




- Molecule 11: V-type proton ATPase subunit c






- Molecule 11: V-type proton ATPase subunit c

Chain Y: 75% 18% 6%



- Molecule 11: V-type proton ATPase subunit c

Chain W: 69% 21% 6%



- Molecule 11: V-type proton ATPase subunit c

Chain Z: 71% 21% 6%



- Molecule 11: V-type proton ATPase subunit c

Chain V: 81% 13% 6%




- Molecule 11: V-type proton ATPase subunit c

Chain a: 78% 14% 6%



- Molecule 11: V-type proton ATPase subunit c

Chain S: 72% 20% 6%




- Molecule 11: V-type proton ATPase subunit c

Chain T: 

72%

21%

6%



## 4 Experimental information

Property	Value	Source
Reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	Depositor
Number of images	17595	Depositor
Resolution determination method	FSC 0.143	Depositor
CTF correction method	Each particle	Depositor
Microscope	FEI TECNAI F20	Depositor
Voltage (kV)	200	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	30	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	7000	Depositor
Magnification	34483	Depositor
Image detector	GATAN K2 (4k x 4k)	Depositor

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >2	RMSZ	# Z  >2
1	b	1.76	22/2578 (0.9%)	2.07	83/3479 (2.4%)
10	P	1.68	25/3766 (0.7%)	1.87	74/5087 (1.5%)
11	R	1.63	3/1086 (0.3%)	2.03	30/1472 (2.0%)
11	S	1.77	7/1086 (0.6%)	1.79	19/1472 (1.3%)
11	T	1.60	4/1086 (0.4%)	1.97	25/1472 (1.7%)
11	U	1.65	6/1086 (0.6%)	2.05	23/1472 (1.6%)
11	V	1.67	4/1086 (0.4%)	1.84	17/1472 (1.2%)
11	W	1.67	9/1086 (0.8%)	1.95	30/1472 (2.0%)
11	X	1.72	7/1086 (0.6%)	1.86	22/1472 (1.5%)
11	Y	1.63	3/1086 (0.3%)	1.90	27/1472 (1.8%)
11	Z	1.68	7/1086 (0.6%)	1.87	18/1472 (1.2%)
11	a	1.68	7/1086 (0.6%)	1.97	24/1472 (1.6%)
2	O	1.68	27/3185 (0.8%)	1.90	63/4314 (1.5%)
3	M	1.76	16/1710 (0.9%)	1.87	43/2295 (1.9%)
4	N	1.68	8/944 (0.8%)	1.83	14/1277 (1.1%)
5	A	1.72	46/4677 (1.0%)	1.95	105/6339 (1.7%)
5	C	1.71	35/4677 (0.7%)	1.96	108/6339 (1.7%)
5	E	1.75	61/4677 (1.3%)	1.97	125/6339 (2.0%)
6	B	1.72	38/3654 (1.0%)	1.99	90/4953 (1.8%)
6	D	1.72	31/3654 (0.8%)	1.98	88/4953 (1.8%)
6	F	1.74	34/3654 (0.9%)	1.91	90/4953 (1.8%)
7	Q	1.74	28/2861 (1.0%)	1.96	78/3880 (2.0%)
8	H	1.70	8/828 (1.0%)	1.76	10/1098 (0.9%)
8	J	1.63	5/828 (0.6%)	1.72	9/1098 (0.8%)
8	L	1.59	3/828 (0.4%)	1.76	13/1098 (1.2%)
9	G	1.68	12/1743 (0.7%)	1.86	33/2338 (1.4%)
9	I	1.70	10/1743 (0.6%)	1.84	30/2338 (1.3%)
9	K	1.67	9/1743 (0.5%)	1.84	29/2338 (1.2%)
All	All	1.70	475/58610 (0.8%)	1.93	1320/79236 (1.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.



Mol	Chain	#Chirality outliers	#Planarity outliers
1	b	0	11
10	P	0	9
11	R	0	3
11	S	0	1
11	T	0	1
11	U	0	3
11	W	0	1
11	X	0	7
11	Y	0	3
11	Z	0	2
11	a	0	3
2	O	0	10
3	M	0	4
5	A	0	14
5	C	0	17
5	E	0	19
6	B	0	20
6	D	0	8
6	F	0	11
7	Q	0	6
8	J	0	2
8	L	0	1
9	G	0	4
9	I	0	3
9	K	0	4
All	All	0	167

The worst 5 of 475 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	F	393	ARG	CZ-NH2	9.92	1.46	1.33
10	P	246	SER	CA-CB	9.62	1.67	1.52
6	B	257	ARG	NE-CZ	8.88	1.44	1.33
5	E	334	TYR	CE1-CZ	8.71	1.49	1.38
6	F	101	ARG	CZ-NH1	8.52	1.44	1.33

The worst 5 of 1320 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	62	ARG	NE-CZ-NH1	19.24	129.92	120.30
11	U	12	PHE	CB-CG-CD1	18.18	133.52	120.80
1	b	285	TYR	CB-CG-CD1	16.96	131.18	121.00
6	D	325	ARG	NE-CZ-NH2	-16.76	111.92	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	b	285	TYR	CB-CG-CD2	-16.54	111.07	121.00

There are no chirality outliers.

5 of 167 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	b	61	ARG	Sidechain
1	b	69	TYR	Sidechain
1	b	73	TYR	Sidechain
1	b	86	GLY	Peptide
1	b	91	TYR	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	b	2540	0	2537	0	0
2	O	3122	0	3155	6	0
3	M	1691	0	1740	7	0
4	N	928	0	926	1	0
5	A	4578	0	4519	15	0
5	C	4578	0	4519	23	0
5	E	4578	0	4519	20	0
6	B	3585	0	3567	11	0
6	D	3585	0	3567	14	0
6	F	3585	0	3567	14	0
7	Q	2802	0	2689	11	0
8	H	824	0	877	1	0
8	J	824	0	877	0	0
8	L	824	0	877	0	0
9	G	1731	0	1797	2	0
9	I	1731	0	1797	6	0
9	K	1731	0	1797	2	0
10	P	3712	0	3829	13	0
11	R	1071	0	1141	5	0
11	S	1071	0	1141	4	0
11	T	1071	0	1141	8	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
11	U	1071	0	1141	0	0
11	V	1071	0	1141	2	0
11	W	1071	0	1141	8	0
11	X	1071	0	1141	7	0
11	Y	1071	0	1141	2	0
11	Z	1071	0	1141	7	0
11	a	1071	0	1141	0	0
All	All	57659	0	58566	169	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 169 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:S:66:TYR:HB3	11:S:144:LEU:HD22	1.32	1.07
11:T:66:TYR:HB3	11:T:144:LEU:HD22	1.36	1.06
11:X:66:TYR:HB3	11:X:144:LEU:HD22	1.38	1.02
11:R:66:TYR:HB3	11:R:144:LEU:HD22	1.45	0.97
11:X:66:TYR:CB	11:X:144:LEU:HD22	2.05	0.86

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	b	306/840 (36%)	283 (92%)	16 (5%)	7 (2%)	8	48
2	O	390/392 (100%)	359 (92%)	19 (5%)	12 (3%)	5	42
3	M	208/256 (81%)	201 (97%)	6 (3%)	1 (0%)	34	77
4	N	113/118 (96%)	103 (91%)	8 (7%)	2 (2%)	11	53

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	A	591/616 (96%)	543 (92%)	34 (6%)	14 (2%)	7	47
5	C	591/616 (96%)	540 (91%)	35 (6%)	16 (3%)	6	45
5	E	591/616 (96%)	536 (91%)	43 (7%)	12 (2%)	9	51
6	B	455/517 (88%)	415 (91%)	32 (7%)	8 (2%)	11	53
6	D	455/517 (88%)	406 (89%)	34 (8%)	15 (3%)	5	40
6	F	455/517 (88%)	405 (89%)	39 (9%)	11 (2%)	7	47
7	Q	343/345 (99%)	312 (91%)	24 (7%)	7 (2%)	9	51
8	H	103/114 (90%)	101 (98%)	0	2 (2%)	10	52
8	J	103/114 (90%)	99 (96%)	2 (2%)	2 (2%)	10	52
8	L	103/114 (90%)	99 (96%)	2 (2%)	2 (2%)	10	52
9	G	215/233 (92%)	205 (95%)	8 (4%)	2 (1%)	21	67
9	I	215/233 (92%)	209 (97%)	6 (3%)	0	100	100
9	K	215/233 (92%)	207 (96%)	5 (2%)	3 (1%)	14	58
10	P	457/478 (96%)	429 (94%)	19 (4%)	9 (2%)	9	51
11	R	148/160 (92%)	138 (93%)	6 (4%)	4 (3%)	6	45
11	S	148/160 (92%)	139 (94%)	7 (5%)	2 (1%)	14	58
11	T	148/160 (92%)	137 (93%)	7 (5%)	4 (3%)	6	45
11	U	148/160 (92%)	135 (91%)	10 (7%)	3 (2%)	9	51
11	V	148/160 (92%)	139 (94%)	7 (5%)	2 (1%)	14	58
11	W	148/160 (92%)	143 (97%)	4 (3%)	1 (1%)	26	71
11	X	148/160 (92%)	138 (93%)	6 (4%)	4 (3%)	6	45
11	Y	148/160 (92%)	140 (95%)	5 (3%)	3 (2%)	9	51
11	Z	148/160 (92%)	138 (93%)	6 (4%)	4 (3%)	6	45
11	a	148/160 (92%)	136 (92%)	8 (5%)	4 (3%)	6	45
All	All	7389/8469 (87%)	6835 (92%)	398 (5%)	156 (2%)	13	50

5 of 156 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	O	167	THR
2	O	172	VAL
5	E	475	TYR
6	F	125	LYS
6	F	207	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	b	275/728 (38%)	261 (95%)	14 (5%)	29	66
2	O	348/348 (100%)	345 (99%)	3 (1%)	84	93
3	M	183/221 (83%)	177 (97%)	6 (3%)	45	76
4	N	102/104 (98%)	100 (98%)	2 (2%)	63	85
5	A	497/515 (96%)	480 (97%)	17 (3%)	44	75
5	C	497/515 (96%)	477 (96%)	20 (4%)	38	71
5	E	497/515 (96%)	473 (95%)	24 (5%)	31	67
6	B	391/444 (88%)	374 (96%)	17 (4%)	35	70
6	D	391/444 (88%)	377 (96%)	14 (4%)	42	74
6	F	391/444 (88%)	373 (95%)	18 (5%)	33	68
7	Q	309/309 (100%)	302 (98%)	7 (2%)	58	83
8	H	87/94 (93%)	86 (99%)	1 (1%)	80	91
8	J	87/94 (93%)	83 (95%)	4 (5%)	33	68
8	L	87/94 (93%)	85 (98%)	2 (2%)	58	83
9	G	194/208 (93%)	191 (98%)	3 (2%)	72	88
9	I	194/208 (93%)	193 (100%)	1 (0%)	92	96
9	K	194/208 (93%)	192 (99%)	2 (1%)	82	92
10	P	426/439 (97%)	413 (97%)	13 (3%)	47	77
11	R	110/119 (92%)	107 (97%)	3 (3%)	52	79
11	S	110/119 (92%)	102 (93%)	8 (7%)	17	54
11	T	110/119 (92%)	106 (96%)	4 (4%)	42	74
11	U	110/119 (92%)	104 (94%)	6 (6%)	27	63
11	V	110/119 (92%)	107 (97%)	3 (3%)	52	79
11	W	110/119 (92%)	104 (94%)	6 (6%)	27	63
11	X	110/119 (92%)	107 (97%)	3 (3%)	52	79
11	Y	110/119 (92%)	110 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
11	Z	110/119 (92%)	107 (97%)	3 (3%)	52	79
11	a	110/119 (92%)	109 (99%)	1 (1%)	84	93
All	All	6250/7122 (88%)	6045 (97%)	205 (3%)	49	76

5 of 205 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
6	B	298	VAL
5	C	476	PRO
11	V	17	CYS
6	B	340	ASP
5	C	175	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 61 such sidechains are listed below:

Mol	Chain	Res	Type
6	B	446	GLN
5	C	554	HIS
11	W	151	ASN
5	C	71	GLN
5	C	151	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.