



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 1, 2016 – 12:03 AM EDT

PDB ID : 4JA9  
Title : Rat PP5 apo  
Authors : Haslbeck, V.; Helmuth, M.; Alte, F.; Popowicz, G.; Schmidt, W.; Weiwad, M.;  
Fischer, G.; Gemmecker, G.; Sattler, M.; Striggow, F.; Groll, M.; Richter, K.  
Deposited on : 2013-02-18  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	488	<div> <div>14%</div> <div>80%</div> <div>15%</div> <div>...</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MG	A	501	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3820 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein phosphatase 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	467	Total	C	N	O	S	0	0	0
			3755	2390	634	709	22			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	GLY	-	EXPRESSION TAG	UNP P53042
A	13	SER	-	EXPRESSION TAG	UNP P53042
A	14	HIS	-	EXPRESSION TAG	UNP P53042
A	15	MET	-	EXPRESSION TAG	UNP P53042

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	63	Total	O	0	0
			63	63		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein phosphatase 5



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.09 Å 50.09 Å 379.19 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.30 14.95 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (15.00-2.30) 99.8 (14.95-2.30)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.25 (at 2.29 Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.238 , 0.272 0.238 , 0.265	Depositor DCC
$R_{free}$ test set	1137 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	59.5	Xtriage
Anisotropy	0.070	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 57.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3820	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.28	0/3837	0.52	0/5176

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3755	0	3692	37	0
2	A	2	0	0	0	0
3	A	63	0	0	0	0
All	All	3820	0	3692	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:312:ILE:HG22	1:A:494:LEU:HD12	1.76	0.68

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:229:THR:OG1	1:A:342:GLN:NE2	2.29	0.65
1:A:130:LYS:O	1:A:134:MET:HB2	1.98	0.63
1:A:126:LYS:N	1:A:127:PRO:HD3	2.17	0.59
1:A:312:ILE:CG2	1:A:494:LEU:HD12	2.34	0.57
1:A:267:ILE:HG12	1:A:298:HIS:HB2	1.88	0.56
1:A:337:TRP:O	1:A:371:ARG:NH1	2.38	0.56
1:A:281:GLU:O	1:A:285:THR:HG23	2.07	0.55
1:A:415:LEU:HD21	1:A:423:ILE:HG23	1.88	0.54
1:A:89:ILE:HD12	1:A:102:ARG:HD2	1.91	0.53
1:A:191:MET:CE	1:A:194:LEU:HD23	2.40	0.52
1:A:191:MET:HA	1:A:191:MET:HE3	1.91	0.52
1:A:217:LYS:HB2	1:A:334:VAL:HG12	1.92	0.50
1:A:89:ILE:CD1	1:A:102:ARG:HD2	2.41	0.50
1:A:73:LEU:HD21	1:A:104:ALA:HB1	1.93	0.50
1:A:247:PHE:CZ	1:A:251:LEU:HD21	2.47	0.49
1:A:436:VAL:HG12	1:A:440:GLY:HA2	1.94	0.49
1:A:191:MET:HE1	1:A:194:LEU:HD23	1.95	0.48
1:A:415:LEU:HD12	1:A:441:ARG:HD2	1.95	0.48
1:A:191:MET:CE	1:A:191:MET:HA	2.43	0.48
1:A:213:LEU:HD21	1:A:331:PHE:CE1	2.47	0.48
1:A:107:MET:HE3	1:A:493:LEU:HD13	1.97	0.46
1:A:125:VAL:C	1:A:127:PRO:HD3	2.37	0.45
1:A:213:LEU:HA	1:A:216:VAL:HG12	1.98	0.45
1:A:166:ASP:N	1:A:166:ASP:OD2	2.49	0.45
1:A:51:TYR:CZ	1:A:67:ASN:HB3	2.52	0.45
1:A:225:THR:HG21	1:A:369:ILE:HB	1.98	0.44
1:A:102:ARG:HD3	1:A:118:ASP:OD1	2.17	0.43
1:A:221:CYS:HA	1:A:337:TRP:CE3	2.53	0.43
1:A:418:ASN:HB2	1:A:420:LEU:HD12	2.01	0.42
1:A:126:LYS:N	1:A:127:PRO:CD	2.81	0.42
1:A:425:ARG:O	1:A:444:THR:HA	2.18	0.42
1:A:123:VAL:HG22	1:A:132:ALA:HA	2.02	0.42
1:A:364:ASP:OD2	1:A:367:ARG:NH1	2.53	0.42
1:A:248:TYR:CE1	1:A:482:PRO:HD2	2.55	0.41
1:A:220:LEU:HD22	1:A:339:PRO:HD3	2.03	0.40
1:A:426:SER:O	1:A:427:HIS:HB3	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	463/488 (95%)	432 (93%)	26 (6%)	5 (1%)	17 18

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	233	GLU
1	A	126	LYS
1	A	132	ALA
1	A	372	ASN
1	A	164	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	407/423 (96%)	383 (94%)	24 (6%)	24 32

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	LYS
1	A	60	SER
1	A	61	ASN
1	A	94	LYS
1	A	113	ARG
1	A	116	LEU

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	A	128	ASN
1	A	130	LYS
1	A	138	GLU
1	A	170	MET
1	A	181	LEU
1	A	191	MET
1	A	229	THR
1	A	232	LYS
1	A	306	THR
1	A	371	ARG
1	A	384	LEU
1	A	396	SER
1	A	399	LYS
1	A	402	VAL
1	A	425	ARG
1	A	477	THR
1	A	479	VAL
1	A	496	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	36	ASN
1	A	61	ASN
1	A	67	ASN
1	A	310	ASN
1	A	326	GLN
1	A	342	GLN
1	A	374	GLN
1	A	450	ASN
1	A	463	HIS
1	A	465	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	467/488 (95%)	0.68	70 (14%) 3 5	44, 71, 113, 149	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	167	ILE	9.4
1	A	133	LYS	8.1
1	A	128	ASN	6.6
1	A	159	ARG	5.9
1	A	182	GLU	5.1
1	A	419	GLN	5.1
1	A	132	ALA	4.7
1	A	136	TYR	4.5
1	A	174	ASP	4.5
1	A	168	GLU	4.4
1	A	467	SER	4.0
1	A	498	MET	4.0
1	A	421	ASP	3.8
1	A	263	THR	3.7
1	A	126	LYS	3.2
1	A	88	ALA	3.2
1	A	175	GLU	3.1
1	A	50	PHE	3.1
1	A	232	LYS	3.0
1	A	464	LEU	3.0
1	A	125	VAL	3.0
1	A	285	THR	2.9
1	A	233	GLU	2.9
1	A	35	ALA	2.8
1	A	470	ARG	2.8
1	A	162	VAL	2.8
1	A	235	GLU	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	127	PRO	2.8
1	A	229	THR	2.8
1	A	131	ASP	2.7
1	A	370	GLU	2.5
1	A	130	LYS	2.5
1	A	87	ARG	2.5
1	A	134	MET	2.5
1	A	416	GLU	2.5
1	A	124	LYS	2.5
1	A	474	HIS	2.5
1	A	166	ASP	2.5
1	A	465	GLN	2.5
1	A	456	GLY	2.4
1	A	169	SER	2.4
1	A	137	GLN	2.4
1	A	222	LYS	2.4
1	A	460	SER	2.4
1	A	31	LEU	2.3
1	A	231	LEU	2.3
1	A	358	GLU	2.2
1	A	347	LYS	2.2
1	A	164	SER	2.2
1	A	411	THR	2.2
1	A	145	GLN	2.2
1	A	47	ALA	2.2
1	A	496	LEU	2.2
1	A	248	TYR	2.2
1	A	183	ASP	2.2
1	A	237	ILE	2.1
1	A	206	ARG	2.1
1	A	262	GLU	2.1
1	A	337	TRP	2.1
1	A	85	ALA	2.1
1	A	100	TYR	2.1
1	A	165	LEU	2.1
1	A	230	THR	2.1
1	A	34	GLN	2.0
1	A	218	GLU	2.0
1	A	372	ASN	2.0
1	A	234	THR	2.0
1	A	40	LYS	2.0
1	A	335	PHE	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	184	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	A	501	1/1	0.90	0.19	2.59	55,55,55,55	0
2	MG	A	502	1/1	0.89	0.15	0.68	57,57,57,57	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.