



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:43 PM GMT

PDB ID : 4JAV  
Title : Structural basis of a rationally rewired protein-protein interface (HK853wt and RR468mutant V13P, L14I, I17M and N21V)  
Authors : Podgornaia, A.I.; Casino, P.; Marina, A.; Laub, M.T.  
Deposited on : 2013-02-19  
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1114 (3.14-3.06)
Clashscore	102246	1222 (3.14-3.06)
Ramachandran outliers	100387	1174 (3.14-3.06)
Sidechain outliers	100360	1174 (3.14-3.06)
RSRZ outliers	91569	1119 (3.14-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	<div> <div></div> <div>80%14%• 5%</div> </div>
1	B	258	<div> <div>%</div> <div>79%14%• 5%</div> </div>
2	C	122	<div> <div></div> <div>88%11%••</div> </div>
2	D	122	<div> <div>10%</div> <div>86%11%••</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	B	503	-	-	X	-

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5944 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histidine kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	246	Total	C	N	O	S	0	0	0
			1968	1253	333	379	3			
1	B	244	Total	C	N	O	S	0	1	0
			1953	1243	333	374	3			

- Molecule 2 is a protein called Response regulator.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
2	C	121	Total	Be	C	F	N	O	S	0	0	0
			965	1	620	3	155	181	5			
2	D	120	Total	Be	C	F	N	O	S	0	2	0
			961	1	619	3	154	179	5			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	13	PRO	VAL	ENGINEERED MUTATION	UNP Q9WYT9
C	14	ILE	LEU	ENGINEERED MUTATION	UNP Q9WYT9
C	17	MET	ILE	ENGINEERED MUTATION	UNP Q9WYT9
C	21	VAL	ASN	ENGINEERED MUTATION	UNP Q9WYT9
D	13	PRO	VAL	ENGINEERED MUTATION	UNP Q9WYT9
D	14	ILE	LEU	ENGINEERED MUTATION	UNP Q9WYT9
D	17	MET	ILE	ENGINEERED MUTATION	UNP Q9WYT9
D	21	VAL	ASN	ENGINEERED MUTATION	UNP Q9WYT9

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	A	1	Total	Mg	0	0
			1	1		
4	D	1	Total	Mg	0	0
			1	1		
4	C	1	Total	Mg	0	0
			1	1		

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
5	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
5	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Cl	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	8	Total	O	0	0
			8	8		
7	B	1	Total	O	0	0
			1	1		
7	C	4	Total	O	0	0
			4	4		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histidine kinase




- Molecule 1: Histidine kinase



- Molecule 2: Response regulator



- Molecule 2: Response regulator



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	119.32Å 143.93Å 138.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.55 – 3.10 44.51 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.55-3.10) 99.8 (44.51-3.10)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.51 (at 3.12Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.202 , 0.253 0.202 , 0.254	Depositor DCC
$R_{free}$ test set	1129 reflections (5.41%)	DCC
Wilson B-factor (Å <sup>2</sup> )	71.0	Xtriage
Anisotropy	0.065	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 44.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 22018 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5944	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BFD, SO4, ADP, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.27	0/2000	0.46	0/2705
1	B	0.29	0/1988	0.59	5/2688 (0.2%)
2	C	0.27	0/966	0.48	0/1294
2	D	0.28	0/968	0.46	0/1298
All	All	0.28	0/5922	0.51	5/7985 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	412	ASN	N-CA-C	-10.35	83.04	111.00
1	B	412	ASN	CB-CA-C	7.96	126.32	110.40
1	B	359	CYS	C-N-CD	-6.73	105.80	120.60
1	B	413	GLY	N-CA-C	-5.44	99.51	113.10
1	B	360	PRO	N-CA-CB	5.18	109.51	103.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	359	CYS	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1968	0	1989	21	0
1	B	1953	0	1976	36	0
2	C	965	0	1010	7	0
2	D	961	0	1014	6	0
3	A	10	0	0	0	0
3	B	15	0	0	2	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	27	0	12	0	0
5	B	27	0	12	0	0
6	B	1	0	0	0	0
7	A	8	0	0	0	0
7	B	1	0	0	0	0
7	C	4	0	0	0	0
All	All	5944	0	6013	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:387:LYS:CE	1:B:435:LEU:HD11	1.69	1.21
1:B:387:LYS:HE3	1:B:435:LEU:HD11	1.43	0.99
1:B:387:LYS:NZ	1:B:435:LEU:HD21	1.81	0.95
1:B:387:LYS:HE2	1:B:435:LEU:HD11	1.47	0.94
1:B:387:LYS:HZ3	1:B:435:LEU:HD21	1.35	0.88
1:B:441:GLY:HA2	3:B:503:SO4:O1	1.74	0.86
2:C:21:VAL:HG12	2:C:113:ILE:HD11	1.60	0.82
1:B:265:PRO:HA	1:B:298:GLN:HE21	1.51	0.75
1:B:387:LYS:NZ	1:B:435:LEU:CD2	2.54	0.71
1:B:387:LYS:CE	1:B:435:LEU:CD1	2.61	0.67
1:B:387:LYS:HZ1	1:B:435:LEU:HG	1.59	0.67

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:387:LYS:HZ1	1:B:435:LEU:CG	2.08	0.66
1:A:386:LYS:NZ	1:A:467:GLY:O	2.24	0.66
1:B:387:LYS:CE	1:B:435:LEU:HD21	2.29	0.63
1:B:435:LEU:C	1:B:435:LEU:HD12	2.19	0.63
1:B:387:LYS:HE2	1:B:435:LEU:CD1	2.24	0.61
1:B:441:GLY:CA	3:B:503:SO4:O1	2.46	0.60
1:B:410:GLU:HG3	1:B:471:ARG:HG2	1.83	0.59
2:C:25:GLU:HG3	2:C:113:ILE:HD12	1.84	0.59
1:A:404:GLY:HA2	1:A:478:LYS:HG2	1.84	0.57
1:A:265:PRO:HG3	1:A:301:HIS:HD2	1.69	0.57
1:B:265:PRO:HG3	1:B:301:HIS:CD2	2.40	0.57
1:B:265:PRO:HG3	1:B:301:HIS:HD2	1.70	0.57
2:D:68:LEU:HD23	2:D:74:TRP:HB3	1.86	0.56
1:A:266:LEU:HD13	1:B:302:LEU:HD22	1.86	0.56
1:A:410:GLU:HG3	1:A:471:ARG:HG2	1.88	0.56
1:A:418:ASP:HA	1:A:421:LYS:HD3	1.90	0.54
2:D:101:LYS:HD2	2:D:115:GLU:HG2	1.89	0.53
1:A:265:PRO:HA	1:A:298:GLN:HE21	1.75	0.52
1:A:310:LEU:HD22	1:B:263:ARG:HH11	1.76	0.51
1:B:327:VAL:HG11	1:B:367:PRO:HB3	1.92	0.50
1:B:435:LEU:HD12	1:B:435:LEU:O	2.12	0.50
1:B:435:LEU:CG	1:B:435:LEU:O	2.60	0.50
1:B:322:ILE:HG12	1:B:456:HIS:CE1	2.47	0.50
2:C:3:LYS:HG2	2:C:120:LEU:HD22	1.95	0.49
1:B:283:LEU:HD11	1:B:288:LEU:HD13	1.95	0.49
1:B:329:LEU:HD13	1:B:405:VAL:HG11	1.95	0.48
1:A:277:TYR:HA	1:A:280:LEU:HD23	1.94	0.48
1:B:387:LYS:HE2	1:B:435:LEU:HD21	1.95	0.48
1:B:387:LYS:NZ	1:B:435:LEU:CG	2.74	0.47
1:A:428:PHE:CE1	1:A:444:LEU:HD11	2.50	0.47
1:A:397:ILE:HB	1:A:408:ILE:HB	1.97	0.47
2:C:82:LEU:HD22	2:C:105:LYS:HG3	1.98	0.46
1:B:435:LEU:O	1:B:435:LEU:HG	2.17	0.45
1:A:404:GLY:HA3	1:A:477:PRO:HA	1.98	0.45
1:B:243:ARG:O	1:B:247:ILE:HD12	2.17	0.45
1:A:276:ILE:HD11	1:A:291:PHE:HB3	1.99	0.45
1:A:276:ILE:HA	1:A:283:LEU:HD11	1.99	0.45
1:B:435:LEU:HB3	1:B:439:VAL:HG22	1.99	0.45
1:A:356:ASN:OD1	1:A:399:ASP:HA	2.17	0.45
1:A:265:PRO:HG3	1:A:301:HIS:CD2	2.50	0.44
1:A:264:THR:HB	1:A:265:PRO:HD3	2.00	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:373:VAL:HG13	1:A:449:THR:HG23	2.00	0.43
2:C:4:LYS:HG2	2:C:28:GLU:HB2	2.00	0.43
2:D:89:GLU:H	2:D:89:GLU:CD	2.22	0.43
1:B:329:LEU:O	1:B:333:VAL:HG22	2.18	0.43
1:A:329:LEU:HD11	1:A:476:ILE:HD11	2.01	0.43
2:D:113:ILE:HD13	2:D:113:ILE:HA	1.87	0.42
1:B:431:VAL:HG13	1:B:434:SER:H	1.84	0.42
2:D:84:ALA:HB2	2:D:105:LYS:HD2	2.01	0.42
1:A:416:ILE:HA	1:A:417:PRO:HD2	1.90	0.42
1:A:329:LEU:HD13	1:A:398:LEU:HD11	2.02	0.41
2:C:21:VAL:CG1	2:C:113:ILE:HD11	2.38	0.41
1:B:365:ILE:HG22	1:B:456:HIS:HD2	1.86	0.40
2:D:5:VAL:HG13	2:D:49:LEU:HB3	2.04	0.40
2:C:77:ILE:HA	2:C:78:PRO:HD3	1.97	0.40
1:B:264:THR:HB	1:B:265:PRO:HD3	2.03	0.40
1:B:251:LYS:O	1:B:255:ILE:HG12	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/258 (95%)	237 (97%)	7 (3%)	0	100	100
1	B	243/258 (94%)	233 (96%)	8 (3%)	2 (1%)	24	63
2	C	118/122 (97%)	117 (99%)	1 (1%)	0	100	100
2	D	119/122 (98%)	116 (98%)	3 (2%)	0	100	100
All	All	724/760 (95%)	703 (97%)	19 (3%)	2 (0%)	46	80

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	360	PRO
1	B	359	CYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	221/232 (95%)	213 (96%)	8 (4%)	42	77
1	B	219/232 (94%)	207 (94%)	12 (6%)	27	63
2	C	108/109 (99%)	104 (96%)	4 (4%)	41	76
2	D	109/109 (100%)	103 (94%)	6 (6%)	27	63
All	All	657/682 (96%)	627 (95%)	30 (5%)	33	70

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	247	ILE
1	A	248	ASP
1	A	280	LEU
1	A	329	LEU
1	A	402	ASP
1	A	422	ASP
1	A	436	THR
1	A	450	LYS
1	B	236	THR
1	B	248	ASP
1	B	279	SER
1	B	284	ASP
1	B	292	LEU
1	B	322	ILE
1	B	329	LEU
1	B	331	ASP
1	B	359	CYS
1	B	391	ASP
1	B	432	ASP
1	B	435	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	C	7	LEU
2	C	42	LEU
2	C	82	LEU
2	C	104	ARG
2	D	7	LEU
2	D	42	LEU
2	D	82	LEU
2	D	107	PHE
2	D	108	SER
2	D	113	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	278	ASN
1	A	298	GLN
1	A	300	ASN
1	A	301	HIS
1	A	323	ASN
1	B	298	GLN
1	B	300	ASN
1	B	301	HIS
1	B	356	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BFD	C	53	2,4	8,11,12	1.04	0	4,15,17	2.40	2 (50%)
2	BFD	D	53	2,4	8,11,12	1.05	0	4,15,17	2.42	3 (75%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BFD	C	53	2,4	-	0/4/11/13	0/0/0/0
2	BFD	D	53	2,4	-	0/4/11/13	0/0/0/0

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	53	BFD	OD2-CG-CB	-2.95	117.91	124.69
2	D	53	BFD	OD2-CG-CB	-2.89	118.05	124.69
2	D	53	BFD	O-C-CA	-2.12	119.97	125.49
2	D	53	BFD	OD1-CG-CB	3.24	119.27	111.50
2	C	53	BFD	OD1-CG-CB	3.24	119.28	111.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 5 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	501	-	4,4,4	0.36	0	6,6,6	0.13	0
3	SO4	A	502	4	4,4,4	0.37	0	6,6,6	0.14	0
5	ADP	A	504	4	22,29,29	1.08	2 (9%)	27,45,45	1.81	5 (18%)
3	SO4	B	501	-	4,4,4	0.33	0	6,6,6	0.07	0
3	SO4	B	502	-	4,4,4	0.33	0	6,6,6	0.10	0
3	SO4	B	503	-	4,4,4	0.33	0	6,6,6	0.08	0
5	ADP	B	505	4	22,29,29	1.06	1 (4%)	27,45,45	2.00	5 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	501	-	-	0/0/0/0	0/0/0/0
3	SO4	A	502	4	-	0/0/0/0	0/0/0/0
5	ADP	A	504	4	-	0/12/32/32	0/3/3/3
3	SO4	B	501	-	-	0/0/0/0	0/0/0/0
3	SO4	B	502	-	-	0/0/0/0	0/0/0/0
3	SO4	B	503	-	-	0/0/0/0	0/0/0/0
5	ADP	B	505	4	-	0/12/32/32	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	504	ADP	O4'-C1'	2.17	1.44	1.41
5	A	504	ADP	C5-C4	3.22	1.47	1.40
5	B	505	ADP	C5-C4	3.23	1.47	1.40

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	505	ADP	N3-C2-N1	-7.06	123.49	128.89
5	A	504	ADP	N3-C2-N1	-6.87	123.64	128.89
5	B	505	ADP	C2'-C1'-N9	-4.20	107.87	114.29
5	B	505	ADP	PA-O3A-PB	-3.32	121.54	132.67
5	B	505	ADP	C4-C5-N7	-3.18	106.55	109.48
5	A	504	ADP	C4-C5-N7	-3.03	106.69	109.48

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	504	ADP	C2'-C1'-N9	-2.61	110.30	114.29
5	A	504	ADP	PA-O3A-PB	-2.34	124.82	132.67
5	A	504	ADP	O4'-C1'-N9	2.02	112.33	108.10
5	B	505	ADP	O3B-PB-O2B	2.14	115.54	107.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	503	SO4	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	246/258 (95%)	-0.10	0 100 100	44, 63, 87, 99	0
1	B	244/258 (94%)	0.17	3 (1%) 81 64	47, 75, 110, 160	0
2	C	120/122 (98%)	0.01	0 100 100	45, 58, 81, 91	0
2	D	119/122 (97%)	0.65	12 (10%) 9 3	49, 82, 120, 128	0
All	All	729/760 (95%)	0.13	15 (2%) 67 44	44, 68, 106, 160	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	69	GLN	4.4
2	D	97	LEU	3.5
1	B	436	THR	3.3
2	D	68	LEU	2.9
2	D	121	ASN	2.6
2	D	95	LEU	2.6
2	D	86	GLY	2.6
1	B	403	GLY	2.6
2	D	65	LEU	2.5
2	D	96	SER	2.5
1	B	477	PRO	2.4
2	D	76	ARG	2.3
2	D	75	LYS	2.2
2	D	87	GLY	2.1
2	D	70	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	BFD	C	53	12/13	0.98	0.17	-	47,49,54,54	0
2	BFD	D	53	12/13	0.95	0.17	-	55,58,64,65	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	B	502	5/5	0.91	0.24	0.06	141,142,143,143	0
5	ADP	A	504	27/27	0.98	0.21	-0.08	48,52,53,54	0
4	MG	C	201	1/1	0.97	0.18	-0.09	40,40,40,40	0
3	SO4	B	503	5/5	0.83	0.21	-0.41	161,162,163,165	0
5	ADP	B	505	27/27	0.98	0.20	-0.94	70,73,75,76	0
3	SO4	A	502	5/5	0.89	0.18	-1.01	110,112,112,113	0
3	SO4	A	501	5/5	0.97	0.13	-2.68	79,81,82,82	0
4	MG	D	201	1/1	0.95	0.15	-	45,45,45,45	0
3	SO4	B	501	5/5	0.96	0.13	-	83,84,86,86	0
4	MG	B	504	1/1	0.92	0.16	-	60,60,60,60	0
6	CL	B	506	1/1	0.87	0.28	-	79,79,79,79	0
4	MG	A	503	1/1	0.94	0.29	-	40,40,40,40	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.