



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 3, 2016 – 01:23 PM EDT

PDB ID : 5JCM
Title : Structure and catalytic mechanism of monodehydroascorbate reductase, MD-HAR, from *Oryza sativa* L. japonica
Authors : Park, A.K.; Kim, H.W.
Deposited on : 2016-04-15
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	<div> <div>3%</div> <div> <div></div> <div>84%</div> <div>10%</div> <div>• 5%</div> </div> </div>
1	B	451	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>9%</div> <div>• 6%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ISD	A	502	-	-	-	X
4	ISD	B	502	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7283 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Os09g0567300 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	429	Total	C	N	O	S	0	0	0
			3256	2099	531	620	6			
1	B	425	Total	C	N	O	S	0	0	0
			3223	2077	526	614	6			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-15	HIS	-	expression tag	UNP Q652L6
A	-14	HIS	-	expression tag	UNP Q652L6
A	-13	HIS	-	expression tag	UNP Q652L6
A	-12	HIS	-	expression tag	UNP Q652L6
A	-11	HIS	-	expression tag	UNP Q652L6
A	-10	HIS	-	expression tag	UNP Q652L6
A	-9	ALA	-	expression tag	UNP Q652L6
A	-8	SER	-	expression tag	UNP Q652L6
A	-7	GLU	-	expression tag	UNP Q652L6
A	-6	ASN	-	expression tag	UNP Q652L6
A	-5	LEU	-	expression tag	UNP Q652L6
A	-4	TYR	-	expression tag	UNP Q652L6
A	-3	PHE	-	expression tag	UNP Q652L6
A	-2	GLN	-	expression tag	UNP Q652L6
A	-1	GLY	-	expression tag	UNP Q652L6
A	0	ALA	-	expression tag	UNP Q652L6
A	1	MET	-	expression tag	UNP Q652L6
A	2	ALA	-	expression tag	UNP Q652L6
A	3	SER	-	expression tag	UNP Q652L6
A	349	PHE	TYR	engineered mutation	UNP Q652L6
B	-15	HIS	-	expression tag	UNP Q652L6
B	-14	HIS	-	expression tag	UNP Q652L6
B	-13	HIS	-	expression tag	UNP Q652L6
B	-12	HIS	-	expression tag	UNP Q652L6
B	-11	HIS	-	expression tag	UNP Q652L6

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-10	HIS	-	expression tag	UNP Q652L6
B	-9	ALA	-	expression tag	UNP Q652L6
B	-8	SER	-	expression tag	UNP Q652L6
B	-7	GLU	-	expression tag	UNP Q652L6
B	-6	ASN	-	expression tag	UNP Q652L6
B	-5	LEU	-	expression tag	UNP Q652L6
B	-4	TYR	-	expression tag	UNP Q652L6
B	-3	PHE	-	expression tag	UNP Q652L6
B	-2	GLN	-	expression tag	UNP Q652L6
B	-1	GLY	-	expression tag	UNP Q652L6
B	0	ALA	-	expression tag	UNP Q652L6
B	1	MET	-	expression tag	UNP Q652L6
B	2	ALA	-	expression tag	UNP Q652L6
B	3	SER	-	expression tag	UNP Q652L6
B	349	PHE	TYR	engineered mutation	UNP Q652L6

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0
			53	27	9	15	2	
2	B	1	Total	C	N	O	P	0
			53	27	9	15	2	

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 44	C 21	N 7	O 14	P 2	0	0
3	B	1	Total 44	C 21	N 7	O 14	P 2	0	0

- Molecule 4 is (5R)-5-[(1R)-1,2-DIHYDROXYETHYL]-3,4-DIHYDROXYFURAN-2(5H)-ONE (three-letter code: ISD) (formula: C₆H₈O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			12	6	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			12	6	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	290	Total	O	0	0
			290	290		
5	B	296	Total	O	0	0
			296	296		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Os09g0567300 protein



• Molecule 1: Os09g0567300 protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.39Å 85.14Å 131.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.90 39.20 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.00-1.90) 99.3 (39.20-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.68 (at 1.91Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.174 , 0.224 0.186 , 0.233	Depositor DCC
R_{free} test set	3409 reflections (5.16%)	DCC
Wilson B-factor (Å ²)	31.9	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 60.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7283	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, FAD, ISD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.03	2/3325 (0.1%)	1.10	14/4497 (0.3%)
1	B	0.98	1/3291 (0.0%)	1.16	17/4450 (0.4%)
All	All	1.00	3/6616 (0.0%)	1.13	31/8947 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	76	GLU	CD-OE1	6.07	1.32	1.25
1	B	41	GLU	CD-OE2	5.91	1.32	1.25
1	A	213	GLU	CD-OE2	5.48	1.31	1.25

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	63	ARG	NE-CZ-NH2	-25.90	107.35	120.30
1	B	63	ARG	NE-CZ-NH1	22.83	131.72	120.30
1	A	63	ARG	NE-CZ-NH2	-17.96	111.32	120.30
1	A	63	ARG	NE-CZ-NH1	16.62	128.61	120.30
1	B	63	ARG	CD-NE-CZ	10.52	138.33	123.60
1	A	351	ARG	NE-CZ-NH2	8.19	124.39	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	351	ARG	NE-CZ-NH1	-7.85	116.38	120.30
1	B	77	ARG	NE-CZ-NH1	-7.63	116.49	120.30
1	B	147	ARG	NE-CZ-NH1	7.11	123.85	120.30
1	B	312	ARG	NE-CZ-NH1	6.93	123.77	120.30
1	A	48	ARG	NE-CZ-NH1	6.55	123.57	120.30
1	A	163	LYS	CD-CE-NZ	6.43	126.48	111.70
1	A	63	ARG	CD-NE-CZ	6.21	132.30	123.60
1	B	372	ASP	C-N-CA	6.13	137.03	121.70
1	A	372	ASP	C-N-CA	6.08	136.90	121.70
1	A	127	VAL	CB-CA-C	-6.07	99.86	111.40
1	A	312	ARG	NE-CZ-NH1	6.05	123.32	120.30
1	B	334	GLU	C-N-CA	5.70	135.95	121.70
1	B	334	GLU	CA-C-N	5.66	129.66	117.20
1	B	334	GLU	N-CA-C	5.61	126.16	111.00
1	B	311	ARG	NE-CZ-NH2	-5.47	117.56	120.30
1	B	77	ARG	NE-CZ-NH2	5.41	123.01	120.30
1	B	235	ASP	CB-CG-OD1	5.30	123.07	118.30
1	B	372	ASP	O-C-N	-5.23	114.33	122.70
1	B	48	ARG	NE-CZ-NH1	5.22	122.91	120.30
1	A	312	ARG	NE-CZ-NH2	-5.21	117.70	120.30
1	B	372	ASP	CA-C-N	5.13	128.50	117.20
1	A	63	ARG	CG-CD-NE	-5.12	101.04	111.80
1	B	254	ASP	CB-CG-OD1	-5.11	113.70	118.30
1	A	144	LEU	CB-CG-CD1	5.07	119.61	111.00
1	A	213	GLU	OE1-CD-OE2	5.02	129.32	123.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	337	GLU	Peptide
1	A	376	THR	Peptide
1	B	334	GLU	Peptide
1	B	63	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3256	0	3257	25	1
1	B	3223	0	3217	17	1
2	A	53	0	31	0	0
2	B	53	0	31	0	0
3	A	44	0	26	1	0
3	B	44	0	25	0	0
4	A	12	0	6	0	1
4	B	12	0	6	1	1
5	A	290	0	0	5	1
5	B	296	0	0	7	1
All	All	7283	0	6599	42	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:179:LEU:HD23	1:A:257:VAL:HG11	1.60	0.82
1:B:71:VAL:CG1	1:B:77:ARG:HG2	2.13	0.79
1:A:233:ASP:OD2	5:A:601:HOH:O	2.01	0.78
1:A:83:TYR:OH	5:A:602:HOH:O	2.05	0.72
1:B:71:VAL:HG13	1:B:77:ARG:HG2	1.72	0.72
1:B:77:ARG:NH1	5:B:603:HOH:O	2.30	0.63
1:B:47:GLU:HG3	5:B:777:HOH:O	1.99	0.63
1:A:105:THR:HG22	1:A:115:THR:HG22	1.82	0.61
1:B:71:VAL:HG23	5:B:777:HOH:O	2.02	0.59
1:A:263:ARG:NE	5:A:608:HOH:O	2.36	0.57
1:A:127:VAL:HG22	1:A:147:ARG:HB2	1.87	0.54
1:A:377:SER:HB3	1:A:378:ALA:CA	2.40	0.52
1:A:107:THR:OG1	5:A:603:HOH:O	2.19	0.51
1:B:19:GLY:HA3	1:B:69:VAL:HG21	1.93	0.51
1:A:377:SER:HB3	1:A:378:ALA:HA	1.92	0.51
1:A:337:GLU:HB3	1:A:338:SER:HA	1.93	0.49
1:A:377:SER:HB3	1:A:378:ALA:HB2	1.94	0.49
1:A:200:MET:HB2	1:A:204:PHE:CD2	2.48	0.48
1:A:55:TYR:O	1:A:63:ARG:NH2	2.42	0.48
1:B:222:LYS:HE2	5:B:646:HOH:O	2.13	0.48
1:B:4:GLU:N	5:B:609:HOH:O	2.46	0.48
1:A:342:TYR:CE2	1:A:344:TYR:HB2	2.49	0.48
1:B:264:PRO:HG3	1:B:312:ARG:NH1	2.29	0.47
1:B:331:LYS:NZ	5:B:611:HOH:O	2.48	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:200:MET:N	1:B:201:PRO:CD	2.79	0.46
1:A:264:PRO:HG2	1:A:278:GLY:O	2.17	0.45
1:A:179:LEU:CD2	1:A:257:VAL:HG11	2.40	0.45
1:B:342:TYR:CE2	1:B:344:TYR:HB2	2.52	0.45
1:A:19:GLY:HA3	1:A:69:VAL:HG21	1.98	0.44
1:B:110:VAL:O	1:B:110:VAL:HG12	2.18	0.44
1:A:180:SER:OG	1:A:192:MET:HG2	2.18	0.43
1:B:145:TYR:CE1	1:B:258:VAL:HB	2.54	0.43
1:A:377:SER:HB3	1:A:378:ALA:CB	2.49	0.43
1:A:213:GLU:OE1	1:B:225:LYS:HB2	2.19	0.43
4:B:502:ISD:H132	5:B:777:HOH:O	2.18	0.43
1:A:174:TYR:HB3	3:A:501:NAD:C4N	2.48	0.43
1:B:331:LYS:O	1:B:335:SER:HB3	2.18	0.43
1:B:68:HIS:O	1:B:71:VAL:HG22	2.19	0.43
1:A:4:GLU:CB	1:A:113:THR:HG23	2.50	0.42
1:A:15:GLY:HA2	5:A:814:HOH:O	2.19	0.41
1:A:103:SER:O	1:A:105:THR:HG23	2.21	0.41
1:A:213:GLU:HG2	1:A:223:ILE:HD12	2.03	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:669:HOH:O	5:B:601:HOH:O[3_454]	1.66	0.54
1:B:133:PHE:O	4:A:502:ISD:O14[4_455]	1.99	0.21
1:A:133:PHE:O	4:B:502:ISD:O14[3_454]	2.14	0.06

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	425/451 (94%)	404 (95%)	15 (4%)	6 (1%)	14 4

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	419/451 (93%)	404 (96%)	13 (3%)	2 (0%)	34	21
All	All	844/902 (94%)	808 (96%)	28 (3%)	8 (1%)	21	9

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	428	GLY
1	B	373	SER
1	B	433	SER
1	A	373	SER
1	A	425	LYS
1	A	337	GLU
1	A	426	LYS
1	A	377	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	336/353 (95%)	329 (98%)	7 (2%)	61	55
1	B	332/353 (94%)	320 (96%)	12 (4%)	42	30
All	All	668/706 (95%)	649 (97%)	19 (3%)	51	41

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	VAL
1	A	144	LEU
1	A	196	GLU
1	A	270	LYS
1	A	373	SER
1	A	423	GLU
1	A	429	LEU
1	B	78	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	84	SER
1	B	85	GLU
1	B	86	LYS
1	B	144	LEU
1	B	184	LYS
1	B	202	ARG
1	B	305	LYS
1	B	338	SER
1	B	379	LYS
1	B	429	LEU
1	B	434	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	315	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	A	500	-	52,58,58	2.04	13 (25%)	52,89,89	3.39	17 (32%)
3	NAD	A	501	-	42,48,48	2.03	10 (23%)	46,73,73	3.43	6 (13%)
4	ISD	A	502	-	12,12,12	1.37	2 (16%)	16,17,17	2.82	9 (56%)
2	FAD	B	500	-	52,58,58	1.84	10 (19%)	52,89,89	2.90	13 (25%)
3	NAD	B	501	-	42,48,48	2.01	10 (23%)	46,73,73	2.24	10 (21%)
4	ISD	B	502	-	12,12,12	2.01	1 (8%)	16,17,17	2.62	6 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	500	-	-	0/30/50/50	0/6/6/6
3	NAD	A	501	-	-	0/22/62/62	0/5/5/5
4	ISD	A	502	-	-	0/6/22/22	0/1/1/1
2	FAD	B	500	-	-	0/30/50/50	0/6/6/6
3	NAD	B	501	-	-	0/22/62/62	0/5/5/5
4	ISD	B	502	-	-	0/6/22/22	0/1/1/1

All (46) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	NAD	C2D-C1D	-5.66	1.44	1.53
3	B	501	NAD	C3N-C7N	-5.42	1.42	1.50
3	A	501	NAD	C3N-C7N	-4.56	1.43	1.50
2	A	500	FAD	C8M-C8	-4.04	1.43	1.51
2	B	500	FAD	C10-N10	-3.75	1.34	1.39
2	B	500	FAD	C8M-C8	-3.51	1.44	1.51
2	B	500	FAD	C10-N1	-3.41	1.29	1.35
3	B	501	NAD	O3B-C3B	-3.03	1.35	1.43
2	B	500	FAD	C7M-C7	-2.94	1.45	1.51
2	A	500	FAD	C3B-C4B	-2.91	1.45	1.53
2	A	500	FAD	C7M-C7	-2.77	1.45	1.51
3	B	501	NAD	PN-O2N	-2.75	1.43	1.55
2	B	500	FAD	C4-C4X	-2.68	1.35	1.41
2	A	500	FAD	P-O1P	-2.67	1.41	1.51
3	B	501	NAD	C2D-C1D	-2.66	1.49	1.53
3	A	501	NAD	C5A-C4A	-2.47	1.34	1.40
3	B	501	NAD	C2D-C3D	-2.36	1.47	1.53
2	A	500	FAD	C2-N3	-2.33	1.33	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	NAD	PN-O2N	-2.30	1.45	1.55
2	B	500	FAD	C4X-C10	-2.26	1.36	1.40
2	A	500	FAD	C6-C5X	-2.00	1.38	1.41
2	A	500	FAD	O2'-C2'	2.01	1.47	1.43
3	B	501	NAD	C7N-N7N	2.08	1.37	1.33
4	A	502	ISD	O11-C9	2.09	1.48	1.43
3	A	501	NAD	O3D-C3D	2.28	1.48	1.43
3	A	501	NAD	C2A-N1A	2.35	1.38	1.33
2	B	500	FAD	C2A-N1A	2.94	1.39	1.33
3	B	501	NAD	O4D-C1D	2.97	1.45	1.41
2	B	500	FAD	C4-N3	2.98	1.38	1.33
3	A	501	NAD	O2D-C2D	3.39	1.50	1.43
4	A	502	ISD	O1-C5	3.40	1.40	1.35
2	A	500	FAD	C6-C7	3.46	1.47	1.37
2	A	500	FAD	C2A-N1A	3.62	1.40	1.33
3	A	501	NAD	C2A-N3A	3.78	1.38	1.32
3	B	501	NAD	C2A-N1A	3.78	1.41	1.33
3	B	501	NAD	C4N-C3N	4.01	1.45	1.39
2	A	500	FAD	C4X-N5	4.18	1.39	1.33
2	A	500	FAD	O4B-C4B	4.34	1.55	1.45
3	A	501	NAD	C2N-N1N	4.42	1.41	1.35
3	A	501	NAD	O2B-C2B	4.61	1.53	1.43
2	B	500	FAD	C4X-N5	4.88	1.40	1.33
3	B	501	NAD	C2A-N3A	5.00	1.41	1.32
2	A	500	FAD	C2A-N3A	5.32	1.41	1.32
2	B	500	FAD	C2A-N3A	5.64	1.42	1.32
2	A	500	FAD	O4B-C1B	5.78	1.49	1.41
4	B	502	ISD	O1-C5	6.47	1.45	1.35

All (61) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	NAD	N3A-C2A-N1A	-16.94	115.56	128.87
2	B	500	FAD	N3A-C2A-N1A	-13.39	118.36	128.87
3	A	501	NAD	C1B-N9A-C4A	-13.16	112.12	126.81
2	A	500	FAD	N3A-C2A-N1A	-12.27	119.23	128.87
3	B	501	NAD	N3A-C2A-N1A	-11.24	120.04	128.87
2	A	500	FAD	C1B-N9A-C4A	-10.45	115.15	126.81
2	B	500	FAD	C1B-N9A-C4A	-6.77	119.25	126.81
4	B	502	ISD	C9-C2-C3	-6.75	104.44	114.17
2	A	500	FAD	N3-C2-N1	-6.61	116.56	127.69
2	B	500	FAD	N3-C2-N1	-6.25	117.17	127.69

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	502	ISD	C13-C9-C2	-5.35	102.90	111.95
2	A	500	FAD	C4B-O4B-C1B	-4.97	104.38	109.64
4	A	502	ISD	C9-C2-C3	-4.87	107.14	114.17
2	A	500	FAD	O4B-C1B-N9A	-4.86	98.92	108.11
4	B	502	ISD	C2-O1-C5	-4.38	105.02	109.29
2	B	500	FAD	C9A-C5X-N5	-3.90	115.83	122.18
4	A	502	ISD	O14-C13-C9	-3.76	102.74	111.07
2	A	500	FAD	C4X-C4-N3	-3.73	118.65	123.52
3	A	501	NAD	O3B-C3B-C2B	-3.48	100.59	111.86
2	B	500	FAD	O4B-C1B-N9A	-3.26	101.95	108.11
4	B	502	ISD	O6-C5-C4	-3.15	125.45	129.30
3	A	501	NAD	O4D-C1D-N1N	-3.14	104.71	108.10
3	B	501	NAD	C4B-O4B-C1B	-3.12	106.33	109.64
2	B	500	FAD	C4B-O4B-C1B	-3.11	106.35	109.64
2	B	500	FAD	C4X-C4-N3	-2.96	119.65	123.52
4	A	502	ISD	O1-C2-C9	-2.93	104.69	109.69
3	A	501	NAD	C2N-C3N-C4N	-2.81	115.08	118.27
4	A	502	ISD	O1-C5-C4	-2.72	107.61	109.94
2	A	500	FAD	O2A-PA-O5B	-2.71	95.32	108.24
2	A	500	FAD	C9A-C5X-N5	-2.69	117.81	122.18
2	B	500	FAD	C9-C9A-C5X	-2.52	115.13	119.65
3	B	501	NAD	O4B-C1B-N9A	-2.51	103.36	108.11
3	B	501	NAD	C2N-C3N-C4N	-2.17	115.80	118.27
3	B	501	NAD	C4D-O4D-C1D	-2.16	107.35	109.64
2	A	500	FAD	C9-C9A-C5X	-2.15	115.78	119.65
3	B	501	NAD	C2B-C3B-C4B	-2.11	98.31	102.64
4	A	502	ISD	C2-O1-C5	-2.09	107.25	109.29
3	A	501	NAD	C2B-C1B-N9A	2.19	119.31	113.47
2	A	500	FAD	C4-C4X-N5	2.23	121.41	118.70
4	B	502	ISD	O1-C2-C3	2.23	105.90	104.13
2	A	500	FAD	O2A-PA-O3P	2.34	115.30	105.27
3	B	501	NAD	O7N-C7N-C3N	2.55	122.43	119.60
2	B	500	FAD	O2'-C2'-C1'	2.68	116.54	109.93
3	B	501	NAD	O3B-C3B-C2B	2.89	121.21	111.86
2	A	500	FAD	C6-C5X-C9A	2.91	122.32	119.11
3	B	501	NAD	O2B-C2B-C1B	2.91	120.73	111.61
4	A	502	ISD	O1-C2-C3	2.96	106.47	104.13
3	B	501	NAD	C3N-C2N-N1N	3.04	123.81	120.34
4	B	502	ISD	O1-C5-O6	3.04	124.72	121.27
2	B	500	FAD	C6-C5X-C9A	3.25	122.69	119.11
4	A	502	ISD	O11-C9-C13	3.34	117.16	109.23
4	B	502	ISD	O11-C9-C2	3.77	116.79	110.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	FAD	C5X-C9A-N10	3.90	120.50	117.58
2	A	500	FAD	O2P-P-O1P	3.94	133.07	112.56
4	A	502	ISD	O11-C9-C2	4.12	117.40	110.33
2	A	500	FAD	C4X-N5-C5X	4.39	121.89	116.72
2	A	500	FAD	C1'-N10-C9A	4.53	124.08	118.83
2	B	500	FAD	C4X-N5-C5X	5.05	122.68	116.72
2	A	500	FAD	C5X-C9A-N10	5.08	121.38	117.58
2	B	500	FAD	C4-N3-C2	5.50	119.74	115.16
2	A	500	FAD	C4-N3-C2	9.34	122.95	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	NAD	1	0
4	A	502	ISD	0	1
4	B	502	ISD	1	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	429/451 (95%)	0.18	13 (3%) 54 57	22, 36, 63, 100	0
1	B	425/451 (94%)	0.04	10 (2%) 62 66	23, 35, 61, 99	0
All	All	854/902 (94%)	0.11	23 (2%) 58 61	22, 36, 63, 100	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	337	GLU	6.4
1	B	110	VAL	4.8
1	A	418	VAL	4.0
1	A	426	LYS	3.3
1	B	387	ILE	3.3
1	B	389	ASP	3.3
1	A	424	LEU	3.2
1	B	434	LYS	3.0
1	B	423	GLU	2.8
1	A	417	PRO	2.7
1	B	366	ASP	2.5
1	A	338	SER	2.5
1	B	424	LEU	2.5
1	A	378	ALA	2.4
1	A	389	ASP	2.4
1	A	98	LYS	2.2
1	A	273	VAL	2.2
1	A	422	GLU	2.1
1	B	379	LYS	2.1
1	B	313	VAL	2.1
1	B	422	GLU	2.0
1	A	105	THR	2.0
1	A	307	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	ISD	A	502	12/12	0.92	0.20	5.16	31,38,66,76	0
4	ISD	B	502	12/12	0.89	0.21	3.89	35,46,80,94	0
2	FAD	B	500	53/53	0.96	0.13	0.85	19,25,34,38	0
3	NAD	B	501	44/44	0.97	0.13	0.44	21,28,36,45	0
3	NAD	A	501	44/44	0.98	0.13	-0.12	21,28,38,60	0
2	FAD	A	500	53/53	0.97	0.11	-0.18	19,24,35,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.