



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:57 AM GMT

PDB ID : 2JHT  
Title : CRYSTAL STRUCTURE OF RHOGDI K135T,K138T,K141T MUTANT  
Authors : Cooper, D.R.; Grelewski, K.; Derewenda, Z.S.  
Deposited on : 2007-02-23  
Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6965 (1.90-1.86)
Clashscore	102246	7778 (1.90-1.86)
Ramachandran outliers	100387	7691 (1.90-1.86)
Sidechain outliers	100360	7692 (1.90-1.86)
RSRZ outliers	91569	6979 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	138	<div> <div>6%</div> <div>83%</div> <div>14%</div> <div>..</div> </div>
1	B	138	<div> <div>19%</div> <div>75%</div> <div>20%</div> <div>..</div> </div>
1	C	138	<div> <div>7%</div> <div>75%</div> <div>20%</div> <div>5%</div> </div>
1	D	138	<div> <div>17%</div> <div>77%</div> <div>17%</div> <div>7%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4701 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RHO GDP-DISSOCIATION INHIBITOR 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	138	Total	C	N	O	S	0	1	0
			1112	710	182	214	6			
1	B	138	Total	C	N	O	S	0	1	0
			1109	709	181	213	6			
1	C	138	Total	C	N	O	S	0	0	0
			1104	705	181	213	5			
1	D	138	Total	C	N	O	S	0	1	0
			1108	709	182	212	5			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	135	THR	LYS	ENGINEERED MUTATION	UNP P52565
A	138	THR	LYS	ENGINEERED MUTATION	UNP P52565
A	141	THR	LYS	ENGINEERED MUTATION	UNP P52565
B	135	THR	LYS	ENGINEERED MUTATION	UNP P52565
B	138	THR	LYS	ENGINEERED MUTATION	UNP P52565
B	141	THR	LYS	ENGINEERED MUTATION	UNP P52565
C	135	THR	LYS	ENGINEERED MUTATION	UNP P52565
C	138	THR	LYS	ENGINEERED MUTATION	UNP P52565
C	141	THR	LYS	ENGINEERED MUTATION	UNP P52565
D	135	THR	LYS	ENGINEERED MUTATION	UNP P52565
D	138	THR	LYS	ENGINEERED MUTATION	UNP P52565
D	141	THR	LYS	ENGINEERED MUTATION	UNP P52565

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is LITHIUM ION (three-letter code: LI) (formula: Li).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Li	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	100	Total	O	0	0
			100	100		
4	B	57	Total	O	0	0
			57	57		
4	C	56	Total	O	0	0
			56	56		
4	D	44	Total	O	0	0
			44	44		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: RHO GDP-DISSOCIATION INHIBITOR 1




#### • Molecule 1: RHO GDP-DISSOCIATION INHIBITOR 1



#### • Molecule 1: RHO GDP-DISSOCIATION INHIBITOR 1



#### • Molecule 1: RHO GDP-DISSOCIATION INHIBITOR 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	190.83Å 34.02Å 103.94Å 90.00° 115.59° 90.00°	Depositor
Resolution (Å)	93.66 – 1.88 34.55 – 1.88	Depositor EDS
% Data completeness (in resolution range)	90.4 (93.66-1.88) 90.4 (34.55-1.88)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.53 (at 1.88Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.222 , 0.259 0.220 , 0.260	Depositor DCC
$R_{free}$ test set	1143 reflections (2.62%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.8	Xtriage
Anisotropy	1.203	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 56.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 44845 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4701	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.29 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 7.2295e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, LI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.00	1/1137 (0.1%)	0.99	2/1539 (0.1%)
1	B	0.78	0/1137	0.94	1/1539 (0.1%)
1	C	0.76	1/1129 (0.1%)	0.85	1/1529 (0.1%)
1	D	1.20	4/1133 (0.4%)	0.82	2/1534 (0.1%)
All	All	0.95	6/4536 (0.1%)	0.90	6/6141 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	99	LYS	CE-NZ	22.68	2.05	1.49
1	D	96	SER	CB-OG	19.29	1.67	1.42
1	D	99	LYS	CG-CD	10.08	1.86	1.52
1	A	193	GLU	CB-CG	6.33	1.64	1.52
1	D	99	LYS	CD-CE	5.83	1.65	1.51
1	C	163	GLU	CD-OE2	5.29	1.31	1.25

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	88	LEU	CA-CB-CG	6.58	130.43	115.30
1	D	134	ARG	N-CA-C	-5.86	95.17	111.00
1	A	66	MET	N-CA-CB	5.50	120.51	110.60
1	A	111	ARG	NE-CZ-NH1	-5.31	117.65	120.30
1	D	135	THR	N-CA-C	-5.27	96.76	111.00
1	C	172	ARG	NE-CZ-NH2	-5.22	117.69	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	65	ALA	Peptide
1	B	138	THR	Peptide
1	C	68	PRO	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1112	0	1099	28	0
1	B	1109	0	1100	29	0
1	C	1104	0	1091	34	1
1	D	1108	0	1094	31	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	1	0	0	0	0
4	A	100	0	0	9	0
4	B	57	0	0	4	0
4	C	56	0	0	6	0
4	D	44	0	0	1	0
All	All	4701	0	4384	117	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (117) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:99:LYS:CD	1:D:99:LYS:CG	1.86	1.48
1:D:96:SER:CB	1:D:96:SER:OG	1.67	1.41
1:B:89:ASP:OD1	1:B:91:THR:HG23	1.35	1.27
1:A:159:LEU:HG	4:A:2079:HOH:O	1.38	1.24
1:D:99:LYS:CE	1:D:99:LYS:NZ	2.05	1.18
1:A:159:LEU:CG	4:A:2079:HOH:O	1.94	1.07
1:A:111:ARG:CG	1:A:111:ARG:HH11	1.70	0.99
1:A:111:ARG:NH1	1:A:111:ARG:HG2	1.63	0.95
1:C:134:ARG:CG	1:C:134:ARG:HH11	1.79	0.93
1:A:111:ARG:HH11	1:A:111:ARG:HG2	0.80	0.93
1:B:178:LYS:HE3	1:B:191:SER:HB3	1.55	0.89
1:C:134:ARG:HH11	1:C:134:ARG:HG3	1.38	0.88
1:D:105:LYS:HG3	1:D:201:ASP:HA	1.56	0.85
1:C:69:ASN:HD21	1:C:121:GLU:HG3	1.43	0.84
1:C:134:ARG:NE	4:C:2029:HOH:O	2.11	0.82
1:D:99:LYS:CD	1:D:99:LYS:CB	2.61	0.79
1:D:99:LYS:CG	1:D:99:LYS:CE	2.67	0.72
1:B:178:LYS:CE	1:B:191:SER:HB3	2.19	0.72
1:B:65:ALA:O	1:B:66[B]:MET:CG	2.38	0.71
1:A:145[A]:MET:HG2	4:A:2065:HOH:O	1.91	0.70
1:A:138:THR:HG21	1:C:169:MET:O	1.91	0.70
1:A:159:LEU:CD1	4:A:2079:HOH:O	2.29	0.70
1:D:117:ARG:HD2	1:D:119:ASN:OD1	1.93	0.68
1:D:126:MET:CE	1:D:181:PHE:CE1	2.76	0.68
1:A:202:TRP:O	4:A:2098:HOH:O	2.10	0.67
1:C:134:ARG:CZ	4:C:2029:HOH:O	2.39	0.67
1:B:112:ILE:HD13	1:B:130:GLN:HE22	1.58	0.67
1:D:123:VAL:HG11	1:D:126:MET:SD	2.36	0.66
1:B:109:GLU:OE1	4:B:2014:HOH:O	2.14	0.66
1:A:69:ASN:HD21	1:A:123:VAL:CG1	2.09	0.65
1:A:69:ASN:HD21	1:A:123:VAL:HG13	1.61	0.65
1:A:145[B]:MET:SD	4:A:2052:HOH:O	2.54	0.65
1:C:69:ASN:HD22	1:C:120:ARG:NH1	1.96	0.64
1:B:180:ARG:NH2	4:B:2048:HOH:O	2.30	0.63
1:A:69:ASN:ND2	1:A:123:VAL:CG1	2.63	0.61
1:B:109:GLU:HG2	1:B:162:VAL:CG1	2.31	0.61
1:C:112:ILE:CD1	1:C:130:GLN:HE22	2.14	0.61
1:D:200:LYS:N	1:D:200:LYS:HD2	2.16	0.60
1:C:69:ASN:HD22	1:C:120:ARG:HH12	1.50	0.60
1:C:69:ASN:ND2	1:C:120:ARG:NH1	2.50	0.59
1:A:127:LYS:HD3	1:A:129:ILE:HD11	1.84	0.59
1:C:112:ILE:HD13	1:C:130:GLN:HE22	1.67	0.58

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:169:MET:HB3	1:B:172:ARG:NH1	2.18	0.58
1:B:178:LYS:NZ	4:B:2047:HOH:O	2.37	0.57
1:C:69:ASN:ND2	1:C:121:GLU:HG3	2.16	0.57
1:C:134:ARG:HG2	1:C:134:ARG:HH11	1.67	0.57
1:A:159:LEU:HD12	4:A:2079:HOH:O	1.99	0.57
1:B:65:ALA:O	1:B:66[B]:MET:HG2	2.05	0.56
1:B:81:SER:HB2	1:B:202:TRP:CZ2	2.41	0.56
1:C:174:SER:OG	4:C:2049:HOH:O	2.18	0.56
1:B:66[B]:MET:HG3	1:B:67:VAL:H	1.70	0.56
1:B:131:HIS:ND1	1:B:141:THR:HB	2.20	0.56
1:D:199:LYS:HD3	1:D:200:LYS:HE2	1.88	0.55
1:C:120:ARG:HB2	4:C:2001:HOH:O	2.07	0.55
1:A:138:THR:CG2	1:C:170:LEU:HA	2.36	0.55
1:B:139:ILE:HG23	1:B:139:ILE:O	2.07	0.54
1:D:200:LYS:CD	1:D:200:LYS:H	2.19	0.54
1:B:112:ILE:CD1	1:B:130:GLN:HE22	2.20	0.54
1:D:111:ARG:HD3	1:D:159:LEU:HD22	1.89	0.54
1:D:105:LYS:O	1:D:108:VAL:HG22	2.08	0.54
1:A:138:THR:HG21	1:C:170:LEU:HA	1.90	0.53
1:B:65:ALA:O	1:B:66[B]:MET:HG3	2.08	0.53
1:D:200:LYS:HD2	1:D:200:LYS:H	1.74	0.53
1:D:134:ARG:HH12	1:D:173:GLY:HA3	1.74	0.52
1:C:128:TYR:OH	1:C:130:GLN:NE2	2.38	0.52
1:B:139:ILE:CG2	1:B:139:ILE:O	2.58	0.52
1:B:131:HIS:HE2	1:B:180:ARG:HH11	1.56	0.52
1:A:111:ARG:HD3	4:A:2010:HOH:O	2.10	0.51
1:D:201:ASP:OD2	1:D:201:ASP:N	2.42	0.51
1:B:131:HIS:HB3	1:B:139:ILE:HD11	1.92	0.51
1:B:67:VAL:HG23	4:B:2019:HOH:O	2.11	0.51
1:C:129:ILE:HG12	1:C:143:ASP:HB3	1.93	0.51
1:D:109:GLU:HG2	1:D:162:VAL:HG12	1.93	0.50
1:C:134:ARG:CG	1:C:134:ARG:NH1	2.52	0.50
1:B:81:SER:HB2	1:B:202:TRP:HZ2	1.75	0.50
1:A:65:ALA:N	4:A:2001:HOH:O	2.43	0.50
1:C:134:ARG:NH1	1:C:134:ARG:HG3	2.13	0.50
1:D:109:GLU:HG2	1:D:162:VAL:CG1	2.42	0.49
1:B:89:ASP:OD1	1:B:91:THR:CG2	2.30	0.48
1:C:109:GLU:HG2	1:C:162:VAL:CG1	2.44	0.48
1:C:76:THR:HG23	1:C:87:GLU:HG2	1.96	0.48
1:C:66:MET:C	1:C:68:PRO:HD3	2.33	0.47
1:A:112:ILE:HD13	1:A:130:GLN:HE22	1.79	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:106:GLU:HB2	1:C:167:LYS:HB3	1.96	0.47
1:A:122:ILE:HG12	1:A:150:GLY:HA2	1.96	0.47
1:A:138:THR:CG2	1:C:169:MET:O	2.60	0.47
1:B:109:GLU:HG3	1:B:164:GLU:HG2	1.96	0.47
1:C:120:ARG:HD2	1:C:121:GLU:HG2	1.96	0.46
1:B:135:THR:OG1	1:D:135:THR:HG23	2.16	0.45
1:D:96:SER:CB	1:D:96:SER:HG	2.10	0.45
1:D:78:VAL:HB	1:D:111:ARG:HB3	1.99	0.45
1:D:126:MET:HE3	1:D:181:PHE:CE1	2.52	0.44
1:A:76:THR:HG23	1:A:87:GLU:HG2	1.99	0.44
1:C:66:MET:O	1:C:68:PRO:HD3	2.18	0.44
1:D:66:MET:HB3	1:D:67:VAL:H	1.50	0.44
1:C:154:GLU:HB2	4:C:2039:HOH:O	2.17	0.44
1:B:66[B]:MET:HG3	1:B:67:VAL:N	2.31	0.43
1:B:112:ILE:HD13	1:B:130:GLN:NE2	2.30	0.43
1:D:180:ARG:HG3	4:D:2019:HOH:O	2.18	0.43
1:D:82:ALA:HA	1:D:83:PRO:HD2	1.81	0.43
1:A:111:ARG:NH2	1:A:159:LEU:HD13	2.34	0.43
1:D:134:ARG:HG2	1:D:135:THR:CG2	2.48	0.43
1:B:117:ARG:HD3	1:B:155:GLU:HA	2.00	0.42
1:C:166:PRO:HG2	1:C:171:ALA:HB1	2.01	0.42
1:C:68:PRO:HA	1:C:120:ARG:NH2	2.34	0.42
1:A:134:ARG:HD3	1:A:175:TYR:CE2	2.54	0.42
1:C:123:VAL:HG21	1:C:126:MET:HE1	2.01	0.42
1:C:70:VAL:CG2	1:C:123:VAL:HG11	2.50	0.42
1:C:77:LEU:HD12	1:C:102:PHE:CD2	2.55	0.42
1:D:96:SER:OG	1:D:100:GLN:NE2	2.54	0.41
1:D:126:MET:HE1	1:D:181:PHE:CE1	2.54	0.41
1:B:137:VAL:O	1:B:137:VAL:HG22	2.19	0.41
1:A:134:ARG:HG3	4:C:2029:HOH:O	2.19	0.41
1:D:134:ARG:HG2	1:D:135:THR:HG22	2.02	0.41
1:A:184:ASP:N	1:A:184:ASP:OD1	2.42	0.41
1:D:134:ARG:HD2	1:D:175:TYR:CE2	2.56	0.41
1:A:123:VAL:O	1:A:123:VAL:HG23	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:91:THR:O	1:C:119:ASN:O[2_554]	1.91	0.29

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	137/138 (99%)	132 (96%)	4 (3%)	1 (1%)	26	13
1	B	137/138 (99%)	129 (94%)	3 (2%)	5 (4%)	4	0
1	C	136/138 (99%)	131 (96%)	5 (4%)	0	100	100
1	D	137/138 (99%)	132 (96%)	4 (3%)	1 (1%)	26	13
All	All	547/552 (99%)	524 (96%)	16 (3%)	7 (1%)	17	4

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	66[A]	MET
1	B	66[B]	MET
1	B	139	ILE
1	D	66	MET
1	B	136	GLY
1	B	137	VAL
1	A	66	MET

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/123 (101%)	120 (97%)	4 (3%)	46	33
1	B	124/123 (101%)	107 (86%)	17 (14%)	4	1
1	C	123/123 (100%)	115 (94%)	8 (6%)	21	9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	D	122/123 (99%)	113 (93%)	9 (7%)	17 6
All	All	493/492 (100%)	455 (92%)	38 (8%)	17 5

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	111	ARG
1	A	143	ASP
1	A	170	LEU
1	A	184	ASP
1	B	66[A]	MET
1	B	66[B]	MET
1	B	67	VAL
1	B	81	SER
1	B	97	PHE
1	B	99	LYS
1	B	126	MET
1	B	134	ARG
1	B	137	VAL
1	B	138	THR
1	B	139	ILE
1	B	141	THR
1	B	169	MET
1	B	170	LEU
1	B	181	PHE
1	B	193	GLU
1	B	200	LYS
1	C	69	ASN
1	C	77	LEU
1	C	96	SER
1	C	109	GLU
1	C	121	GLU
1	C	134	ARG
1	C	167	LYS
1	C	186	LYS
1	D	87	GLU
1	D	101	SER
1	D	109	GLU
1	D	117	ARG
1	D	127	LYS
1	D	167	LYS
1	D	193	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	200	LYS
1	D	201	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	ASN
1	A	130	GLN
1	B	130	GLN
1	C	69	ASN
1	C	130	GLN
1	D	100	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	1203	-	4,4,4	0.48	0	6,6,6	0.76	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	B	1203	-	4,4,4	0.36	0	6,6,6	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1203	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1203	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	138/138 (100%)	0.81	8 (5%) 26 28	25, 32, 47, 52	0
1	B	138/138 (100%)	1.38	26 (18%) 2 2	28, 43, 66, 74	1 (0%)
1	C	138/138 (100%)	1.04	9 (6%) 22 24	37, 45, 56, 77	0
1	D	138/138 (100%)	1.23	23 (16%) 2 2	41, 59, 69, 78	0
All	All	552/552 (100%)	1.11	66 (11%) 6 6	25, 44, 67, 78	1 (0%)

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	65	ALA	19.5
1	C	67	VAL	14.0
1	C	66	MET	11.9
1	C	68	PRO	7.7
1	B	66[A]	MET	5.7
1	B	138	THR	5.3
1	B	93	ASP	5.2
1	A	65	ALA	4.9
1	B	174	SER	4.7
1	D	65	ALA	4.7
1	D	67	VAL	4.7
1	B	83	PRO	4.6
1	D	184	ASP	4.4
1	D	66	MET	4.3
1	B	92	GLY	4.2
1	B	94	LEU	4.1
1	B	137	VAL	3.8
1	B	65	ALA	3.8
1	B	99	LYS	3.6
1	B	135	THR	3.5
1	B	98	LYS	3.4

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	97	PHE	3.2
1	B	79	CYS	3.1
1	C	83	PRO	3.1
1	B	67	VAL	3.1
1	A	122	ILE	2.9
1	B	91	THR	2.9
1	B	139	ILE	2.9
1	D	85	PRO	2.9
1	C	187	THR	2.8
1	D	159	LEU	2.8
1	B	166	PRO	2.8
1	D	94	LEU	2.7
1	C	154	GLU	2.7
1	A	124	SER	2.7
1	A	79	CYS	2.7
1	B	145	MET	2.7
1	D	185	ASP	2.6
1	B	200	LYS	2.5
1	D	154	GLU	2.5
1	D	88	LEU	2.5
1	D	183	ASP	2.5
1	D	102	PHE	2.4
1	B	133	TYR	2.4
1	C	164	GLU	2.4
1	D	78	VAL	2.4
1	B	167	LYS	2.4
1	D	196	LEU	2.4
1	D	103	VAL	2.4
1	D	198	ILE	2.3
1	D	108	VAL	2.3
1	D	173	GLY	2.3
1	B	193	GLU	2.3
1	B	184	ASP	2.3
1	A	120	ARG	2.2
1	A	118	VAL	2.2
1	D	192	TRP	2.2
1	B	181	PHE	2.2
1	D	193	GLU	2.1
1	D	187	THR	2.1
1	B	144	TYR	2.1
1	C	78	VAL	2.1
1	A	184	ASP	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	199	LYS	2.0
1	D	201	ASP	2.1
1	A	146	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	LI	A	1204	1/1	0.97	0.62	-	16,16,16,16	0
2	SO4	B	1203	5/5	0.88	0.32	-	56,58,61,62	0
2	SO4	A	1203	5/5	0.95	0.29	-	39,44,46,46	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.