



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 26, 2016 – 08:58 PM BST

PDB ID : 2JNW
Title : Solution structure of a ERCC1-XPA heterodimer
Authors : Tsodikov, O.V.; Ivanov, D.; Orelli, B.; Staresincic, L.; Scharer, O.D.; Wagner, G.
Deposited on : 2007-02-07

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	133	
2	B	14	

2 Ensemble composition and analysis ⓘ

This entry contains 10 models. Model 7 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:99-A:142, A:150-A:214, B:67-B:69 (112)	0.09	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 1 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 5, 7, 8, 9, 10
Single-model clusters	6

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2045 atoms, of which 1035 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called DNA excision repair protein ERCC-1.

Mol	Chain	Residues	Atoms						Trace
1	A	116	Total	C	H	N	O	S	0
			1878	601	947	162	164	4	

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	82	MET	-	EXPRESSION TAG	UNP P07992
A	83	GLY	-	EXPRESSION TAG	UNP P07992
A	84	SER	-	EXPRESSION TAG	UNP P07992
A	85	SER	-	EXPRESSION TAG	UNP P07992
A	86	HIS	-	EXPRESSION TAG	UNP P07992
A	87	HIS	-	EXPRESSION TAG	UNP P07992
A	88	HIS	-	EXPRESSION TAG	UNP P07992
A	89	HIS	-	EXPRESSION TAG	UNP P07992
A	90	HIS	-	EXPRESSION TAG	UNP P07992
A	91	HIS	-	EXPRESSION TAG	UNP P07992
A	92	SER	-	EXPRESSION TAG	UNP P07992
A	93	GLN	-	EXPRESSION TAG	UNP P07992
A	94	ASP	-	EXPRESSION TAG	UNP P07992
A	95	PRO	-	EXPRESSION TAG	UNP P07992

- Molecule 2 is a protein called DNA-repair protein complementing XP-A cells.


Mol	Chain	Residues	Atoms					Trace
2	B	11	Total	C	H	N	O	0
			167	53	88	12	14	

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA excision repair protein ERCC-1




- Molecule 2: DNA-repair protein complementing XP-A cells



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 7. Colouring as in section 4.1 above.

- Molecule 1: DNA excision repair protein ERCC-1



- Molecule 2: DNA-repair protein complementing XP-A cells



K67	I68	D70	I71	G72	G73	F74	F75	I76	L77	GLU	GLU	GLU
I69	D70	I71	G72	G73	F74	F75	I76	L77	GLU	GLU	GLU	GLU

5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	refinement	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.17±0.00	0±0/883 (0.0±0.0%)	1.36±0.00	2±0/1198 (0.2±0.0%)
2	B	1.04±0.08	0±0/25 (0.0±0.0%)	1.15±0.07	0±0/33 (0.0±0.0%)
All	All	1.17	0/9080 (0.0%)	1.36	20/12310 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	3.8±0.4
All	All	0	38

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	100	SER	N-CA-C	-6.60	93.18	111.00	1	10
1	A	156	ARG	NE-CZ-NH1	5.09	122.84	120.30	1	10

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	108	ARG	Sidechain	10
1	A	117	ARG	Sidechain	10
1	A	167	ARG	Sidechain	10
1	A	106	ARG	Sidechain	8

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	864	883	883	77±4
2	B	25	37	34	6±3
All	All	8890	9200	9170	829

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 46.

5 of 154 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:208:TYR:CZ	1:A:212:TYR:HE2	1.23	1.52	7	10
1:A:208:TYR:CZ	1:A:212:TYR:CE2	1.19	2.30	7	10
1:A:208:TYR:OH	1:A:212:TYR:CE2	0.95	2.19	7	10
1:A:168:VAL:HG11	1:A:208:TYR:OH	0.93	1.64	7	10
1:A:208:TYR:CE1	1:A:212:TYR:CE2	0.90	2.59	7	10

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	107/133 (80%)	92±1 (86±1%)	11±1 (10±1%)	5±1 (4±1%)	6	30
2	B	2/14 (14%)	0±0 (20±24%)	1±1 (65±39%)	0±0 (15±23%)	1	5
All	All	1090/1470 (74%)	919 (84%)	120 (11%)	51 (5%)	5	28

5 of 8 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	108	ARG	10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	101	ILE	10
1	A	110	ASN	7
1	A	150	PRO	7
1	A	109	GLY	7

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	95/117 (81%)	85±1 (90±1%)	10±1 (10±1%)	14	58
2	B	3/11 (27%)	1±1 (47±22%)	2±1 (53±22%)	0	1
All	All	980/1280 (77%)	868 (89%)	112 (11%)	11	54

5 of 18 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	211	THR	10
1	A	166	LEU	10
1	A	209	LEU	10
1	A	180	GLN	10
1	A	212	TYR	10

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided