



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 21, 2016 – 12:18 PM EDT

PDB ID : 5JNL
Title : Structure of Plasmodium falciparum DXR in complex with a beta-substituted fosmidomycin analogue, LC54 and manganese
Authors : Sooriyaarachchi, S.; Bergfors, T.; Jones, T.A.; Mowbray, S.L.
Deposited on : 2016-04-30
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION



The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2475 (1.60-1.60)
Clashscore	102246	2732 (1.60-1.60)
Ramachandran outliers	100387	2654 (1.60-1.60)
Sidechain outliers	100360	2653 (1.60-1.60)
RSRZ outliers	91569	2479 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	
1	B	422	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	A	507	-	-	-	X
5	EDO	B	506	-	-	-	X
5	EDO	B	508	-	-	-	X
5	EDO	B	510	-	-	-	X

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 7269 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 1-deoxy-D-xylulose 5-phosphate reductoisomerase, apicoplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	410	3328	2132	545	630	21	0	8	0
1	B	410	3321	2129	543	629	20	0	8	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	67	MET	-	initiating methionine	UNP Q8IKG4
A	68	ALA	-	expression tag	UNP Q8IKG4
A	69	HIS	-	expression tag	UNP Q8IKG4
A	70	HIS	-	expression tag	UNP Q8IKG4
A	71	HIS	-	expression tag	UNP Q8IKG4
A	72	HIS	-	expression tag	UNP Q8IKG4
A	73	HIS	-	expression tag	UNP Q8IKG4
A	74	HIS	-	expression tag	UNP Q8IKG4
B	67	MET	-	initiating methionine	UNP Q8IKG4
B	68	ALA	-	expression tag	UNP Q8IKG4
B	69	HIS	-	expression tag	UNP Q8IKG4
B	70	HIS	-	expression tag	UNP Q8IKG4
B	71	HIS	-	expression tag	UNP Q8IKG4
B	72	HIS	-	expression tag	UNP Q8IKG4
B	73	HIS	-	expression tag	UNP Q8IKG4
B	74	HIS	-	expression tag	UNP Q8IKG4

- Molecule 2 is {(2R)-2-{2-[hydroxy(methyl)amino]-2-oxoethyl}-5-[4-(trifluoromethyl)phenyl]pentyl}phosphonic acid (three-letter code: L54) (formula: C₁₅H₂₁F₃NO₅P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf		
			Total	C	F	N	O			P	
2	A	1	Total	25	15	3	1	5	1	0	0
2	B	1	Total	25	15	3	1	5	1	0	0

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mn		
3	B	1	Total	1	0	0
3	A	1	Total	1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	234	Total O 234 234	0	0
6	B	274	Total O 274 274	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 1-deoxy-D-xylulose 5-phosphate reductoisomerase, apicoplactic



- Molecule 1: 1-deoxy-D-xylulose 5-phosphate reductoisomerase, apicoplactic



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	51.47Å 56.03Å 86.04Å 103.10° 102.89° 100.90°	Depositor
Resolution (Å)	30.00 – 1.60 47.88 – 1.60	Depositor EDS
% Data completeness (in resolution range)	94.3 (30.00-1.60) 85.5 (47.88-1.60)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.178 , 0.199 0.187 , 0.203	Depositor DCC
R_{free} test set	5526 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	18.6	Xtrriage
Anisotropy	0.250	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.008 for -h,-k,h+k+l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7269	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MN, L54, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/3394	0.58	0/4583
1	B	0.37	0/3393	0.59	1/4583 (0.0%)
All	All	0.37	0/6787	0.58	1/9166 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	98	ARG	NE-CZ-NH1	5.14	122.87	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3328	0	3363	5	0
1	B	3321	0	3363	4	0
2	A	25	0	0	0	0
2	B	25	0	0	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	12	0	16	2	0
4	B	12	0	16	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	12	0	18	0	0
5	B	24	0	36	0	0
6	A	234	0	0	1	0
6	B	274	0	0	1	0
All	All	7269	0	6812	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:464[A]:MET:HE3	1:A:464[A]:MET:O	1.89	0.72
4:A:504:GOL:O2	6:A:601:HOH:O	2.09	0.60
1:B:339:ILE:HG22	1:B:355:MET:HG2	1.94	0.50
1:B:359:ASP:N	2:B:501:L54:F25	2.36	0.48
1:B:114:VAL:HG22	1:B:121:LEU:HD22	1.98	0.46
1:A:340:ILE:HD12	1:A:391:PHE:HZ	1.80	0.46
1:A:191:MET:HG3	1:A:218:LEU:HD11	1.99	0.45
1:A:339:ILE:HG22	1:A:355:MET:HG2	2.01	0.42
4:A:504:GOL:H11	6:B:602:HOH:O	2.20	0.42
1:B:340:ILE:HD12	1:B:391:PHE:HZ	1.85	0.41
1:A:340:ILE:HG12	1:A:355:MET:HG3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	416/422 (99%)	404 (97%)	11 (3%)	1 (0%)	52 28
1	B	416/422 (99%)	405 (97%)	11 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	832/844 (99%)	809 (97%)	22 (3%)	1 (0%)	56 31

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	293	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	385/388 (99%)	378 (98%)	7 (2%)	66 41
1	B	385/388 (99%)	382 (99%)	3 (1%)	86 75
All	All	770/776 (99%)	760 (99%)	10 (1%)	76 56

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	116	LYS
1	A	155	LYS
1	A	160	CYS
1	A	292	LYS
1	A	297	LYS
1	A	349	LYS
1	A	469	GLN
1	B	297	LYS
1	B	349	LYS
1	B	360	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	459	ASN
1	B	197	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 2 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	L54	A	501	3	24,25,25	1.05	3 (12%)	28,36,36	1.13	1 (3%)
4	GOL	A	503	-	5,5,5	0.27	0	5,5,5	0.56	0
4	GOL	A	504	-	5,5,5	0.34	0	5,5,5	0.55	0
5	EDO	A	505	-	3,3,3	0.47	0	2,2,2	0.22	0
5	EDO	A	506	-	3,3,3	0.54	0	2,2,2	0.25	0
5	EDO	A	507	-	3,3,3	0.43	0	2,2,2	0.30	0
2	L54	B	501	3	24,25,25	1.13	4 (16%)	28,36,36	1.12	1 (3%)
4	GOL	B	503	-	5,5,5	0.29	0	5,5,5	0.29	0
4	GOL	B	504	-	5,5,5	0.26	0	5,5,5	0.30	0
5	EDO	B	505	-	3,3,3	0.41	0	2,2,2	0.37	0
5	EDO	B	506	-	3,3,3	0.42	0	2,2,2	0.27	0
5	EDO	B	507	-	3,3,3	0.46	0	2,2,2	0.21	0
5	EDO	B	508	-	3,3,3	0.53	0	2,2,2	0.20	0
5	EDO	B	509	-	3,3,3	0.43	0	2,2,2	0.38	0
5	EDO	B	510	-	3,3,3	0.45	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	L54	A	501	3	-	0/24/25/25	0/1/1/1
4	GOL	A	503	-	-	0/4/4/4	0/0/0/0
4	GOL	A	504	-	-	0/4/4/4	0/0/0/0
5	EDO	A	505	-	-	0/1/1/1	0/0/0/0
5	EDO	A	506	-	-	0/1/1/1	0/0/0/0
5	EDO	A	507	-	-	0/1/1/1	0/0/0/0
2	L54	B	501	3	-	0/24/25/25	0/1/1/1
4	GOL	B	503	-	-	0/4/4/4	0/0/0/0
4	GOL	B	504	-	-	0/4/4/4	0/0/0/0
5	EDO	B	505	-	-	0/1/1/1	0/0/0/0
5	EDO	B	506	-	-	0/1/1/1	0/0/0/0
5	EDO	B	507	-	-	0/1/1/1	0/0/0/0
5	EDO	B	508	-	-	0/1/1/1	0/0/0/0
5	EDO	B	509	-	-	0/1/1/1	0/0/0/0
5	EDO	B	510	-	-	0/1/1/1	0/0/0/0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	L54	P5-O6	-2.47	1.48	1.54
2	B	501	L54	P5-O7	-2.07	1.49	1.54
2	A	501	L54	P5-O6	-2.06	1.49	1.54
2	A	501	L54	C11-N9	2.12	1.48	1.44
2	B	501	L54	C11-N9	2.16	1.48	1.44
2	B	501	L54	P5-C1	2.70	1.81	1.78
2	A	501	L54	P5-C1	2.77	1.81	1.78

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	L54	C3-C4-N9	2.69	120.81	117.59
2	A	501	L54	C3-C4-N9	2.89	121.05	117.59

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	504	GOL	2	0
2	B	501	L54	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	410/422 (97%)	0.26	30 (7%) 18 16	11, 21, 46, 100	0
1	B	410/422 (97%)	-0.01	11 (2%) 58 56	10, 18, 35, 96	0
All	All	820/844 (97%)	0.12	41 (5%) 32 29	10, 19, 44, 100	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	296	TRP	16.0
1	B	296	TRP	15.2
1	B	184	PHE	8.1
1	A	293	HIS	7.4
1	A	295	LYS	7.1
1	B	183	SER	6.7
1	B	297	LYS	6.3
1	B	294	PRO	5.8
1	A	294	PRO	5.5
1	B	293	HIS	5.3
1	A	183	SER	5.1
1	B	295	LYS	4.8
1	A	141	TYR	4.6
1	A	153	ASP	4.4
1	A	116	LYS	4.3
1	A	160	CYS	4.2
1	A	150	ASN	3.9
1	A	126	ARG	3.7
1	A	122	TYR	3.7
1	A	148	VAL	3.4
1	A	297	LYS	3.3
1	A	154	TYR	3.3
1	A	157	ILE	3.1
1	A	138	LYS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	149	LYS	2.8
1	A	144	LEU	2.8
1	A	155	LYS	2.8
1	B	185	GLN	2.8
1	A	142	GLU	2.8
1	A	184	PHE	2.7
1	A	484	LYS	2.7
1	A	152	LYS	2.7
1	B	485	HIS	2.6
1	B	153	ASP	2.6
1	A	158	ILE	2.5
1	A	145	LYS	2.5
1	B	188	TYR	2.1
1	A	486	ASN	2.1
1	A	151	ILE	2.1
1	A	136	HIS	2.1
1	A	135	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	EDO	A	507	4/4	0.80	0.17	2.69	36,37,38,40	0
5	EDO	B	510	4/4	0.92	0.15	2.57	42,42,42,43	0
5	EDO	B	508	4/4	0.78	0.16	2.52	26,31,33,36	0
5	EDO	B	506	4/4	0.91	0.13	2.32	24,24,24,26	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	GOL	A	504	6/6	0.88	0.14	1.99	24,28,30,30	0
4	GOL	B	504	6/6	0.75	0.16	1.94	30,33,35,37	0
5	EDO	A	506	4/4	0.89	0.11	1.21	28,29,29,30	0
5	EDO	B	507	4/4	0.83	0.14	0.79	46,46,46,47	0
5	EDO	B	509	4/4	0.87	0.11	0.78	34,35,36,37	0
4	GOL	A	503	6/6	0.87	0.11	0.57	27,29,29,30	0
5	EDO	B	505	4/4	0.92	0.09	0.45	25,27,27,27	0
2	L54	B	501	25/25	0.93	0.14	0.20	16,28,42,44	0
2	L54	A	501	25/25	0.93	0.14	-0.01	22,31,54,55	0
5	EDO	A	505	4/4	0.76	0.10	-0.92	50,51,51,51	0
4	GOL	B	503	6/6	0.93	0.09	-1.18	21,22,23,23	0
3	MN	B	502	1/1	1.00	0.05	-2.00	13,13,13,13	0
3	MN	A	502	1/1	1.00	0.02	-3.24	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.