



# Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 09:11 PM BST

PDB ID : 2JOL  
Title : Average NMR structure of the catalytic domain of guanine nucleotide exchange factor BopE from Burkholderia pseudomallei  
Authors : Wu, H.; Upadhyay, A.; Williams, C.; Galyov, E.E.; van den Elsen, J.M.H.; Bagby, S.  
Deposited on : 2007-03-14

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	186	 69% 23% 6% ..

## 2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2775 atoms, of which 1370 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Putative G-nucleotide exchange factor.

Mol	Chain	Residues	Atoms						Trace
1	A	184	Total	C	H	N	O	S	0
			2775	862	1370	263	276	4	


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	76	GLY	-	CLONING ARTIFACT	UNP Q63K41
A	77	SER	-	CLONING ARTIFACT	UNP Q63K41

## 4 Residue-property plots [i](#)

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Putative G-nucleotide exchange factor



## 5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *simulated annealing*.

Of the 40 calculated structures, 1 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	
X-PLOR NIH	refinement	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality

### 6.1 Standard geometry

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1405	1370	1369	42
All	All	1405	1370	1369	42

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:251:GLU:O	1:A:252:LYS:O	0.73	2.06
1:A:85:ILE:HD13	1:A:85:ILE:O	0.69	1.88
1:A:189:MET:SD	1:A:194:ALA:HB2	0.65	2.32
1:A:108:ALA:O	1:A:112:ARG:NH1	0.60	2.34
1:A:251:GLU:O	1:A:256:ASN:ND2	0.58	2.36
1:A:174:THR:O	1:A:175:ASN:CB	0.57	2.53
1:A:81:ALA:O	1:A:85:ILE:HG22	0.56	2.01
1:A:112:ARG:CD	1:A:112:ARG:N	0.55	2.70
1:A:177:LEU:O	1:A:181:ILE:HG23	0.54	2.02
1:A:101:THR:N	1:A:102:PRO:CD	0.53	2.71
1:A:129:ARG:NE	1:A:203:GLN:OE1	0.53	2.42
1:A:175:ASN:N	1:A:176:PRO:CD	0.53	2.72
1:A:107:ASP:O	1:A:109:GLU:N	0.52	2.42
1:A:254:ALA:O	1:A:258:ALA:N	0.51	2.44
1:A:189:MET:C	1:A:191:ALA:H	0.50	2.09

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:201:GLU:O	1:A:203:GLN:N	0.50	2.40
1:A:129:ARG:NE	1:A:203:GLN:CD	0.50	2.65
1:A:189:MET:O	1:A:191:ALA:N	0.50	2.45
1:A:177:LEU:HD12	1:A:177:LEU:H	0.50	1.66
1:A:80:ASP:C	1:A:82:LYS:N	0.48	2.67
1:A:83:GLN:O	1:A:87:HIS:CD2	0.48	2.67
1:A:140:ASN:C	1:A:141:VAL:HG22	0.47	2.30
1:A:86:ARG:HB2	1:A:202:LEU:HD21	0.46	1.88
1:A:193:LEU:N	1:A:193:LEU:HD12	0.46	2.24
1:A:153:ALA:O	1:A:156:THR:OG1	0.45	2.35
1:A:248:SER:O	1:A:252:LYS:N	0.45	2.49
1:A:189:MET:C	1:A:191:ALA:N	0.45	2.70
1:A:193:LEU:N	1:A:193:LEU:CD1	0.44	2.80
1:A:207:ARG:O	1:A:211:ILE:CG1	0.44	2.66
1:A:131:CYS:SG	1:A:142:ARG:NE	0.43	2.91
1:A:201:GLU:C	1:A:203:GLN:H	0.43	2.17
1:A:198:GLU:OE1	1:A:198:GLU:CA	0.43	2.66
1:A:205:TYR:C	1:A:205:TYR:CD1	0.43	2.91
1:A:129:ARG:O	1:A:133:ASP:OD2	0.43	2.37
1:A:80:ASP:O	1:A:82:LYS:N	0.43	2.51
1:A:131:CYS:SG	1:A:142:ARG:CZ	0.42	3.07
1:A:203:GLN:N	1:A:204:PRO:CD	0.42	2.82
1:A:79:GLY:O	1:A:80:ASP:CG	0.42	2.58
1:A:254:ALA:O	1:A:255:THR:C	0.42	2.58
1:A:204:PRO:CB	1:A:230:ARG:NH1	0.41	2.83
1:A:251:GLU:O	1:A:252:LYS:C	0.41	2.58
1:A:260:GLY:O	1:A:261:ALA:C	0.41	2.59

## 6.3 Torsion angles ⓘ

### 6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	182/186 (98%)	160 (88%)	12 (7%)	10 (5%)	4	24
All	All	182/186 (98%)	160 (88%)	12 (7%)	10 (5%)	4	24



All 10 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	254	ALA
1	A	108	ALA
1	A	80	ASP
1	A	175	ASN
1	A	202	LEU
1	A	221	GLU
1	A	84	ALA
1	A	252	LYS
1	A	141	VAL
1	A	190	GLY

### 6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	137/138 (99%)	123 (90%)	14 (10%)	14	58
All	All	137/138 (99%)	123 (90%)	14 (10%)	14	58

All 14 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	184	ARG
1	A	256	ASN
1	A	89	VAL
1	A	198	GLU
1	A	235	THR
1	A	175	ASN
1	A	112	ARG
1	A	252	LYS
1	A	85	ILE
1	A	141	VAL
1	A	189	MET
1	A	110	PHE
1	A	101	THR
1	A	181	ILE

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided