



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 09:27 PM BST

PDB ID : 2JSY
Title : Solution structure of Tpx in the oxidized state
Authors : Jin, C.; Lu, J.
Deposited on : 2007-07-17

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 86%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	167	 89% 5% 6%

2 Ensemble composition and analysis

This entry contains 21 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:3-A:84, A:93-A:167 (157)	0.38	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 6 single-model clusters were found.

Cluster number	Models
1	1, 3, 5, 7, 9, 12, 13, 15, 18, 19
2	4, 10, 16
3	14, 21
Single-model clusters	2; 6; 8; 11; 17; 20

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 2557 atoms, of which 1276 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Probable thiol peroxidase.


Mol	Chain	Residues	Atoms						Trace
1	A	167	Total	C	H	N	O	S	0
			2557	810	1276	213	253	5	

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Probable thiol peroxidase




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: Probable thiol peroxidase




4.2.2 Score per residue for model 2

- Molecule 1: Probable thiol peroxidase




4.2.3 Score per residue for model 3

- Molecule 1: Probable thiol peroxidase




4.2.4 Score per residue for model 4

- Molecule 1: Probable thiol peroxidase




4.2.5 Score per residue for model 5

- Molecule 1: Probable thiol peroxidase




4.2.6 Score per residue for model 6

- Molecule 1: Probable thiol peroxidase




4.2.7 Score per residue for model 7

- Molecule 1: Probable thiol peroxidase




4.2.8 Score per residue for model 8

- Molecule 1: Probable thiol peroxidase




4.2.9 Score per residue for model 9

- Molecule 1: Probable thiol peroxidase




4.2.10 Score per residue for model 10

- Molecule 1: Probable thiol peroxidase




4.2.11 Score per residue for model 11

- Molecule 1: Probable thiol peroxidase




4.2.12 Score per residue for model 12

- Molecule 1: Probable thiol peroxidase




4.2.13 Score per residue for model 13

- Molecule 1: Probable thiol peroxidase




4.2.14 Score per residue for model 14

- Molecule 1: Probable thiol peroxidase




4.2.15 Score per residue for model 15

- Molecule 1: Probable thiol peroxidase




4.2.16 Score per residue for model 16

- Molecule 1: Probable thiol peroxidase




4.2.17 Score per residue for model 17

- Molecule 1: Probable thiol peroxidase




4.2.18 Score per residue for model 18

- Molecule 1: Probable thiol peroxidase




4.2.19 Score per residue for model 19

- Molecule 1: Probable thiol peroxidase




4.2.20 Score per residue for model 20

- Molecule 1: Probable thiol peroxidase



4.2.21 Score per residue for model 21

- Molecule 1: Probable thiol peroxidase



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 21 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.0
AMBER	refinement	7

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 15798
Number of chemical shift lists	1
Total number of shifts	1895
Number of shifts mapped to atoms	1895
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	86%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.61±0.02	0±0/1223 (0.0±0.0%)	0.95±0.11	2±3/1655 (0.1±0.2%)
All	All	0.61	0/25683 (0.0%)	0.96	36/34755 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.4±1.1
All	All	0	9

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	109	ARG	NE-CZ-NH2	-11.65	114.47	120.30	1	4
1	A	65	ARG	NE-CZ-NH2	-11.57	114.52	120.30	1	8
1	A	129	ARG	NE-CZ-NH2	-9.54	115.53	120.30	1	2
1	A	129	ARG	NE-CZ-NH1	9.00	124.80	120.30	1	1
1	A	65	ARG	NE-CZ-NH1	7.74	124.17	120.30	1	1
1	A	125	ARG	NE-CZ-NH2	-7.50	116.55	120.30	1	1
1	A	120	TYR	CB-CG-CD2	-7.33	116.60	121.00	1	6
1	A	66	ARG	NE-CZ-NH2	-7.21	116.70	120.30	1	7
1	A	93	TRP	CA-CB-CG	5.72	124.56	113.70	1	1
1	A	109	ARG	CA-CB-CG	-5.62	101.04	113.40	1	1
1	A	27	PHE	CB-CG-CD2	-5.61	116.87	120.80	1	1
1	A	137	ASN	CA-CB-CG	5.24	124.92	113.40	1	1
1	A	111	MET	CG-SD-CE	5.10	108.36	100.20	1	1
1	A	142	TYR	CB-CG-CD1	-5.09	117.95	121.00	1	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	145	TYR	Sidechain	3
1	A	129	ARG	Sidechain	2
1	A	66	ARG	Sidechain	1
1	A	27	PHE	Sidechain	1
1	A	16	GLN	Peptide	1
1	A	17	GLU	Peptide	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1204	1198	1198	0±1
All	All	25284	25158	25158	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:30:LEU:CD2	1:A:109:ARG:HD3	0.58	2.28	1	1
1:A:27:PHE:HB2	1:A:112:SER:HB3	0.51	1.82	1	1
1:A:30:LEU:HD22	1:A:109:ARG:HD3	0.44	1.88	1	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	156/167 (93%)	142±3 (91±2%)	12±3 (7±2%)	2±2 (1±1%)	19	64
All	All	3276/3507 (93%)	2985 (91%)	245 (7%)	46 (1%)	19	64

All 13 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	26	ASP	9
1	A	16	GLN	8
1	A	23	GLN	8
1	A	17	GLU	6
1	A	93	TRP	4
1	A	96	ALA	2
1	A	14	VAL	2
1	A	63	GLN	2
1	A	100	ASP	1
1	A	64	THR	1
1	A	75	GLY	1
1	A	94	CYS	1
1	A	55	ILE	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/138 (95%)	122±3 (93±2%)	9±3 (7±2%)	24	69
All	All	2751/2898 (95%)	2561 (93%)	190 (7%)	24	69

All 52 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	140	VAL	20
1	A	99	ILE	12
1	A	79	VAL	12
1	A	151	ASN	8
1	A	137	ASN	8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	57	THR	7
1	A	111	MET	7
1	A	129	ARG	6
1	A	131	VAL	6
1	A	150	THR	6
1	A	64	THR	5
1	A	93	TRP	5
1	A	120	TYR	5
1	A	104	THR	5
1	A	66	ARG	4
1	A	28	THR	4
1	A	29	VAL	4
1	A	94	CYS	3
1	A	50	SER	3
1	A	13	LEU	3
1	A	19	LYS	3
1	A	63	GLN	3
1	A	65	ARG	3
1	A	112	SER	3
1	A	55	ILE	3
1	A	97	ASN	3
1	A	135	ASP	3
1	A	124	LEU	2
1	A	38	SER	2
1	A	20	VAL	2
1	A	126	LEU	2
1	A	60	CYS	2
1	A	156	GLU	2
1	A	16	GLN	2
1	A	108	HIS	2
1	A	54	SER	2
1	A	106	SER	2
1	A	32	ASN	2
1	A	152	HIS	1
1	A	141	VAL	1
1	A	17	GLU	1
1	A	101	LYS	1
1	A	39	LEU	1
1	A	109	ARG	1
1	A	107	ASP	1
1	A	127	LEU	1
1	A	46	VAL	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	18	VAL	1
1	A	23	GLN	1
1	A	130	SER	1
1	A	68	ASN	1
1	A	3	GLU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 86% for the well-defined parts and 85% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 15798

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1895
Number of shifts mapped to atoms	1895
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	165	-0.16 ± 0.16	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	153	0.07 ± 0.17	None needed (< 0.5 ppm)
$^{13}\text{C}'$	142	0.09 ± 0.13	None needed (< 0.5 ppm)
^{15}N	145	0.86 ± 0.36	Should be applied

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 86%, i.e. 1598 atoms were assigned a chemical shift out of a possible 1850. 29 out of 29 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	739/775 (95%)	302/309 (98%)	296/314 (94%)	141/152 (93%)
Sidechain	817/955 (86%)	496/551 (90%)	313/367 (85%)	8/37 (22%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	42/120 (35%)	41/64 (64%)	0/53 (0%)	1/3 (33%)
Overall	1598/1850 (86%)	839/924 (91%)	609/734 (83%)	150/192 (78%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 85%, i.e. 1677 atoms were assigned a chemical shift out of a possible 1974. 30 out of 30 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	770/823 (94%)	318/328 (97%)	307/334 (92%)	145/161 (90%)
Sidechain	865/1022 (85%)	526/591 (89%)	330/390 (85%)	9/41 (22%)
Aromatic	42/129 (33%)	41/69 (59%)	0/57 (0%)	1/3 (33%)
Overall	1677/1974 (85%)	885/988 (90%)	637/781 (82%)	155/205 (76%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

